

Sex Differences in Estrogen Receptor α and β Levels and Activation Status in LPS-Stimulated Human Macrophages

Questa è la versione Post print del seguente articolo:

Original

Sex Differences in Estrogen Receptor α and β Levels and Activation Status in LPS-Stimulated Human Macrophages / Campesi, Ilaria; Marino, Maria; Montella, Andrea Costantino Mario; Pais, Sara; Franconi, Flavia. - In: JOURNAL OF CELLULAR PHYSIOLOGY. - ISSN 0021-9541. - 232:2(2017), pp. 340-345. [10.1002/jcp.25425]

Availability:

This version is available at: 11388/173560 since: 2022-06-01T09:41:20Z

Publisher:

Published

DOI:10.1002/jcp.25425

Terms of use:

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

Publisher copyright

note finali coverpage

(Article begins on next page)

Sex differences in estrogen receptor α and β levels and activation status in LPS-stimulated human macrophages

Sex-dependent LPS modulation of estrogen receptors

Ilaria Campesi ^{1,2*}, Maria Marino ^{3*}, Andrea Montella ¹, Sara Pais ¹, Flavia Franconi ^{1,4}

¹ Department of Biomedical Sciences, University of Sassari, Sassari, Italy

² Laboratory of Sex-Gender Medicine, National Institute of Biostructures and Biosystems, Osilo, Italy

³ Cell Physiology Lab, Department of Science, University Roma Tre, Rome, Italy

⁴ Assessorato alle Politiche per la Persona of Basilicata Region, Potenza, Italy;

***Co-corresponding authors:**

Ilaria Campesi, Department of Biomedical Science, Via Muroni 23, 07100 Sassari, Italy.

Phone (+39) 079-228709 E-mail: ilacampesi79@yahoo.it

Maria Marino, Department of Science, University Roma Tre, V.le G. Marconi, 446, 00146

Rome, Italy Phone (+39) 06-57336345 E-mail: maria.marino@uniroma3.it

Email addresses

Ilaria Campesi : ilacampesi79@yahoo.it

Maria Marino: maria.marino@uniroma3.it

Andrea Montella: montella@uniss.it

Sara Pais: sara.pais@hotmail.it

Flavia Franconi: franconi@iol.it

Abstract

Immune function, inflammation and atherosclerosis display sex differences and are influenced by 17 β -estradiol through estrogen receptors subtypes ER α and ER β . Male tissues express active ERs, but their possible involvement in inflammation in males has never been assessed. Macrophages express both ER α and ER β and offer the opportunity to evaluate the role of ER levels and activation in inflammation. We assessed the ability of lipopolysaccharide (LPS) to modulate, in a sex-specific way, the expression and the activation status of ER α and ER β in blood monocytes-derived macrophages (MDMs) from men and women. MDMs were incubated with 100 ng / ml LPS for 24 h and used to evaluate ER α , ER β , P- ER α , p38 and P-p38 expression by Western Blotting.

In basal conditions, ER α and ER β were significantly higher in female MDMs than in male MDMs. LPS up-regulated ER α and ER α phosphorylation in both sexes, with a significantly higher effect observed in male MDMs, and down-regulated ER β level only in female MDMs. p38 and P-p38 proteins, indicative of ER β activity, did not show sex differences both in basal conditions and after LPS treatment. Finally, ER α / ER β and P-ER α / ER α ratios were significantly higher in male MDMs than in female ones.

Our data indicate, for the first time, that LPS affects ER α but not ER β activation status. We identify a significant role of ER α in LPS-mediated inflammatory responses in MDMs, which represents an initial step in understanding the influence of sex in the relationship between LPS and ER α .

Keywords: estrogen receptors, lipopolysaccharide, inflammation, sex differences

Number of Figures: 3

Introduction

Macrophages play a significant role in immunity and immune responses. These cells reside in every tissue of the body and are activated in the initial steps of inflammation (Fujiwara and Kobayashi, 2005) and atherosclerosis (Moore et al., 2013; Moore and Tabas, 2011). **A large body of clinical and preclinical evidence has demonstrated that** 17 β -estradiol (E2), the most active of estrogen hormones, plays a key role **in both inflammation and atherosclerosis** (Arnal et al., 2004; Lizcano and Guzman, 2014). In particular, the menstrual cycle and pregnancy influence the etiology and course of chronic inflammatory diseases and the increase in the incidence of disorders characterized by a strong inflammatory component (e.g., osteoporosis, atherosclerosis, diabetes, and arthritis) is typical in post-menopausal women, **whereas** pre-menopausal women are less prone than males to these diseases (Villa et al., 2015). Furthermore, E2 reduces the production of pro-inflammatory cytokines including interleukin-6, tumor necrosis factor- α (TNF- α) and macrophage inhibitory factor (Hsieh et al., 2007; Messingham et al., 2001; Suzuki et al., 2008).

All E2 anti-inflammatory effects are mediated by both estrogen receptors subtypes (ER α and ER β), members of nuclear receptor family, which are widely distributed in several **parts** of male and female reproductive system as well as in various non-reproductive tissues such as brain, bone and the cardiovascular system (Ascenzi et al., 2006). ER-mediated cellular responses to E2 have been loosely grouped into two interconnected categories: nuclear and rapid extra-nuclear signals. It is now known that both ER subtypes share the same molecular action mechanisms which are strictly dependent on the intracellular localization of ERs (Ascenzi et al., 2006). In addition, it has been reported that ERs have a key role in inflammation, as the down-regulation of their expression has been reported in mouse microglia (Sierra et al., 2008), and the activation of ER α led to an anti-inflammatory effect in lipopolysaccharide (LPS) -induced brain inflammation (Vegeto et al., 2003). It is widely accepted that abnormalities in ER levels and signaling lead to different types of pathological conditions. For example, ER α expression increases at early stages of cancer where acts as a tumor promoter, **whereas**, ER β levels are reduced during carcinogenesis and cancer progression acting as a tumor suppressor (Jia et al., 2015). At the present, the contribution of the balance of ER α and ER β levels and activation status to E2 anti-inflammatory effects is completely unknown.

Macrophages represent a good experimental model to evaluate the role played by ER levels and activation in inflammation in that these cells express both ER α and ER β (Jensen and Pioli, 2008; Murphy et al., 2009). Furthermore, ER α levels are higher in monocytes derived macrophages (MDMs) from women taking oral contraceptives versus non-users, **whereas** ER β is more expressed and is also more active in untreated women (Campesi et al., 2012). ERs play, also, a pivotal role in monocyte survival regulation (Vegeto et al., 2001) and in inflammation in different cells systems (Cuzzocrea et al., 2000). In particular, ERs activity has been shown to augment and dampen innate immune signaling pathways in macrophages (Cunningham and Gilkeson, 2011; Cutolo et al., 1995).

Several sex differences have been reported in immune function, inflammation and atherosclerosis (Berkley et al., 2006; Imahara et al., 2005; Lansky et al., 2012; Ngo et al., 2014; Oertelt-Prigione, 2012). As example, the LPS-induced TNF- α release (the most central inflammatory mediator rapidly secreted after interaction with pathogens) is different in men and women (Campesi et al., 2013). Moreover, normal human female alveolar macrophages **are able to kill the ingested pneumococci better than** men and authors suggest a role for E2 (Yang et al., 2014).

Altogether, these data sustain the ability of E2 to modulate ER levels in macrophages and the anti-inflammatory role of this hormone in females. However, no information is available on the possible involvement of ERs in inflammation in male, although male tissues express active ERs [(Ascenzi et al., 2006) and literature cited therein]. Therefore, we aimed to assess the ability of LPS, a bacterial cell-wall component of Gram-negative and a component of tobacco (Sebastian et al., 2006) to modulate, in a sex-specific way, the expression and the activity of ER α and ER β in MDMs from young adult, and healthy men and women.

Materials and methods

Population

The study was approved by the local ethical committee of ASL1 of Sassari. Informed consent was obtained, and blood sampling was performed during a voluntary blood donation. Seven healthy adult men and 7 healthy adult women aged between 21-35 years were enrolled. All women were fertile and premenopausal, with regular menstrual cycles (28-30 days) without reported precocious and /or surgically-induced menopause, were free from contraceptives use for at least 3 months to ensure an adequate wash-out period and were all analysed

during the follicular phase of their menstrual cycles. All procedures were conducted in accordance with the Declaration of Helsinki.

Monocytes isolation, differentiation and treatment

Monocytes of individual subjects were isolated from 30 ml of blood withdrawn from healthy men and women according to Campesi et al. (2012). Purified monocytes were obtained by adhesion; non-adherent cells (mainly lymphocytes) were removed by gentle washes with phosphate buffer. MDMs were prepared from monocytes cultured for 8 days in a 5% CO₂ incubator at 37°C in RPMI 1640 medium containing 20% fetal bovine serum, 2 mM glutamine, 10 mM HEPES and antibiotics / antimycotics. This procedure allowed to obtain an homogeneous population of MDMs, which appeared as adherent cells with their typical elongated morphology: prominent nucleus with flatly outspread cytoplasm and multiple pseudopodia. After 8 day of culture, MDMs were incubated with 100 ng / ml LPS for 24 h and subsequently lysed for western blotting analysis.

Western blotting

MDMs cells were cultured for 8 days and then were lysed (Cell Lysis Buffer, Cell Signalling Technology, Danvers, MA, USA). The protein concentration was quantified using the BCA protein assay kit (Thermo Scientific, Waltham, MA, USA). For the Western blot analysis, 30 µg of solubilized proteins were electrophoretically resolved by 4% – 15% SDS-PAGE (100 V, 2 h, 24°C) and then transferred to a PVDF membrane (250 mA, 65 min, 4°C) using a mini-PROTEAN tetracell system (Bio-Rad, Milano, Italy). The membranes were blocked in 5% (w/v) skim milk (Sigma-Aldrich, Milano, Italy) in 150 mM Tris buffer (Sigma-Aldrich, Milano, Italy) and 20 mM Tris-HCl, pH 7.2 (Sigma-Aldrich, Milano, Italy,) at 24°C for 1 h. Then membranes were incubated overnight at 4°C with the following antibodies: rabbit ER α and mouse ER β (Santa Cruz Biotechnology, Segrate, Milano Italy) (1:100), rabbit P-ER α (Ser118), rabbit p38, and rabbit P-p38 (Cell Signalling Technology, Danvers, MA, USA) (1 : 1,000). After P-ER α (Ser118) and P-p38 determination, the nitrocellulose membranes were stripped by Restore Western Blot Stripping Buffer (Pierce Chemical Company, Rockford, IL, USA) for 10 min at room temperature and then probed with anti-ER α and anti-p38 antibodies, respectively. The level of phosphorylated / unphosphorylated protein has been assayed on the same electrophoresis, but the relative figures have been splitted to better evidence the

differences between male and female samples. The rabbit polyclonal antibody anti- β -tubulin (Cell Signalling Technology, Danvers, MA, USA) (1:1,000) was used as loading control to normalize ER levels; no difference in β -tubulin levels were reported between sexes as demonstrated by Western blot images. After washing, the blots were incubated for 1 h with anti-rabbit IgG and anti-mouse IgG, horseradish peroxidase (HRP)-conjugated secondary antibodies (Cell Signalling Technology, Milano, Italy) (1:2,000). Antibody binding was detected using a chemiluminescence reaction (Cell Signalling Technology, Danvers, MA, USA) with the Bio-Rad Chemi Doc instrument (Berkeley, CA, USA). Band volume analysis was performed using the Image Lab 4.0 software (Bio-Rad Laboratories, Berkeley, CA, USA).

Statistical analysis

Data were reported as the mean \pm SD. Statistical analysis was performed by Two Way Analysis of Variance (**2-Way ANOVA**) followed by the Pairwise Multiple Comparison Procedures to analyse the effect of sex and treatment with LPS, using Sigma-Stat 3.1 software (Systat Software, Erkrath, Germany). For the analysis of P-ER α , which was undetectable in basal conditions, an unpaired Student's t –tests was used to compare males and females after LPS exposure. The distribution of samples was assessed by the Kolmogorov-Smirnov and Shapiro tests. A $P \leq 0.05$ was considered statistically significant.

Results

Population

The 2 cohorts were well-matched for age and body mass index, and, as expected, men were taller and weight more than females.

LPS effect on ER α levels in male and female MDMs

ER α has a key role in the E2 anti-inflammatory effects (Sierra et al., 2008; Vegeto et al., 2003), this prompted us to first evaluate the effect of LPS on the levels and activation status of this ER subtype in male and female MDMs. The **2-Way ANOVA** demonstrated that there is a statistically significant interaction between sex and treatment ($P < 0.001$). In particular, under basal conditions, the expression of ER α was significantly different in MDMs obtained from adult healthy men and adult healthy women (**Fig. 1**). Indeed, ER α expression was 2 times higher in MDMs obtained from women in comparison with MDMs obtained from men

(**Fig. 1A**). LPS exposure (100 ng / ml, 24 h) up-regulated ER α level by 26 times in male and 9 times in female MDMs (**Fig. 1A**).

It is well recognized that ER α activities (both nuclear and extra-nuclear) should be preceded by receptor phosphorylation at 118 serine residue (ER α S118) (Ali et al., 1993; La Rosa et al., 2012). In unstimulated cells, P-ER α S118 levels were undetectable in MDMs of both sexes, indicating that the receptor is inactive. However, LPS exposure (100 ng / ml, 24 h) up-regulated ER α phosphorylation in both sexes even if the effect was significantly higher in male MDMs than in female ones (**Fig. 1B**).

LPS effect on ER β levels in male and female MDM cells.

The **2-Way ANOVA** demonstrated that there is a statistically significant interaction between sex and treatment ($P < 0.001$). In basal conditions, ER β expression appeared four times higher in female MDMs than in male cells sustaining the possible role played by this receptor subtype in E2 anti-inflammatory effects (**Fig. 2A**). **In contrast** to the effect on ER α , LPS exposure (100 ng / ml, 24 h) did not affect ER β expression in male MDMs, whereas it significantly down-regulated ER β level in female MDMs (**Fig. 2A**).

We previously identified the E2-dependent activation of the p38 component of the MAPK family as a conserved pathway in ER β -based E2 signaling (Galluzzo et al., 2007). The ER β -mediated activation of the p38/MAPK, which occurs in several E2 target cells (Acconcia et al., 2005; Galluzzo et al., 2007), is important both for the regulation of ER expression and for the transcriptional activity of this receptor subtype (Caiazza et al., 2007). LPS treatment significantly induced the levels of p38, without sex differences, although this compound did not affect the phosphorylation status of this kinase and thus its activation status (**Figs. 2B and 2C**). Consequently, no significant variation in the level of P-p38 / p38 ratio were reported (**Fig. 2D**), indicating that the level of activation of ER β was not different between sexes after LPS exposure (**Figs. 2B, 2C, and 2D**).

LPS effect on ER α /ER β ratio in male and female MDM cells.

The differences above reported between male and female ERs levels prompted us to calculate the ratio between receptor subtypes. **Figure 3A** shows that the ER α / ER β ratio was significantly higher in male MDMs than in female ones. Intriguingly, the activation status of ER α (i.e P-ER α / ER α ratio) was significantly higher in male MDMs (**Fig. 3B**).

Discussion

A plethora of literature have demonstrated that ER α and ER β exert opposite effects on E2-regulated cellular proliferation; these data have evidenced the strict correlation existing between the perturbation of ER subtype-specific expression and different clinical outcomes in cancer patients (Ascenzi et al., 2006; Kim and Kim, 2014; Malorni et al., 2008). Less information is available on the contribution of the balance of ER α and ER β levels and their activation status to the well-known E2 anti-inflammatory effects. In addition, none of this information has ever evaluated the possible involvement of ERs in inflammation in male, although male tissues express active ERs (Ascenzi et al., 2006). Here, we assessed the ability of LPS to modulate, in a sex-specific way, the expression of ER α and ER β in MDMs from young adult, and healthy men and women. Moreover, when activated, both ERs trigger different and complex signal transduction pathways, thus, we selected ER α Ser118 phosphorylation and ER β -dependent p38 activation as markers of receptor activation (Acconcia et al., 2005; Ali et al., 1993; Galluzzo et al., 2007; La Rosa et al., 2012). These two parameters represent a new tool in the study of ER biology being involved and, thus, allowing to investigate both nuclear (transcriptional) and extra-nuclear (rapid activation of signaling cascade) receptor activities.

Results reported here show that 24 h LPS exposure modifies the expression of the ER α and ER β , the activity of the ER α in MDMs and these changes are sex-dependent. In particular, LPS significantly up-regulates the ER α level and this up-regulation is more evident in male than in female cells. On the other hand, the ER β level is down-regulated only in female MDMs. The above data also indicate that LPS strongly perturbs the ER α / ER β balance inducing a net increase of ER α level in both sexes. Considering that ER α can be activated in a ligand-dependent manner, via E2, or in a ligand-independent manner, via several signal transduction pathways including p44/42 MAPK (McGlynn et al., 2013) and PI3K/AKT (Ascenzi et al., 2006) it is likely that LPS activates ER α in a ligand independent way. Interestingly, human MDMs maintain sex differences after 8 days in culture. This result is in line with previous data that demonstrated that freshly isolated rat female- and male-derived vascular smooth muscle cells maintain several sex differences at least for the first eight passages in culture (Malorni et al., 2008; Pellegrini et al., 2014; Straface et al., 2009).

It has been reported that LPS binds to the Toll like receptor 4 (TLR4) (Takeda and Akira, 2005) which, in turn, enhances ER α expression whereas neither E2, nor ER α , impacted TLR4 expression, at least, in mesangial cells (Svenson et al., 2014). In contrast, Smith et al. (2011) demonstrated that LPS decreases the levels of ER α mRNA in microglia without any effect on ER β mRNA, and Holm et al. (2010) reported that 4 days of LPS stimulation reduced both ER α and ER β mRNA levels in endothelial cells. Finally, Toniolo et al. (2015) found that the association of LPS with interferon gamma decreases ER α levels. Although it is possible that LPS effects differ in dependence of animal species or tissue, data reported in literature and present data suggest that LPS effects could rely on the modulation of ER synthesis and degradation. Moreover, the evidence that the number of TLR4 is higher in blood monocytes isolated from boys than in girls (Bannister et al., 2013) well supports the higher LPS effect in male than in female MDMs.

The most important result of this research is the LPS ability to increase ER α phosphorylation without any effect on ER β activation status. The ability of TLR ligands, including LPS, to increase ER α phosphorylation has been already reported in mesangial cells where it is associated to the increase of inflammatory response (Svenson et al., 2014). Moreover, LPS induces higher expression of TLR4 and greater release of TNF- α in male neutrophils than in female ones (Aomatsu et al., 2013), which is paralleled by a stronger activation of mitogen-activated protein kinases and phosphatidylinositol 3-kinase (Aomatsu et al., 2013). Together these data sustain the pivotal role played by ligand-independent ER α activation in LPS-induced inflammatory response and suggest that males are prone to the inflammation cascade as females.

Conclusions

Our findings sustain the key role played by ER α in LPS-induced inflammatory response and indicate that males are more sensitive to the LPS effects than females. Considering the role of macrophages in atherosclerosis the higher response of male MDMs respect to female ones can impact on a variety of key molecular mechanisms linked with important processes such as immune response and cardiovascular diseases (Moore et al., 2013; Moore and Tabas, 2011).

As a whole, the results reported here identify a significant impact / interaction of ER α in LPS-mediated inflammatory responses in MDMs representing an initial step in understanding the influence of sex in the relationship between LPS, its receptor and ER α .

Abbreviations

E2: 17 β -estradiol; ER α : estrogen receptor alpha; ER β : estrogen receptor beta; IL-6: interleukin 6; LPS: lipopolysaccharide; MDMs: monocytes-derived macrophages; P-ER α : phosphorylated estrogen receptor alpha; P-p38: phosphorylated p38; TLR4: Toll like receptor 4; TNF- α : tumor necrosis factor- α

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We thank the Mayor of Osilo for the granting of the structure devoted to the National Laboratory of Gender Medicine.

References

- Acconcia F, Totta P, Ogawa S, Cardillo I, Inoue S, Leone S, Trentalance A, Muramatsu M, Marino M. 2005. Survival versus apoptotic 17 β -estradiol effect: role of ER alpha and ER beta activated non-genomic signaling. *J Cell Physiol* 203(1):193-201.
- Ali S, Metzger D, Bornert JM, Chambon P. 1993. Modulation of transcriptional activation by ligand-dependent phosphorylation of the human oestrogen receptor A/B region. *Embo J* 12(3):1153-1160.
- Aomatsu M, Kato T, Kasahara E, Kitagawa S. 2013. Gender difference in tumor necrosis factor-alpha production in human neutrophils stimulated by lipopolysaccharide and interferon-gamma. *Biochem Biophys Res Commun* 441(1):220-225.
- Arnal JF, Gourdy P, Elhage R, Garmy-Susini B, Delmas E, Brouchet L, Castano C, Barreira Y, Couloumiers JC, Prats H, Prats AC, Bayard F. 2004. Estrogens and atherosclerosis. *Eur J Endocrinol* 150(2):113-117.
- Ascenzi P, Bocedi A, Marino M. 2006. Structure-function relationship of estrogen receptor alpha and beta: impact on human health. *Mol Aspects Med* 27(4):299-402.
- Bannister EG, Smith C, Visvanathan K, Thompson A, Hardikar W. 2013. TLR2 and TLR4 in healthy children: age and gender differences. *J Paediatr Child Health* 49(12):1082-1083.
- Berkley KJ, Zalcman SS, Simon VR. 2006. Sex and gender differences in pain and inflammation: a rapidly maturing field. *Am J Physiol Regul Integr Comp Physiol* 291(2):R241-244.

- Caiazza F, Galluzzo P, Lorenzetti S, Marino M. 2007. 17Beta-estradiol induces ERbeta up-regulation via p38/MAPK activation in colon cancer cells. *Biochem Biophys Res Commun* 359(1):102-107.
- Campesi I, Carru C, Zinellu A, Occhioni S, Sanna M, Palermo M, Tonolo G, Mercurio G, Franconi F. 2013. Regular cigarette smoking influences the transsulfuration pathway, endothelial function, and inflammation biomarkers in a sex-gender specific manner in healthy young humans. *Am J Transl Res* 5(5):497-509.
- Campesi I, Sanna M, Zinellu A, Carru C, Rubattu L, Bulzomi P, Seghieri G, Tonolo G, Palermo M, Rosano G, Marino M, Franconi F. 2012. Oral contraceptives modify DNA methylation and monocyte-derived macrophage function. *Biol Sex Differ* 3:4.
- Cunningham M, Gilkeson G. 2011. Estrogen receptors in immunity and autoimmunity. *Clin Rev Allergy Immunol* 40(1):66-73.
- Cutolo M, Sulli A, Seriola B, Accardo S, Masi AT. 1995. Estrogens, the immune response and autoimmunity. *Clin Exp Rheumatol* 13(2):217-226.
- Cuzzocrea S, Santagati S, Sautebin L, Mazzon E, Calabro G, Serraino I, Caputi AP, Maggi A. 2000. 17beta-estradiol antiinflammatory activity in carrageenan-induced pleurisy. *Endocrinology* 141(4):1455-1463.
- Fujiwara N, Kobayashi K. 2005. Macrophages in inflammation. *Curr Drug Targets Inflamm Allergy* 4(3):281-286.
- Galluzzo P, Caiazza F, Moreno S, Marino M. 2007. Role of ERbeta palmitoylation in the inhibition of human colon cancer cell proliferation. *Endocr Relat Cancer* 14(1):153-167.
- Holm A, Andersson KE, Nordstrom I, Hellstrand P, Nilsson BO. 2010. Down-regulation of endothelial cell estrogen receptor expression by the inflammation promoter LPS. *Mol Cell Endocrinol* 319(1-2):8-13.
- Hsieh YC, Frink M, Hsieh CH, Choudhry MA, Schwacha MG, Bland KI, Chaudry IH. 2007. Downregulation of migration inhibitory factor is critical for estrogen-mediated attenuation of lung tissue damage following trauma-hemorrhage. *Am J Physiol Lung Cell Mol Physiol* 292(5):L1227-1232.
- Imahara SD, Jelacic S, Junker CE, O'Keefe GE. 2005. The influence of gender on human innate immunity. *Surgery* 138(2):275-282.
- Jensen AL, Pioli PA. 2008. Estrogen induces the expression of estrogen receptor in human macrophages. *FASEB J* 22:672.643.
- Jia M, Dahlman-Wright K, Gustafsson JA. 2015. Estrogen receptor alpha and beta in health and disease. *Best Pract Res Clin Endocrinol Metab* 29(4):557-568.
- Kim HM, Kim HS. 2014. [Gender-specific colorectal cancer: epidemiologic difference and role of estrogen]. *Korean J Gastroenterol* 63(4):201-208.
- La Rosa P, Pesiri V, Leclercq G, Marino M, Acconcia F. 2012. Palmitoylation regulates 17beta-estradiol-induced estrogen receptor-alpha degradation and transcriptional activity. *Mol Endocrinol* 26(5):762-774.
- Lansky AJ, Ng VG, Maehara A, Weisz G, Lerman A, Mintz GS, De Bruyne B, Farhat N, Niess G, Jankovic I, Lazar D, Xu K, Fahy M, Serruys PW, Stone GW. 2012. Gender and the extent of coronary atherosclerosis, plaque composition, and clinical outcomes in acute coronary syndromes. *JACC Cardiovasc Imaging* 5(3 Suppl):S62-72.
- Lizcano F, Guzman G. 2014. Estrogen Deficiency and the Origin of Obesity during Menopause. *Biomed Res Int* 2014:757461.

- Malorni W, Straface E, Matarrese P, Ascione B, Coinu R, Canu S, Galluzzo P, Marino M, Franconi F. 2008. Redox state and gender differences in vascular smooth muscle cells. *FEBS Lett* 582(5):635-642.
- McGlynn LM, Tovey S, Bartlett JM, Doughty J, Cooke TG, Edwards J. 2013. Interactions between MAP kinase and oestrogen receptor in human breast cancer. *Eur J Cancer* 49(6):1176-1186.
- Messingham KA, Heinrich SA, Kovacs EJ. 2001. Estrogen restores cellular immunity in injured male mice via suppression of interleukin-6 production. *J Leukoc Biol* 70(6):887-895.
- Moore KJ, Sheedy FJ, Fisher EA. 2013. Macrophages in atherosclerosis: a dynamic balance. *Nat Rev Immunol* 13(10):709-721.
- Moore KJ, Tabas I. 2011. Macrophages in the pathogenesis of atherosclerosis. *Cell* 145(3):341-355.
- Murphy AJ, Guyre PM, Wira CR, Pioli PA. 2009. Estradiol regulates expression of estrogen receptor ERalpha46 in human macrophages. *PLoS One* 4(5):e5539.
- Ngo ST, Steyn FJ, McCombe PA. 2014. Gender differences in autoimmune disease. *Front Neuroendocrinol* 35(3):347-369.
- Oertelt-Prigione S. 2012. The influence of sex and gender on the immune response. *Autoimmun Rev* 11(6-7):A479-485.
- Pellegrini M, Bulzomi P, Lecis M, Leone S, Campesi I, Franconi F, Marino M. 2014. Endocrine disruptors differently influence estrogen receptor beta and androgen receptor in male and female rat VSMC. *J Cell Physiol* 229(8):1061-1068.
- Sebastian A, Pehrson C, Larsson L. 2006. Elevated concentrations of endotoxin in indoor air due to cigarette smoking. *J Environ Monit* 8(5):519-522.
- Sierra A, Gottfried-Blackmore A, Milner TA, McEwen BS, Bulloch K. 2008. Steroid hormone receptor expression and function in microglia. *Glia* 56(6):659-674.
- Smith JA, Das A, Butler JT, Ray SK, Banik NL. 2011. Estrogen or estrogen receptor agonist inhibits lipopolysaccharide induced microglial activation and death. *Neurochem Res* 36(9):1587-1593.
- Straface E, Vona R, Gambardella L, Ascione B, Marino M, Bulzomi P, Canu S, Coinu R, Rosano G, Malorni W, Franconi F. 2009. Cell sex determines anoikis resistance in vascular smooth muscle cells. *FEBS Lett* 583(21):3448-3454.
- Suzuki T, Yu HP, Hsieh YC, Choudhry MA, Bland KI, Chaudry IH. 2008. Estrogen-mediated activation of non-genomic pathway improves macrophages cytokine production following trauma-hemorrhage. *J Cell Physiol* 214(3):662-672.
- Svenson J, Cunningham M, Dasgupta S, Gilkeson GS. 2014. Estrogen receptor alpha modulates mesangial cell responses to toll-like receptor ligands. *Am J Med Sci* 348(6):492-500.
- Takeda K, Akira S. 2005. Toll-like receptors in innate immunity. *Int Immunol* 17(1):1-14.
- Toniolo A, Fadini GP, Tedesco S, Cappellari R, Vegeto E, Maggi A, Avogaro A, Bolego C, Cignarella A. 2015. Alternative activation of human macrophages is rescued by estrogen treatment in vitro and impaired by menopausal status. *J Clin Endocrinol Metab* 100(1):E50-58.
- Vegeto E, Belcredito S, Eteri S, Ghisletti S, Brusadelli A, Meda C, Krust A, Dupont S, Ciana P, Chambon P, Maggi A. 2003. Estrogen receptor-alpha mediates the brain antiinflammatory activity of estradiol. *Proc Natl Acad Sci U S A* 100(16):9614-9619.

- Vegeto E, Bonincontro C, Pollio G, Sala A, Viappiani S, Nardi F, Brusadelli A, Viviani B, Ciana P, Maggi A. 2001. Estrogen prevents the lipopolysaccharide-induced inflammatory response in microglia. *J Neurosci* 21(6):1809-1818.
- Villa A, Rizzi N, Vegeto E, Ciana P, Maggi A. 2015. Estrogen accelerates the resolution of inflammation in macrophagic cells. *Sci Rep* 5:15224.
- Yang Z, Huang YC, Koziel H, de Crom R, Ruetten H, Wohlfart P, Thomsen RW, Kahlert JA, Sorensen HT, Jozefowski S, Colby A, Kobzik L. 2014. Female resistance to pneumonia identifies lung macrophage nitric oxide synthase-3 as a therapeutic target. *Elife* 3.

Figures legends

Fig. 1. Representative Western blot and densitometric analysis of ER α (**A**) and P-ER α (**B**) expression in male (black bar) and female (white bar) MDMs in basal conditions and after 24h exposure to 100 ng / ml LPS. The values are expressed as the means \pm SD of at least 4 independent experiments for each sex, normalised on β -tubulin levels. The level of phosphorylated / unphosphorylated protein has been assayed on the same electrophoresis, but the relative figures have been splitted to better evidence the differences between male and female samples. Connectors represent statistical difference between basal condition and LPS stimulation; * and ** indicate a $P < 0.05$ and a $P < 0.001$ between male and female MDMs, respectively.

Fig. 2. Representative Western blot and densitometric analysis of ER β (**A**), p38 (**B**), P-p38 (**C**) expression and P-p38 / p38 ratio (**D**) in male (black bar) and female (white bar) MDMs in basal conditions and after 24h exposure to 100 ng / ml LPS. The values are expressed as the means \pm SD of at least 4 independent experiments for each sex, normalised on β -tubulin levels. The level of phosphorylated / unphosphorylated proteins has been assayed on the same electrophoresis, but the relative figures have been splitted to better evidence the differences between male and female samples. Connectors represent statistical difference between basal condition and LPS stimulation; * indicates a $P < 0.05$ between male and female MDMs.

Fig. 3. ER α / ER β (**A**) and P-ER α / ER α (**B**) ratios calculated in in male (black bar) and female (white bar) MDMs in basal conditions and after 24h exposure to 100 ng / ml LPS. The values are expressed as the means \pm SD of at least 4 independent experiments for each sex. Connectors represent statistical difference between basal condition and LPS stimulation; * and ** indicate a P < 0.05 and a P < 0.001 between male and female MDMs, respectively.