

Post-IR IRSL290 dating of K-rich feldspar sand grains in a wind-dominated system on Sardinia

Questa è la versione Post print del seguente articolo:

*Original*

Post-IR IRSL290 dating of K-rich feldspar sand grains in a wind-dominated system on Sardinia / Andreucci, S., Sechi, D., Buylaert, J.P., Sanna, L., Pascucci, V.. - In: MARINE AND PETROLEUM GEOLOGY. - ISSN 0264-8172. - 87:(2017), pp. 91-98. [10.1016/j.marpetgeo.2017.03.025]

*Availability:*

This version is available at: 11388/175000 since: 2022-06-11T11:19:42Z

*Publisher:*

*Published*

DOI:10.1016/j.marpetgeo.2017.03.025

*Terms of use:*

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

*Publisher copyright*

note finali coverpage

(Article begins on next page)

# Post-IR IRSL<sub>290</sub> dating of K-feldspar in a wind-dominated system on Sardinia

S. Andreucci<sup>1</sup>, D. Sechi<sup>1</sup>, J.-P. Buylaert<sup>2,3</sup>, L. Sanna<sup>4</sup>, V. Pascucci<sup>5</sup>

1. Department of Chemical and Geological Sciences, University of Cagliari, Cagliari, Italy

2. Nordic Laboratory for Luminescence dating, Department of Earth Sciences, University of Aarhus, Risø DTU, Roskilde, Denmark

3. Radiation Research Division, Risø DTU, Roskilde, Denmark

4. IBIMET-CNR, Sassari, Italy

5. Department of Architecture, Design and Planning, University of Sassari, Sassari, Italy

## ABSTRACT

The reliability of a post-IR elevated temperature IRSL (290°C; pIRIR<sub>290</sub>) is tested on wind-blown, sand-sized (180-250 μm) K-rich feldspar grains. The pIRIR<sub>290</sub> dates were compared with quartz SAR-OSL data, other independent age controls and historical information.

Three study areas along the coast of Sardinia (Italy) were selected: The Bue Marino cave (NE Sardinia), south Alghero coast and the Alghero bay (NW Sardinia).

The sedimentary succession at Bue Marino cave includes a sandy wind-blown unit sandwiched between two calcareous crusts. U-series dates of these crusts constrain the aeolianite formation between ~125 and ~65 ka. Quartz SAR-OSL samples were saturated (> 2xD<sub>0</sub>) instead, fading-uncorrected pIRIR<sub>290</sub> ages on K-feldspar extracts point to a formation between ~100 and ~80 ka, in good agreement with the U-series data.

Along the south Alghero coast a thick dunefield system is widely recognised in the literature to represent the beginning of the last glacial phase (*post* 80 ka). From a single block sand-sized grains for quartz SAR-OSL and K-feldspars post-IR IRSL-OSL purposes were collected. Quartz SAR-OSL sample lies below the saturation limit (< 2xD<sub>0</sub>) giving a reliable age of 76 ± 6 ka. The fading-uncorrected post-IR IRSL<sub>290</sub> age of 73 ± 5 ka is in good agreement with the quartz result.

Samples from a modern coastal cordon system backing Alghero bay was collected for quartz SAR-OSL and K-feldspars post-IR IRSL-OSL purposes. This dunefield was stabilized by plantation during the 1950s. The quartz SAR-OSL ages span from 2500±150 years to 60±20 years ago, consistent with the known cordon stabilization. The pIRIR<sub>290</sub> ages indicate an age offset of ~1500 years.

We can conclude that the pIRIR<sub>290</sub> method on sand-sized K-feldspar grains shows great promise for samples at or beyond the quartz OSL age limit but should not be applied to Late Holocene or modern deposits.

Key words: late Quaternary, Aeolian deposits, OSL

## 1. INTRODUCTION

Since the discovery of optical dating on quartz ((Huntley et al., 1985) and feldspar (Hutt et al., 1988) there have been many methodological developments. Especially the development of the Single Aliquot Regenerative-dose (SAR) protocol for quartz (Murray & Wintle, 2000), has improved the reliability of quartz optical dating and since then it has been extensively used to established chronological framework for Quaternary sedimentary successions of both shallow-marine and continental

1  
2  
3  
4 environments (e.g. Spencer and Robinson, 2008; Jacobs, 2008; Lancaster, 2008; Galli et al., 2009;  
5 Andreucci et al., 2010; Thiel et al., 2011; Pascucci et al., 2014; Lamothe 2016).

6 The range of quartz OSL dating is usually limited to by the saturation of the quartz OSL dose response  
7 curve which typically occurs at ~200 Gy Wintle and Murray (2006) suggested an upper dating limit for  
8 the quartz OSL (fast component dominated samples) to ~2D0 (i.e. 86% of the saturation level of the  
9 dose response curve). For samples with doses beyond 2D0 the reliability is not yet proven and results  
10 should be interpreted with caution. There are published examples that quartz SAR-OSL dating method  
11 underestimates the age of older deposits (e.g. Buylaert et al., 2007; Lowick et al., 2010).

12 Feldspar signals are attractive because their dose response curves saturate at much higher dose than  
13 quartz OSL- However, feldspar infrared stimulated luminescence (IRSL) signals measured at ambient  
14 temperatures are affected by anomalous fading (Spooner, 1994) which causes age underestimations  
15 unless an adequate correction is made (Huntley and Lamothe, 2001). However, it is claimed in Huntley  
16 and Lamothe (2001) that it should not be applied to samples older than 20-50 ka. Furthermore, fading  
17 corrections are based on untestable assumptions (Morthekai et al., 2008). Recently, research effort was  
18 made to identify a feldspar signal which shows less fading (Thomsen et al., 2008) and to develop a  
19 feldspar dating protocol with a reduced or negligible fading correction. (e.g. post-IR IRSL protocols in  
20 Buylaert et al., 2009, 2012; Thiel et al., 2011). In this paper we adopt the post-IR IR<sub>290</sub> protocol put  
21 forward by Thiel et al. (2011) and tested extensively in Buylaert et al. (2012) to further test this method  
22 on samples from Sardinia with aeolian and thus well-bleached sediment. The post-IR IR<sub>290</sub> results are  
23 compared with independent age control (U-Th ages), quartz SAR-OSL ages and historical information.  
24  
25  
26  
27  
28

## 29 30 **2. REGIONAL SETTING AND STRATIGRAPHY**

31  
32 The Island of Sardinia is located in the north-western Mediterranean Sea (Fig. 1), emplaced by  
33 eastward rotation of the European plate during the Oligocene-early Miocene (Carmignani et al., 2001).  
34 In association with this rotation normal and transcurrent faults dissected Sardinian pre-Miocene  
35 deposits and allowed the formation of a series of half graben that were filled with Miocene continental  
36 and marine deposits and calc-alkaline volcanic bodies (Casula et al., 2001). During the early Pliocene  
37 widespread volcanism and basin uplift occurred, but north-west Sardinia is considered to have  
38 remained tectonically stable since the Late Pliocene (Lambeck et al., 2004; Ferranti et al., 2006).  
39 The Quaternary deposits crops out quasi-continuously along the study areas and are characterized by an  
40 alternation of shallow marine and aeolian/colluvial strata spanning in time from Marine Isotopic Stage  
41 (MIS 8; ca 275 ka) to MIS 1 (ca 5 ka; Andreucci et al., 2010; 2011; Pascucci et al., 2014).  
42  
43  
44

### 45 **2.1 STUDY AREA**

46  
47 Three study areas have been selected along the Sardinian coasts: The Bue Marino cave (Orosei Gulf;  
48 NE Sardinia), south Alghero coast and the Alghero bay (NW Sardinia; Fig. 1A).

49 The area of Bue Marino cave is characterized by relatively high (50–400 m) and steep carbonate sea-  
50 cliffs variable affected by karst phenomena (Fig. 1B). The Bue Marino is coastal cave carved on  
51 Jurassic limestone and almost completely filled by wind-blown sediment. The sedimentary succession  
52 is sandwiched between calcareous crusts (Carobene and Pasini, 1980).

53 The south Alghero coast is characterized by small low-cliff-bounded embayments with basal marine  
54 terraces backed by 40 m high hills oriented almost parallel to the present-day coastline (Fig. 1C). The  
55 sedimentary succession which rests on the bedrock, is characterized by an alternation of thin (3 m-  
56 thick) shallow-marine strata and thick (20 m-thick) aeolianites. The bedrock is composed of mainly  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Triassic limestone, Oligo-Miocene andesitic pyroclastic deposits and minor Permian, red, sandy to  
5 gravelly bodies (Pascucci et al., 2014).

6 The Alghero Gulf is sea-ward open on the West and laterally bordered by Mesozoic cliffs. The  
7 embayment is dominated by 5 km long sandy beach-ridge system backing a N-S oriented lagoon  
8 system named Calick (Fig. 1D; Ginesu et al 2001; Zucca et al., 2014). The dune cordon cropping out  
9 on the study area has a maximum thickness of 10 metres and was human stabilized by plantation during  
10 the 1950s.  
11  
12

### 13 14 **3. METHODS**

15  
16 A total of nine samples from the three study areas have been collected for luminescence analyses  
17 (Tables 1). Moreover, two samples from the calcareous crusts cropping out at the Bue Marine cave  
18 were collected for U-series measurements.  
19  
20

#### 21 **3.1 LUMINESCENCE ANALYSES**

22  
23 Samples were prepared under red light conditions at the University of Sassari (Italy) to extract quartz  
24 and K-rich feldspar grains in the size range between 180-250  $\mu\text{m}$ , following the standard procedure  
25 (Stokes, 1992), and were analysed at the Nordic Laboratory for Luminescence Dating (Institute for  
26 Geoscience, Aarhus University, Risø DTU, Denmark).  
27

28 All OSL measurements were conducted on an automated Risø TL/OSL model DA-20 (Bøtter-Jensen et  
29 al., 2002). Quartz was stimulated using blue LEDs (470 nm) and the luminescence detected through a U-  
30 340 filter. Feldspar IRSL was stimulated using IR LEDs (875 nm) and detection was through the  
31 standard blue filter pack (Schott BG-39 + Corning 7-59). For the quartz measurements large (8mm)  
32 aliquots were used. Because of the bright luminescence signals, feldspar measurements were made on  
33 small (2 mm) aliquots. OSL signal from quartz in this region is known to be dominated by a fast  
34 component (Andreucci et al., 2010; Thiel et al. 2010).  
35

36 Initial checks using infra-red stimulated luminescence were conducted on quartz materials to test for  
37 residual feldspar contamination (Smith et al., 1990). These checks proved negative, so a blue-light  
38 stimulation simple SAR protocol was used for quartz equivalent dose ( $D_e$ ) measurements (Murray &  
39 Wintle, 2000). OSL measurements (80 s) were made at 125°C, after pre-heating aliquots for 10 s at  
40 260°C for samples from the Bue Marino cave and South Alghero coast. The pre-heat value was  
41 experimentally derived on the basis of the results of a dose recovery pre-heat plateau test. A cut-heat of  
42 220°C was applied to each aliquot before the test dose measurement and a high temperature blue-light  
43 clean-out for 40 s at 280°C after every SAR cycle. Samples from the modern dune of Alghero bay  
44 underwent, instead, to a lower pre-heat (180°C) and cut-heat (160°C) temperatures for the same  
45 amount of time to avoid a thermal transfer (Madsen and Murray, 2009). Five regeneration points were  
46 measured during the SAR procedure, including a recycling point, which was used to determine the  
47 effectiveness of the sensitivity corrections. A dose recovery was carried out for some samples of the  
48 different study areas. The initial ~2 s of the luminescence signal, less a background derived from the 4  
49 – 8 s, was used for all calculations.  
50  
51  
52  
53  
54

55 SAR post infrared stimulation (post-IR IRSL) at elevated temperature (290°C) protocol (Thiel et al.,  
56 2011; Buylaert et al., 2012) was applied to K-rich feldspar grains in order to evaluate the  $D_e$  of the  
57 same sample measured with quartz SAR-OSL method. After a preheat at 320°C for 60 sec, the samples  
58 were bleached with IR diodes at first at 50°C for 200 s and then again at 290°C for 200 s to measure  
59 the post-IR IR<sub>290</sub> signal. The same preheat conditions were used for natural, regenerative and test dose  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 measurements. An IR illumination at 325°C for 200 s was inserted at the end of each SAR  
5 measurement cycle. Fading measurements (*g* value, Aitken, 1985) were carried out using the same  
6 SAR protocol used for *D<sub>e</sub>* determinations. The *g* values were normalised to a *t<sub>c</sub>* = 2 days (Huntley and  
7 Lamothe, 2001). The initial ~2 s of the luminescence signal, less a background derived from the last  
8 ~10 s, was used for all calculations. A dose recovery test was also undertaken by adding a dose on top  
9 of a very young sample.

10  
11  
12 For each sample, approx. 200 gr of sediment were collected to estimate the water content and to  
13 calculate the natural radioactivity (dose rate, *Dr*, Table 1A).

14 Dose rate calculations are based on high-resolution gamma spectrometry following the methodology  
15 described by Murray et al. (1987) to give natural radionuclide concentrations, and knowledge of burial  
16 depth to give cosmic ray contribution (Prescott and Hutton, 1994). The required corrections on the *Dr*  
17 value for cementation (lithified samples) were taken into account and calculated following Andreucci  
18 et al. (2009). The lifetime average water content since deposition is assumed to lay between present-  
19 day and saturated values and has been estimated for each sample based on sedimentological and  
20 hydrogeological conditions. K-feldspar grains have an internal radioactivity due to the presence of  
21 structural <sup>40</sup>K in the crystal structure. Hence, for the K-feldspars dose rates a contribution to the final  
22 dose rate from <sup>40</sup>K was incorporated based on the assumption that the internal K content is 12.5±0.5 %  
23 (Huntley and Baril, 1997).  
24  
25  
26  
27

### 28 **3.2 U-SERIES METHOD**

29  
30  
31 U/Th extraction from two calcareous crusts (flowstone) were carried out at the U-series Laboratory of  
32 the University of Bergen (Norway) using a actinide-specific, phosphate/phosphine based (Eichrom  
33 TRU) liquid ion exchanger mini-columns, supported on an inert substrate (Amberlite XAD-7)  
34 (Hellstrom, 2003; Peterson et al., 2007; Yang, 2009). Isotopic measurements were performed on a Nu  
35 Plasma HR multicollector ICP-MS with a U-Pb collector block at the Department of Geology,  
36 University of Oslo. Analyses were done in dry plasma using a DSN-100 desolvating nebuliser with a  
37 sample uptake rate of 0.1 mL/min. The mixed U and Th solution was analysed in two separate  
38 procedures. First, uranium isotopes with mass 236, 235 and 234 were determined in ion counters and  
39 thorium with mass 232 in a Faraday cup. The second procedure measures thorium mass 229 and 230 in  
40 an ion counter. Tailing from 238 and 232 was corrected by measuring half masses and using an  
41 exponential interpolation. Fractionation of the instrument has been determined on a daily basis by  
42 analysing mass 235 and 238 of a natural uranium solution in Faraday cups using  $238/235 = 137.88$ . The  
43 reproducibility of each measured was 0.11% ( $2\sigma$ ). Repeated analyses of BR5 (a high uranium  
44 speleothem powder standard) gave an age of  $125.862 \pm 1.546$  kyr ( $n = 12$ ), with a reproducibility of  
45 measured  $234\text{U}/238\text{U}$  ratio of 0.59% ( $2\sigma$ ). Age determinations were based on measured atomic mass  
46 ratios of  $235\text{U}/236\text{U}$ ,  $235\text{U}/234\text{U}$ ,  $236\text{U}/234\text{U}$ ,  $232\text{Th}/229\text{Th}$ ; and  $229\text{Th}/230\text{Th}$ . Data  
47 reduction, error optimization and propagation were done using tailored software (Lauritzen and  
48 Lundberg, 1997) which has been rewritten for the Windows environment. The presence of the non-  
49 authigenic  $230\text{Th}$  in the carbonate samples as indicated by the low  $230\text{Th}/232\text{Th} < 20$ , requires a  
50 correction for detrital  $230\text{Th}$  contamination. This correction was performed directly by the  
51 software, using initial  $234\text{U}/238\text{U}$  calculated from  $2230\text{Th}/234\text{U}$  ratio (Table 3).  
52  
53  
54  
55  
56  
57  
58

## 59 **4. RESULTS**

#### 4.1 Study area 1: Bue Marino cave

At Bue Marino, one opaque PVC tube ( $D = 8$  cm;  $L = 40$  cm) of freshly exposed reddish colluvial strata (BUE1) and three sample blocks (ca  $50 \times 50 \times 40$  cm) of aeolianites (BUE2, BUE4, BUE5) were collected (Figs 2; Table 1). Samples were collected along an artificial scarp composed of a basal sandy unit (8.5 m-thick) capped by a 50 cm-thick reddish colluvial strata (Figs 2). The succession is bounded at the base and at the top by 50 cm-thick calcareous crusts.

Aeolianites are characterized by medium to coarse-grained carbonate-rich sandstone made up of: 70-82%  $\text{CaCO}_3$ , 16-28% siliciclastics, 1% fines and 1% organics. Instead, colluvial strata is composed of a matrix-supported sandy to pebbly deposit made up of: 31%  $\text{CaCO}_3$ , 13% siliciclastics, 55% fines and 1% organics.

Quartz equivalent dose ( $D_e$ ) values lay between 165 and 141 Gy without any significant decrease from the base to the top of the succession. A dose recovery test on sample BUE4 gives a satisfactory dose ratio of  $0.99 \pm 0.05$  ( $n=12$ ). The quartz dose rates range from 2.21 to 3.05 Gy/ka which is consistent with the dose rates of palaeosols in the area but relatively high for Sardinian aeolianites (Andreucci et al., 2011). The quartz OSL signals for all samples are close to saturation ( $D_e > 2D_0$ ) so the dates between 70 and 50 ka probably give only minimum ages (Table 1).

Post-IR IR290 K-feldspar  $D_e$  values range between  $281 \pm 10$  and  $320 \pm 9$  Gy and the K-feldspar dose rates vary between 3.02 to 3.86 Gy/ka. A dose response curve of sample (BUE2,  $D_e = 315 \pm 6$  Gy,  $n=6$ ) shows that the natural feldspar signal lies far below the saturation level of the curve (Fig. 2D). The dose recovery tests for K-feldspar pIRIR290 dating was carried out by adding a large beta dose (49 Gy) on top of a modern K-feldspar sample (MP7; study area 3, see following section) and subtracting the natural  $D_e$  ( $2.3 \pm 0.2$  Gy) from the measured dose. The dose recovery ratio is  $0.97 \pm 0.08$  ( $n=16$ ) confirming the suitability of this protocol for measuring dose. Fading measurements were carried out on three aliquots from each sample after measuring the  $D_e$  values. The mean pIRIR290  $g_{2\text{days}}$  value is  $1.50 \pm 0.10$  %/decade ( $n=12$ ). Buylaert et al. (2012) suggest that these fading rates may be artefacts of our measurement procedure and that the evidence that these apparent fading rates reflect loss of charge during storage is not convincing. Therefore, the ages have not been corrected for fading. The uncorrected pIRIR<sub>290</sub> ages range from the base to the top:  $104 \pm 12$  ka,  $81 \pm 9$  ka,  $83 \pm 9$  ka and  $83 \pm 9$  ka. U-series dates of the floatstone that are bounding the sedimentary succession give ages of 100 ka at the base and 80 ka at the top which is in good agreement with the pIRIR290 ages.

#### 4.2 Study area 2: South Alghero coast

One sample block (ca  $50 \times 50 \times 40$  cm) of aeolianite (AHO16) was collected (Figs 3; Table 2). Sample was collected at the base of a 20 m-thick lithified transgressive dune body. Sampled block is characterized by medium to coarse-grained carbonate-rich sandstone made up of: 90%  $\text{CaCO}_3$ , 8% siliciclastics, 1% fines and 1% organics.

Quartz  $D_e$  value is  $67 \pm 4$  Gy and the quartz dose rate is  $0.88 \pm 0.04$  Gy/ka; both values are typical for Sardinian aeolianites (Andreucci et al., 2009, 2011). The studied sample shows good OSL characteristics (fast component, recycling values close to unity ( $1.12 \pm 0.14$   $n = 12$ ; dose recovery ( $1.02 \pm 0.04$ ,  $n = 18$ )). Because of the low dose rate the quartz OSL signal lays below the saturation limit (69 % of saturation) and AHO16 has a reliable age of  $76 \pm 6$  ka.

K-feldspars  $D_e$  value is of  $124 \pm 4$  Gy and the  $D_r$  of  $1.69 \pm 0.07$  Gy/ka. The K-feldspar sample has a fading-uncorrected age of  $73 \pm 5$  ka.

### 4.3 Study area 3: Alghero bay

Four opaque PVC tubes ( $D = 8$  cm;  $L = 40$  cm) of freshly exposed modern coastal deposits (MP4, MP5, MP6, MP7) were collected (Figs 4; Table 3).

Three samples (MP5, MP6, MP7) were collected on a natural 5 m-thick scarp along the stabilized coastal cordon, one (MP4), instead on the modern backshore, in front of an active blow out (Fig. 4A). Samples are characterized by medium to coarse-grained carbonate-rich sandstone made up of: 55-75%  $\text{CaCO}_3$ , 33-43% siliciclastics, 1% fines and 1% organics.

Quartz  $D_e$  values (MP7:  $1.69 \pm 0.06$  Gy, MP6:  $1.22 \pm 0.07$  Gy, MP5:  $0.04 \pm 0.01$  Gy, MP4:  $0.006 \pm 0.01$  Gy) decrease significantly from the base to the top of the succession, as expected. The  $D_r$  values of all samples are around  $0.07 \pm 0.01$  Gy/ka as expected for beach-dune deposits (ref?).

The studied sample shows good OSL characteristics such as recycling values within  $1.0 \pm 0.05$  and dose recovery measurements on 9 aliquots were performed with a mean value of  $1.03 \pm 0.08$ .

Samples collected on the natural scarp (MP5, MP6, MP7) from the base to the top have ages of  $2450 \pm 150$  yr,  $1670 \pm 100$  yr and  $60 \pm 20$  years respectively. The sample (MP4) from the modern backshore zone shows a zero age ( $9 \pm 14$  years). The pIRIR290 age for sample MP7 is  $1520 \pm 150$  years which is clearly too old. The corresponding  $D_e$  value of  $2.3 \pm 0.2$  Gy ( $n=6$ ) may represent the true unbleachable residual of the pIRIR290 signal or it may be the result of thermal transfer during preheating.

## 5. DISCUSSION

The results from the study area 1 (Bue Marino cave) clearly indicate that the quartz-OSL signal can be saturated already for a  $D_e$  of 141 Gy, corresponding to a minimum age of about 50 ka (Fig. 2C). Instead the pIRIR290 measurements on K-feldspar give uncorrected ages between  $\sim 100$  and  $\sim 80$  ka, in good agreement with the U-series data.

Along the south Alghero coast (study area 2) the thick lithified transgressive dune system is widely recognized in the literature to represent the beginning of the last glacial phase (*post* 80 ka; Andreucci et al., 2010; Pascucci et al., 2014 Figs 3). The coupled quartz-OSL and K-feldspar post-IR IRSL protocols on sediments coming from the same sampled block show interesting results. Firstly, the quartz OSL  $D_e$  ( $67 \pm 4$  Gy) lays below the 2D0 saturation limit giving a reliable age of  $76 \pm 6$  ka, that is in perfect agreement with the previous study (Andreucci et al., 2010; 2011). The fading-uncorrected post-IR IRSL290 age of  $73 \pm 5$  ka confirms the quartz result.

Finally, the third study area (Alghero modern dune system), shows instead some limitations for the pIRIR290 measurements on K-feldspar grains when applied to very young samples. We report a age of  $1520 \pm 150$  a for a sample which is only  $60 \pm 20$  a old based on historical information and quartz OSL; the feldspar pIRIR290  $D_e$  is  $2.3 \pm 0.2$  Gy; this is significant when dating young (Late Holocene, modern) samples but becomes very quickly unimportant for older samples. We suggest that, if possible, modern or very young samples with quartz OSL age control should always be collected during fieldwork to be able to place some lower limit to the residual pIRIR290 doses for investigated sediments.

## 6. CONCLUSIONS

This study has contributed to the further testing of the post-IR elevate temperature ( $290^\circ\text{C}$ ) IRSL protocol applied to coarse-grained K-feldspar for dating well-bleached sedimentary samples. We conclude that:

- 1
- 2
- 3
- 4 1. The dating technique shows great promise for dating coastal deposits in Sardinia that are at or
- 5 beyond the quartz OSL dating limit ( $D_e > 85\%$  of  $2D_0$ ) as indicated by the agreement between
- 6 the uncorrected post-IR IRSL results and the bracketing U-Th ages.
- 7
- 8 2. The pIRIR<sub>290</sub> protocol shows a significant age offset for young (Holocene) samples but
- 9 becomes quickly insignificant as deposits get older.
- 10
- 11

## 12 **Acknowledgements**

13  
14  
15 We are indebted to Prof. Lars Clemmensen for the field assistance and the stimulating discussions.  
16 Financial support was provided by a young researcher grant from Regione Autonoma Sardegna: PO  
17 Sardegna FSE 2007–2013 and L.R. 7/2007 “Promozione della ricerca scientifica e dell’innovazione  
18 tecnologica in Sardegna” (resp. Stefano Andreucci).  
19  
20

## 21 **References:**

22  
23 Andreucci, S., Pascucci, V., Murray, A., Clemmensen, L.B., 2009. Late Pleistocene coastal evolution  
24 of San Giovanni di Sinis, west Sardinia (Western Mediterranean). *Sedimentary . Geology*. 216, 104–  
25 116.  
26

27  
28 Andreucci, S., Clemmensen, L.B., Pascucci, V., 2010. Transgressive dune formation along a cliffed  
29 coast at 75 ka in Sardinia, Western Mediterranean: a record of sea-level fall and increased windiness.  
30 *TerraNova* 22, 424-433.  
31

32  
33 Andreucci, S., Bateman, M.D., Zucca, C., Kapur, S., Akaşit, İ., Dunajko, A., Pascucci, V., 2011.  
34 Evidence of Saharan dust in Upper Pleistocene reworked palaeosols of Northwest Sardinia, Italy:  
35 Palaeoenvironmental implications. *Sedimentology* doi: 10.1111/j.1365-3091.2011.01285.x, in press.  
36  
37

38 Auclair, M., Lamothe, M., Huot, S., 2003: Measurement of anomalous fading for feldspar IRSL using  
39 SAR. *Radiation Measurements* 37, 487-492.  
40

41  
42 Buylaert, J.P., Huot, S., Murray, A.S., Van den haute, P., 2011: Infrared stimulated luminescence  
43 dating of an Eemian (MIS 5e) site in Denmark using K-feldspar. *Boreas* 40, 46-56.  
44

45 Buylaert J-P, Jain M, Murray AS, Thomsen KJ, Thiel C and Sohbaty R, 2012. A robust method for  
46 increasing the age range of feldspar IRSL dating. *Boreas* 41(3): 435–451.  
47

48  
49 Buylaert, JP., Murray, AS., Thomsen, KJ., Jain, M., 2009. Testing the potential of an elevated  
50 temperature IRSL signal from K-feldspar. *Radiation. Measurements* 44, 560-565.  
51

52  
53 Bøtter-Jensen, L., Thomsen, K.J., Jain, M., 2010: Review of optically stimulated luminescence (OSL)  
54 instrumental developments for retrospective dosimetry. *Radiation Measurements* 45, 253-257.  
55

56 Carmignani, L., Barca, S., Oggiano, G., Pertusati, P.C., Salvadori, I., Conti, P., Eltrudis, A., Funedda,  
57 A., Pasci, S., 2001. Note illustrative della Carta Geologica della Sardegna a scala 1:200.000. *Memorie*  
58 *descrittive Carta Geologica Italiana*: Roma.  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 Carobene, C., Pasini G., 1980. Contributo alla conoscenza del Pleistocene superiore e dell'Olocene del  
5 Golofa di Orosei (Sardegna Orientale). *Boll. Soc Adriatic. Scienze* 64, 3-36.  
6  
7  
8 Casula, G., Cherchi, A., Montandert, L., Murru, M., Sarria, E., 2001. The Cenozoic grabens system of  
9 Sardinia: geodynamic evolution from new seismic and field data. *Marine and Petroleum Geology* 18,  
10 863– 888.  
11  
12  
13 Ferranti, L., Antonioli, F., Mauz, B., Amorosi, A., Dai Pra, G., Mastronuzzi, G., Monaco, C., Orrù, P.,  
14 Pappalardo, M., Radtke, U., Renda, P., Romano, P., Sansò, P., Verrubbi, V., 2006. Markers of the last  
15 interglacial sea-level high stand along the coast of Italy: Tectonic implications. *Quatern. Int.* 146, 30-54.  
16  
17  
18 Galli, A., Panzeri, L., Martini, M., Sibilina, E., Vignola, P., Andò, S., Pini, R., Rossi, P.M., 2009.  
19 Optically stimulated luminescence dating of a stratigraphic Late Glacial–Holocene sequence in the Po  
20 plain (Bubano quarry, Bologna, Italy). *Quaternary International* 199, 45-55.  
21  
22  
23 Huntley, D. J., Baril, M. R., 1997. The K content of the K-feldspars being measured in optical dating  
24 or in thermoluminescence dating. *Ancient TL* 15, 11–13.  
25  
26  
27 Huntley, D.J., Lamothe, M., 2001. Ubiquity of anomalous fading in K-feldspars, and the measurement  
28 and correction for it in optical dating. *Can. J. Earth Sci.* 38, 1093–1106.  
29  
30  
31 Huntley, D.J., Godfrey-Smith, D.I., Thewalt, M.L.W., 1985. Optical dating of sediments. *Nature* 313,  
32 105–107.  
33  
34  
35 Hutt, G., Jaek, I., Tchonka, J., 1988. Optical dating: K-feldspars optical response stimulation spectra.  
36 *Quaternary Science Reviews* 7, 381–385.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Jacobs Z., 2008. Luminescence chronologies for coastal and marine sediments. *Boreas* 37, 508–535.
- Lambeck, K., Antonioli, F., Purcell, A., Silenzi, S., 2004. Sea-level change along the Italian coast for the past 10,000 yr. *Quatern. Sci. Rev.* 23, 1567-1598.
- Lamothe, M. (2016) - Luminescence dating of interglacial coastal depositional systems: Recent developments and future avenues of research. *Quaternary Science Reviews* 146, 1-27.
- Lamothe, M., Auclair, M., Hamzaoui, C., Huot, S., 2003. Towards a prediction of longterm anomalous fading of feldspar IRSL. *Radiation Measurements* 37, 493–498.
- Lancaster N., 2008. Desert dune dynamics and development: insights from luminescence dating. *Boreas* 37, 559-573.
- Lowick S.E., Preusser, F., Pini, R., Ravazzi, C., 2010. Underestimation of fine grain quartz OSL dating towards the Eemian: Comparison with palynostratigraphy from Azzano Decimo, northeastern Italy. *Quaternary Geochronology* 5, 583–590.
- Madsen, A.T., Murray, A.S., 2009: Optically stimulated luminescence dating of young sediments: a review. *Geomorphology* 109, 3-16.

- 1  
2  
3  
4  
5 Murray, A.S., Wintle, A.G., 2000. Luminescence dating of quartz using an improved single-aliquot  
6 regenerative-dose protocol. *Radiation Measurements* 32, 57–73.  
7  
8  
9 Morthekai, P., Jain, M., Murray A.S., Thomsen, K.J., Bøtter-Jensen, L., 2008: Fading characteristics of  
10 martian analogue materials and the applicability of a correction procedure. *Radiation Measurements* 43,  
11 672-678.  
12  
13  
14 Pascucci, V., Sechi, D. & Andreucci, S. (2014) - Middle Pleistocene to Holocene coastal evolution of  
15 NW Sardinia (Mediterranean Sea, Italy). *Quaternary International*, 328-329, 3–20.  
16  
17  
18 Spencer, J.Q.G., Robinson R.A.J., 2008. Dating intramontane alluvial deposits from NW Argentina  
19 using luminescence techniques: Problems and potential. *Geomorphology* 93, 144-155.  
20  
21  
22 Spooner, N. A., 1994: The anomalous fading of infrared-stimulated luminescence from feldspars.  
23 *Radiation Measurements* 23, 625-632.  
24  
25  
26 Stokes, S., 1992. Optical dating of young (modern) sediments using quartz: results from a selection of  
27 26 depositional environments. *Quaternary Sci. Rev.* 11, 153-159.  
28  
29  
30 Thiel, C., Coltorti, M., Tsukamoto, S., Frechen, M., 2010. Geochronology for some key sites along the  
31 coast of Sardinia (Italy). *Quat. Int.* 222, 36–47.  
32  
33  
34 Thiel, C., Buylaert, J-P., Murray, AS., Terhorst, B., Hofer, I., Tsukamoto, S., Frechen, M., 2011.  
35 Luminescence dating of the Stratzing loess profile (Austria) - Testing the potential of an elevated  
36 temperature post-IR IRSL protocol. *Quatern. Int.* 234, 23-31.  
37  
38  
39 Thomsen, K.J., Murray, A.S., Jain, M., Bøtter-Jensen, L., 2008. Laboratory fading rates of various  
40 luminescence signals from feldspar-rich sediment extracts. *Radiation Measurements* 43, 1474–1486.  
41  
42  
43 Wallinga, J., Bos, A.J.J., Dorenbos, P., Murray, A.S., Schokker, J., 2007. A test case for anomalous  
44 fading correction in IRSL dating. *Quaternary Geochronology* 2, 216–221.  
45  
46  
47 Wintle, AG. and Murray, AS. (2006) A review of quartz optically stimulated luminescence  
48 characteristics and their relevance in single-aliquot regeneration dating protocols. *Radiation*  
49 *Measurements* 41, 369–391.  
50  
51  
52 Zucca, C., Sechi, D., Andreucci, S., Shaddad, S.M., Deroma, M., Madrau, S., Previtali, F., Pascucci, V.,  
53 & Kapur, S. (2014) - Pedogenic and palaeoclimatic evidence from an Eemian calcrete in North-western  
54 Sardinia (Italy). *European Journal of Soil Sciences*, 65, 420-435.  
55  
56

## 57 **Captures to the figures**

58  
59 Fig. 1 – Location map of the studied areas. A) Satellite view of the Sardinia Island and inset map of the  
60 west Mediterranean region. B) Detailed view of the Bue Marino cave entrances (studied area 1). C)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Satellite view of the South Alghero rocky coast (Study area 2). D) Satellite view of the SE side of the Alghero bay and the modern coastal cordon (Study area 3), fragmented by several blow out, and stabilized by artificial plantation.


Fig. 2 – Study area 1: Bue Marino cave. A) Detailed view of the basal part of the aeolian deposits. Note that the sedimentary body drapes the cave flank encrusted by a calcareous crust (floatstone). The black square indicates the sample position for U/Th analysis and italic letters the relative age. The black circle the OSL sample position. Scale on the left side is a hammer (40 cm high). B) Detail view of central to top parts of the succession characterized, from bottom up, by aeolianites and thin colluvial strata. The succession is capped by a 50 cm-thick floatstone. The white square indicates the sample position for U/Th analysis and italic letters the age. The black circle the OSL sample positions. Scale on the right side is a sitting woman (1 m high). C) Schematic drawing showing the relationship between samples at the Bue Marino cave. The white square indicates the sample position for U/Th and italic letters the age. The black circle the OSL sample positions D) K-feldspar post-IR IRSL290 growth curve for sample BUE2 showing a recycling point (grey circle) and a recuperation point (white triangol). The insets shows the natural pIRIR290 decay curve. Note that sensitivity corrected natural signal (white circle) is well below the saturation limit.

Fig. 3 – Study area 2: south Alghero coast. A) Panoramic view of the 20-m thick transgressive dune body. Note that the aeolian deposits span in time from the Marine Isotopic Stage (MIS 4) to MIS 3. On the top of the hill a MIS 8 (250 ka) dunes crop out. The black circles indicate previous OSL date and italic letters the relative ages (modified from Andreucci et al., 2010). Scale on the left side is a white line. B) Detail view of the basal part of the dune body. The white square indicates the new OSL ages and italic letters the relative age. The white circles previous OSL samples and italic letters the relative ages. Scale on the left side is a black line.

Fig. 4 – Study area 3: Alghero bay. A) Panoramic view of the Alghero bay, sandy beach system and dune cordon human stabilized during the 50<sup>th</sup>. The white circle indicates OSL sample and italic letters the relative age B) Detail view of the basal part of the dune body. The white circle indicates the OSL sample position and italic letters the relative age. C) Detail view of the top part of the dune body. The white circles indicate the OSL sample positions and italic letters the relative ages.



Figure



[Click here to download high resolution image](#)




Figure

[Click here to download high resolution image](#)







**Table 1**

Table 1. Summary of principal radionuclides, water contents and total quartz and K-feldspar dose rates for all samples. The quartz and K-feldspar grain size range was 180-250  $\mu\text{m}$ . The conversion factors from activity concentrations to dose rate are taken from Olley et al. (1996). For the K-feldspar dose rates an internal K content of  $12.5 \pm 0.5\%$  was assumed (Huntley and Baril, 1997).

Site & Location	Sample code	Elevation <sup>(1)</sup> (m)	<sup>238</sup> U (Bq kg <sup>-1</sup> )	<sup>226</sup> Ra (Bq kg <sup>-1</sup> )	<sup>232</sup> Th (Bq kg <sup>-1</sup> )	<sup>40</sup> K (Bq kg <sup>-1</sup> )	Moisture <sup>(2)</sup> %	Quartz Dose rate (Gy ka <sup>-1</sup> )	K-feldspar Dose rate (Gy ka <sup>-1</sup> )
Alghero bay	MP4	1	14 ± 4	5.0±0.3	8.9±0.3	70 ± 3	25	0.65 ± 0.04	1.53±0.07
	MP5	3	14 ± 3	15.3 ± 0.2	5.9 ± 0.2	93 ± 4	25	0.69 ± 0.04	1.57±0.07
	MP6	4	10 ± 3	11.2 ± 0.2	14.3 ± 0.2	126 ± 5	25	0.73 ± 0.04	1.61±0.07
	MP7	4.5	4 ± 3	5.4 ± 0.2	5.9 ± 0.2	226 ± 3	25	0.73 ± 0.04	1.61±0.07
South Alghero	AHO16 <sup>(3)</sup>	2	-	-	-	-	-	0.88 ± 0.04	1.73±0.08
Bue Marino	BUE1	9	5 ± 5	8.1 ± 0.4	16.2 ± 0.5	642 ± 13	15	2.80 ± 0.13	3.68±0.14
	BUE2	2.5	10 ± 6	6.5 ± 0.4	10.3 ± 0.5	580 ± 13	5	2.20 ± 0.11	3.08±0.12
	BUE4	6.5	10 ± 5	9.5 ± 0.4	19.0 ± 0.6	546 ± 13	5	2.66 ± 0.12	3.54±0.13
	BUE5	7.5	10 ± 5	8.3 ± 0.5	11.9 ± 0.4	815 ± 16	5	2.93 ± 0.14	3.81±0.15

Notes:

<sup>(1)</sup> Elevation corresponds to the elevation of samples collected above the present sea-level.

<sup>(2)</sup> Moisture corresponds to water content chosen as life-time average for the site. An absolute error of 4% is assumed. The effect of cementation is incorporated in the final dose rate values following Andreucci et al. (2009).

<sup>(3)</sup> For this sample only dose rates are given, other information can be found in Buylaert et al. (2012).

Table 2. Summary of quartz OSL and K-feldspar pIRIR<sub>290</sub> D<sub>e</sub> values, pIRIR<sub>290</sub> fading rates ( $g_{2\text{days}}$ ), quartz OSL and uncorrected pIRIR<sub>290</sub> ages. The number of individual aliquots contributing to D<sub>e</sub> is denoted with n. Fading rates are determined on three aliquots previously used for D<sub>e</sub> determination. Uncertainties represent one standard error.

Site & Location	Sample code	OSL D <sub>e</sub> (Gy)	n	pIRIR <sub>290</sub> D <sub>e</sub> (Gy)	n	$g_{2\text{days}}$ (%/decade)	OSL age (ka)	pIRIR <sub>290</sub> age (ka)
Alghero bay	MP4	0.011±0.006	32	1.65 ± 0.10	3	1.05±0.34	16±10 <sup>(3)</sup>	1080±80 <sup>(3)</sup>
	MP5	1.82 ± 0.05	34	4.34 ± 0.10	3	1.54±0.11	2640±180 <sup>(3)</sup>	2770±150 <sup>(3)</sup>
	MP6	1.21 ± 0.05	34	3.74 ± 0.12	3	1.42±0.03	1660±120 <sup>(3)</sup>	2320±130 <sup>(3)</sup>
	MP7	0.060 ± 0.008	32	1.92 ± 0.16	5	1.58±0.05	83±13 <sup>(3)</sup>	1200±160 <sup>(3)</sup>
South Alghero	AHO16 <sup>(2)</sup>	67±4	30	127±7	9	1.44±0.19	76 ± 6	73 ± 5
Bue Marino	BUE1	>200 <sup>(1)</sup>	4	317±7	6	1.77±0.13	> 70 <sup>(1)</sup>	86 ± 4
	BUE2	“	4	332±6	6	1.19±0.20	> 90 <sup>(1)</sup>	108 ± 5
	BUE4	“	4	296±10	5	1.63±0.26	> 75 <sup>(1)</sup>	84 ± 4
	BUE5	“	4	338±9	6	1.45±0.09	> 70 <sup>(1)</sup>	89 ± 4

## Notes:

<sup>(1)</sup> The natural signal lies above the 86% of the saturation level of the dose response curve (see Fig. 2a). The dose corresponding to 2\*D<sub>0</sub> (86% saturation) is given and the quartz age is considered a minimum age.

<sup>(2)</sup> Data from Buylaert et al. (2012).

<sup>(3)</sup> Age expressed in years from 2010.

**Table 3**

Table3. Summary of Uranium concentration, U/Th ratios, raw and corrected ages. Uncertainties represent two standard error ( $2\sigma$ ).

<b>ID Lab</b>	<b>ID sample</b>	<b>U (ppm)</b>	$^{234}\text{U}/^{238}\text{U}$	<b>2<math>\sigma</math>+ (%)</b>	<b>2<math>\sigma</math>- (%)</b>	$^{230}\text{Th}/^{234}\text{U}$	<b>2<math>\sigma</math>+ (%)</b>	<b>2<math>\sigma</math>- (%)</b>	$^{230}\text{Th}/^{232}\text{Th}$	<b>2<math>\sigma</math>+ (%)</b>	<b>2<math>\sigma</math>- (%)</b>	<b>Age (ka)</b>	<b>Initial <math>^{234}\text{U}/^{238}\text{U}</math>*</b>	<b>Corrected Age (ka)</b>
857	AL3-2	0.0257	1.035	2.79	0.03	0.647028	6.67	0.04	4	6.11	0.27	113,3 $\pm$ 14	1.04 $\pm$ 0.029	<b>86,1<math>\pm</math>13</b>
855	AL2-stm	0.0280	0.958	3.99	0.04	0.702972	10.94	0.08	21	10.19	2.15	139,5 $\pm$ 32	0.93 $\pm$ 0.037	<b>134,2<math>\pm</math>32</b>

\* = data deriving from  $^{230}\text{Th}/^{234}\text{U}$  ratio.