

Analyzing the recent dynamics of wildland fires in *Quercus suber* L. woodlands in Sardinia (Italy), Corsica (France) and Catalonia (Spain)

Questa è la versione Post print del seguente articolo:

Original

Analyzing the recent dynamics of wildland fires in *Quercus suber* L. woodlands in Sardinia (Italy), Corsica (France) and Catalonia (Spain) / Salis, M.; Arca, B.; Alcasena-Urdiroz, F.; Massaiu, A.; Bacciu, V.; Bosseur, F.; Caramelle, P.; Dettori, S.; Fernandes de Oliveira, A. S.; Molina-Terren, D.; Pellizzaro, G.; Santoni, P. A.; Spano, D.; Vega-Garcia, C.; Duce, P.. - In: EUROPEAN JOURNAL OF FOREST RESEARCH. - ISSN 1612-4669. - 138:3(2019), pp. 415-431. [10.1007/s10342-019-01179-1]

Availability:

This version is available at: 11388/219478 since: 2019-07-11T11:53:51Z

Publisher:

Published

DOI:10.1007/s10342-019-01179-1

Terms of use:

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

Publisher copyright

note finali coverpage

(Article begins on next page)

[Click here to view linked References](#)

1 Analyzing the recent dynamics of wildland fires in *Quercus suber* L. 2 woodlands in Sardinia (Italy), Corsica (France) and Catalonia (Spain)

3 Michele Salis^{1,*}, Bachisio Arca¹, Fermin Alcasena-Urdiroz², Antonella Massaiu³, Valentina Bacciu⁴, Frédéric
4 Bosseur⁵, Philippe Caramelle³, Sandro Dettori⁶, Ana Sofia Fernandes de Oliveira⁶, Domingo Molina-Terren⁷,
5 Grazia Pellizzaro¹, Paul-Antoine Santoni⁵, Donatella Spano^{4,6}, Cristina Vega-Garcia^{2,8}, Pierpaolo Duce¹

6 ¹ National Research Council (CNR), Institute of Biometeorology (IBIMET), Traversa La Crucca 3, 07100,
7 Sassari (Italy); miksalis@ibimet.cnr.it; b.arca@ibimet.cnr.it; g.pellizzaro@ibimet.cnr.it;
8 p.duce@ibimet.cnr.it

9 ² University of Lleida, Department of Agricultural and Forest Engineering, Avenida da Rovira Roure 191,
10 25198 Lleida (Spain); ferminalcasena@eagrof.udl.cat; cvega@eagrof.udl.cat

11 ³ Office National des Forest (ONF), Résidence la Pietrina, Avenue de la Grande Armée, 20000 Ajaccio
12 (France); antonella.massaiu@onf.fr; philippe.caramelle@onf.fr

13 ⁴ Euro-Mediterranean Center on Climate Change (CMCC), IAFES Division, Via De Nicola 9, 07100 Sassari
14 (Italy); valentina.bacciu@cmcc.it; donatella.spano@cmcc.it

15 ⁵ University of Corsica, SPE-UMR 6134 CNRS, Campus Grossetti, B.P. 52, 20250 Corte (France);
16 bosseur@univ-corse.fr; santoni@univ-corse.fr

17 ⁶ University of Sassari, Department of Agriculture, Via De Nicola 9, 07100, Sassari (Italy); sdettori@uniss.it;
18 acortez@uniss.it; spano@uniss.it

19 ⁷ University of Lleida, Department of Crop and Forest Science, Avenida da Rovira Roure 191, 25198 Lleida
20 (Spain); dmolina@pvcf.udl.es

21 ⁸ Forest Sciences Centre of Catalonia (CTFC), Carretera de St. Llorenç de Morunys, km 2, 25280 Solsona
22 (Spain); cvega@eagrof.udl.cat

23 * Corresponding author

24 Abstract

25 Wildland fires represent a major threat to *Quercus suber* L. ecosystems, which provide relevant socio-
26 economic and ecological services in the Mediterranean Basin. In this work, we analyzed recent wildland fire
27 dynamics in cork oak woodlands along the fire-prone areas of Sardinia (Italy), Corsica (France), and Catalonia
28 (Spain). We first characterized geographic extent and main characteristics of cork oak woodlands in these
29 regions and analyzed how environmental (climate and elevation) and socio-economic factors (population and
30 land uses) vary in the areas covered by *Quercus suber* L. We then evaluated how wildfires affected cork oak
31 stands and, by logistic regression analysis, to what extent wildfires in cork oak areas were related to the above
32 set of environmental and anthropic explanatory variables. Results revealed specific variations across study areas
33 in cork oak characteristics as well as in environmental and social factors. We highlighted the spatial and

34 temporal patterns of wildfires on cork oak woodlands, in terms of extent, seasonality, frequency, and main
35 driving factors. In the period 2003-2015, the percentage of cork oak woodlands burned ranged from 3.42% in
36 Corsica to 11.30% in Sardinia. A few large and severe wildfires accounted for most of the area burned in cork
37 oak woodlands. The most significant predictive variable that explained the spatial variation in wildland fire
38 ignitions inside or nearby cork oaks was summer precipitation, while the weight of other factors varied
39 depending on the region. This study provides evidence on recent fire dynamics in cork oak woodlands and gives
40 valuable information and insights for the implementation of forest management and planning strategies in the
41 Mediterranean area.

42 **Keywords**

43 Cork oaks, Mediterranean Basin, fire regime, fire management, land management

44 **1. Introduction**

45 Cork oak (*Quercus suber* L.) is a long-lived (up to hundreds of years) evergreen tree, endemic to the
46 Mediterranean basin (EEA 2007). Cork oak woodlands are mostly concentrated in a few Mediterranean
47 countries and cover about 2.2 million hectares (FAO 2013). The area of *Quercus suber* L. in Europe is estimated
48 to be close to 1.5 million hectares, the majority of which located in the Iberian Peninsula (FAO 2013; Dettori
49 and Filigheddu 2016). The high presence of *Quercus suber* L. in the Mediterranean Basin is linked to its high
50 economic and cultural value for rural communities (Urbieta *et al.* 2008; FAO 2013; Corona *et al.* 2018): cork
51 is the second most important marketable non-wood forest product in the western Mediterranean Basin, with
52 annual world cork market exports close to € 1.45 billion (Aronson *et al.* 2009; Catry *et al.* 2012a; APCOR
53 2016). *Quercus suber* L. stands have been intensively managed, protected and shaped by human activities over
54 millennia for the nutritional values of acorns, to provide shelter and shade for livestock, and for the production
55 of wood for domestic uses. In addition, cork oak forests supply a number of other goods and ecosystem services,
56 including habitats for many animal species and plant endemisms, leisure activities, tourism, beekeeping,
57 aromatic herbs, or edible mushrooms. The social-ecological values that humans have imprinted on cork oak
58 environments in the Mediterranean Basin generated unique cultural landscapes, which in turn influence local
59 and regional culture and identity (Aronson *et al.* 2009; Plieninger *et al.* 2014). The landscape, cultural and
60 ecological value of cork oak areas is also acknowledged in the Natura 2000 Network by the European
61 Commission (DG Environment 2007).

62 As other Mediterranean forest systems, a number of previous studies have reported that cork oak
63 ecosystems are in decline or endangered. The spatial distribution and vitality of *Quercus suber* L. woodlands
64 are affected by several complex and sometimes interrelated disturbance factors (Luciano and Roversi 2001;
65 Dettori and Filigheddu 2003, 2016; Silva and Catry 2006; EEA 2007; Pereira 2007; Acacio *et al.* 2009; Bugalho
66 *et al.* 2011; FAO 2013; Oubrahim *et al.* 2015; Acha and Newing 2015; Oliveira *et al.* 2016; Dettori *et al.* 2018).
67 Wildland fires represent one of the most relevant disturbances for cork oaks, as well as for other forests in the
68 Mediterranean Basin. Nonetheless, it is widely accepted that fires are an integral part of the dynamics of many
69 forest ecosystems, as they play a fundamental role in forest renewal and on insect and disease control (FAO
70 2013, Moritz *et al.* 2014). In Europe, forest fires mainly affect the Southern areas: in the period 2000-2015,
71 about 51,000 fires per year have been recorded, burning approximately 380,000 ha of land (San-Miguel-Ayanz
72 *et al.* 2016). In the Mediterranean Basin, most of the ignitions have anthropogenic origin, while area burned
73 and fire intensity are primarily determined by weather conditions, topography, fuel load and continuity
74 (Romero-Calcerrada *et al.* 2008; Pausas and Paula 2012; Ager *et al.* 2014b; Oliveira *et al.* 2014; Levin *et al.*
75 2016; San-Miguel-Ayanz *et al.* 2016; Turco *et al.* 2017; Rodrigues *et al.* 2018). Climate change is expected to
76 increase future wildfire risk, area burned and the frequency of high severity events all over Southern Europe
77 (Moriondo *et al.* 2006; Lung *et al.* 2013; Bedia *et al.* 2014; Kovats *et al.* 2014; Lozano *et al.* 2017). Furthermore,
78 climate change can exacerbate stress conditions related to droughts, heat waves, pests and diseases, thereby
79 contributing to the potential decline of cork oak woodlands in the next decades (Bergot *et al.* 2004; Luciano *et al.*
80 2005; Catry *et al.* 2012a; Costa *et al.* 2014). The recent trends of forest, agricultural and livestock activity
81 abandonment, as well as the changes in the domestic energetic sources and the afforestation of abandoned lands,
82 are responsible for the increase in highly flammable fuel accumulation and continuity, with potential effects on
83 fire regime (Bonet and Pausas 2007; Baeza *et al.* 2011; Pausas and Fernandez-Munoz 2012; Madrigal *et al.*
84 2016). These factors could further increase the vulnerability to wildland fires of Mediterranean forests in future
85 decades (Moreira *et al.* 2011; Salis *et al.* 2016b; Alcasena *et al.* 2016a).

86 *Quercus suber* L. is considered a fire-resistant and fire-resilient plant, due to its stem and crown
87 resprouting capability (through epicormic buds), even after intense crown fires, and to the thick bark that ensures
88 protection and insulation to the cambium from the heat produced by fires. Therefore, cork oaks are very
89 competitive in the post-fire recovery phases in comparison to several other species (Pausas 1997; Curt and
90 Pausas 2010; Catry *et al.* 2012a). For this reason, summer fires, even induced by landowners, have favored the
91 expansion of cork oaks in Sardinia at the expense of *Quercus ilex* L. and *Quercus pubescens* Willd. (Vogiatzakis
92 and Careddu 2003). However, there is evidence that post-fire cork oak response and recovery capabilities may

93 largely vary depending on several factors such as bark thickness, debarking interval, tree diameter, fire intensity
 94 and spread rate (Pausas 1997; Barberis *et al.* 2003; Moreira *et al.* 2007; Catry *et al.* 2009, 2012b). Moreover,
 95 an important issue for burned cork oaks is that the bark often becomes unsuitable for the most profitable
 96 industrial uses for a long period, which results in important economic losses to forest and rural communities.
 97 Hence, the incidence and effects of wildland fires on cork oak stands in the Mediterranean area are highly
 98 relevant to land managers, planners, and policymakers, according to the strategic economic, ecological and
 99 cultural importance of the *Quercus suber* L. woodlands.

100 In this work, we investigated the recent wildland fire dynamics on cork oak stands in three Mediterranean
 101 regions, namely Sardinia (Italy), Corsica (France), and Catalonia (Spain), where *Quercus suber* L. woodlands
 102 cover approximately 265,000 ha (about 18% of the European cork oak surface area). This research addresses
 103 the limited knowledge on recent wildfire trends in cork oak areas and on the geographic distribution of *Quercus*
 104 *suber* L. in the areas under investigation. The main research questions of this work are: (1) What are the main
 105 geographic, environmental and socio-economic characteristics of the areas covered by *Quercus suber* L.
 106 woodlands? (2) To what extent have wildfires threatened cork oaks? (3) Do environmental and socio-economic
 107 factors explain the historical wildfire incidence in *Quercus suber* L. woodlands? The answers to these questions
 108 provide insights and data on wildland fire dynamics that can support the implementation of forest management
 109 strategies and policies specific for cork oak forests.

110 2. Materials and Methods

111 2.1. Study areas

112 The study was conducted in the fire-prone regions of Sardinia (Italy), Corsica (France), and Catalonia
 113 (Spain), located in the north-western part of the Mediterranean Basin (Fig. 1). For this study, we used “region”
 114 and “province” as reference terms to identify NUTS-2 and NUTS-3 levels (EU Classification of Territorial
 115 Units for Statistics) for the study areas. The main characteristics of these areas are summarized in Table 1.

116 **Table 1.** Summary of the main characteristics of the study areas. The sources of this information are provided
 117 in the next sub-sections

	Sardinia (Italy)	Corsica (France)	Catalonia (Spain)
<i>Size (km²)</i>	24,100	8,700	32,100
<i>Inhabitants</i>	1.65 M	0.33 M	7.50 M
<i>Population</i>	mostly concentrated nearby the main towns of Cagliari and Sassari	mostly concentrated nearby the main towns of Ajaccio and Bastia	mostly concentrated in the metropolitan area of Barcelona and the coastal areas
<i>Climate Type</i>	hot-summer Med.; warm-summer Med. above 750 m a.s.l.	hot-summer Med.; warm-summer Med. above 500 m	hot-summer Med.; warm-summer Med. in NW Catalonia; from cold-summer

		a.s.l.; cold-summer Med. above 1,500 m. a.s.l.	Med. until alpine above 1,200 m a.s.l.
<i>Annual Precipitation</i>	from ~400 mm in the southern coasts to ~1,100 mm in the mountain areas	from ~500 mm in some coastal areas to more than 1,100 mm in the mountain areas	from ~400 mm in the inner areas of Lleida to more than 1,500 mm in the Pyrenees areas
<i>Mean Annual Temperature</i>	from ~7 °C in the mountain areas to ~17 °C along the southern coasts	from ~3 °C at the highest elevations to ~16 °C in the coastal areas	from ~0 °C in the Pyrenees to ~17 °C in the southern coasts




Fig. 1. Digital elevation model (DEM) of the study areas (Catalonia, a; Sardinia and Corsica, b), along with the spatial distribution of *Quercus suber* L. woodlands (green polygons) and the administrative NUTS-3 boundaries (provinces of Cagliari, Sassari, Nuoro and Oristano in Sardinia; provinces of Barcelona, Tarragona, Girona, and Lleida in Catalonia, departments of Corse du Sud and Haute-Corse in Corsica) used for the present study. The perimeters of the wildland fires (red color) that affected the study areas in the period 2003-2015 are also shown.

In the period 2003-2015, the total area burned in the study area was about 218,000 ha in Sardinia, 65,000 ha in Corsica and 61,000 ha in Catalonia; the average area burned per year was about 17,000 ha in Sardinia, 5,000 ha in Corsica, and 4,700 ha in Catalonia (Fig. 1 and Suppl. Fig. 1). The worst fire seasons in terms of total area burned (more than 30,000 ha) were observed in 2003 (Corsica), and in 2007 and 2009 (Sardinia). Overall, in the study period wildland fires affected less than 4% of the provincial areas in Catalonia as well as in Southern Corsica (Suppl. Fig. 1). Northern Corsica and Cagliari provinces were the most affected by fires (respectively 11.5% and 10.4% of the provincial areas in the 13-years study period). The highest provincial area burned by year was observed in Northern Corsica (6.1%) in 2003, Sassari (2.8%) in 2009, and Nuoro (2.3%) in 2007 (Suppl. Fig. 1). A peak in burned area was observed in July for all provinces, except for Tarragona (May), Northern Corsica and Barcelona (August) (Suppl. Fig. 2).

135 2.2. Data compilation

136 2.2.1. Cork oak woodland data

137 The *Quercus suber* L. woodlands data for the study areas were derived from the most accurate spatial
138 input data available at national or regional levels. In Sardinia, the information source was the “Land Use Map
139 of Sardinia” (2008), which is a polygon shapefile that maps the Sardinian land uses with minimum mapping
140 unit of 0.75 ha (www.sardegnaeoportale.it). The information of this database is hierarchically organized
141 following the CORINE Land Cover classification at the fifth level. The codes associated with cork oaks are:
142 2.4.1.3 (Annual crops associated with permanent crops (pastures and grasslands with *Quercus suber* L. cover
143 below 25%) and 3.1.1.2.2 (*Quercus suber* L. forests). The spatial location of cork oak woodlands in Corsica
144 was derived from the “Inventaire Forestier National 2003” (<https://inventaire-forestier.ign.fr/>, minimum
145 mapping unit of 20 ha). The map has three specific categories related to cork oak: *Quercus suber* L. forests,
146 Mediterranean maquis with *Quercus suber* L., mixed *Quercus suber* L. woodlands and coppice. The cork oak
147 information for Catalonia was obtained from the “Pure and Mixed Forest Type Map of Catalonia” (Vericat *et*
148 *al.* 2010, minimum mapping unit of 25 ha). The classes associated to cork oaks, or to areas where cork oaks are
149 the dominant forest types, are the following: a) *Quercus suber* L. forests; b) *Quercus suber* L. and other *Quercus*
150 spp.; c) *Quercus suber* L. and *Pinus* spp.; d) *Quercus suber* L. and other species.

151 2.2.2. Wildland fire data

152 We gathered the perimeters of the wildland fires that affected Sardinia, Corsica, and Catalonia for the
153 study period 2003-2015. These data were provided by regional forest administrations: the Sardinia Forest
154 Service (CFVA), the Office National de Foret (ONF), and the Generalitat de Catalunya (GenCat), for Sardinia,
155 Corsica and Catalonia, respectively. We then merged and, in the case of unclosed polygons, rectified (about
156 0.01% of the data series) the wildland fire perimeters and the associated databases. Furthermore, we collected
157 a longer time series (1990-2015) of wildland fire occurrence to analyze the spatial and temporal patterns of fire
158 ignitions in the study areas. The ignition data of Catalonia were obtained from MAGRAMA (Spanish Ministry
159 of Agriculture, Food and Environment), while for Sardinia and Corsica the data were provided by CFVA and
160 ONF. We first checked the fire databases to evaluate their consistency, and we encountered a certain number
161 of records (about 25%), particularly in the period 1990-2002, where ignition points coordinates were missing
162 or incorrectly reported. In these cases, we randomly distributed the fire ignitions within municipalities. We then
163 used the ignition point database to derive reference raster maps (100-m resolution) of historical ignition point

164 densities for the periods 1990-2002 and 2003-2015. A radius search distance of 5,000 m was set to generate
165 smoothed maps of historical ignition density. This distance has been found significant in other works (i.e. Vega-
166 Garcia *et al.* 1995), and compensates for possible errors in the ignition point coordinates of the historic
167 databases.

168 2.2.3. Other input data

169 Terrain elevation information for the study areas was obtained from the EU-DEM data provided by the
170 EEA (2017b). The EU-DEM is a 25-m resolution digital terrain model (DTM), and covers the whole EU
171 countries plus some neighboring areas. Climate conditions of the study areas and of the cork oak stands were
172 derived from the 1-km resolution “WorldClim version 2” database (Fick and Hijmans 2017), which provides
173 monthly climate data for the period 1970-2000. Population density data at municipality level for the study areas
174 were gathered from the Italian, French and Spanish census data, and refer to the years 1990, 2000 (1999 for
175 Corsica) and 2010. Finally, we used the Corine land cover data (CLC, years 1990, 2000, and 2012, EEA 2017a)
176 to analyze temporal variations in land use patterns for the study areas. The CLC data are seamless vector
177 coverages of land covers for several European countries, with a minimum mapping unit of 25 ha. The CLC
178 classification consists of an inventory of 44 land cover classes, grouped in a three-level hierarchy. In this work,
179 we focused our attention on the following classes: artificial surfaces (codes 111, 112, 121, 122, 123, 124, 131,
180 132, 133, 141, 142), arable land (211, 212), permanent crops (221, 222, 223), heterogeneous agricultural areas
181 (241, 242, 243, 244), forests (311, 312, 313), scrub and/or herbaceous vegetation associations (321, 322, 323,
182 324).

183 *2.3. Data analysis*

184 2.3.1 Geographic extent and environmental and socio-economic conditions in the cork oak areas

185 To characterize *Quercus suber* L. areas, we first mapped the cork oak woodlands and determined their
186 total extent at provincial and regional levels while taking into account elevation classes and climate conditions.
187 For terrain elevation, we used 250-m as reference range to discriminate the cork oak distribution in the study
188 areas as a function of the elevation. We then applied the non-parametric Kruskal-Wallis statistical test to
189 evaluate the significance of differences ($p = 0.05$) among provinces. Regarding climatic conditions, we
190 computed average, minimum and maximum values of temperatures and total precipitation at the provincial level
191 and on the areas covered by cork oak woodlands, using as temporal reference: 1) the months from June to
192 September (which essentially represent the fire season in the majority of the study areas); and 2) the annual

193 averages. We then applied the Kruskal-Wallis test to evaluate the significance of differences ($p = 0.05$) in
194 climatic conditions among provinces. To evaluate demographic and land use variations from 1990 to 2010 in
195 the cork oak zones of the study areas, we selected municipalities characterized by a presence of *Quercus suber*
196 L. woodlands above 5% of the total municipality area (Suppl. Fig. 3, and Fig. 2 and 3).

197 2.3.2 Historical trends in wildfires on cork oak areas

198 We intersected wildland fire perimeters and cork oak woodlands polygons to quantify the hectares of
199 *Quercus suber* L. burned at provincial and regional scales in the period 2003-2015. Moreover, we determined
200 the cork oak area burned by year and by month in the study areas. The role played by the size of historical
201 wildland fires (using as reference three size classes: ≤ 1 ha; 1-10 ha; 10-100 ha; > 100 ha) on cork oak area
202 burned in the period 2003-2015, at the regional and provincial levels, was also investigated. We then evaluated
203 the distance between ignition locations of large wildland fires and *Quercus suber* L. woodlands polygons by
204 using the “Near” command of ArcGIS 10.3. For this purpose, we used 100 ha as a threshold to identify large
205 events. The analysis was carried out considering two reference periods (2003-2015 and 1990-2002) and three
206 reference distance classes (< 1 km, 1-10 km, and > 10 km).

207 2.3.3 Relationships between wildfire occurrence in cork oaks and environmental and socio-economic factors

208 We applied logistic regression to analyze, at the regional scale, the relationships between the 1990-2015
209 wildland fires ignited inside or in the proximity of cork oak woodlands (using a 1-km buffer area around the
210 *Quercus suber* L. polygons) and the following explanatory environmental and socio-economic variables:
211 climate, topography, population density, and land use. We rasterized the explanatory variables using a reference
212 grid of 5-km resolution for the whole study areas and quantified the number of annual fire ignitions for each
213 cell. The logistic regression was used for binary classification (presence/absence) of fire ignitions inside the
214 cork oak grid cells. The estimates of the binary response probability were based on the values of the explanatory
215 variables. The final multivariate models were obtained by using a stepwise approach and considering all
216 possible combinations of the variables for each region. The null hypothesis (the effects of the explanatory
217 variables in the logistic models are equal to zero) was tested by the p-value of the Wald Chi-Square statistics.
218 The Wald statistic was also used to evaluate the significance of both intercept and predictors used in the models.
219 We evaluated the adequacy of the fitted models by the Hosmer and Lemeshow (2000) test, a Pearson Chi-
220 Square statistic that reflects the differences between observed and expected probabilities and provides an
221 estimate of the lack of fit ($p = 0.01$). In addition, the association between observed and predicted data was
222 measured by the c-statistic (Bamber 1975), which is the percentage of pairs with concordant prediction and is

223 commonly used as a standard measure of the predictive accuracy of a logistic regression model. C-statistic
 224 values were used to compare the different models, considering that c values range from 0.5 (random
 225 classification) to 1 (perfect association). We used the overall model accuracy to test the improvement
 226 determined by the use of explanatory variables over the intercept-only model.

227 3. Results

228 3.1. Characterizing geographic extent and environmental and socio-economic conditions of 229 the cork oak areas

230 As shown in Table 2 and Fig. 1, the presence of *Quercus suber* L. woodlands differs at the provincial
 231 level. Sardinia has about 138,400 ha of lands covered by cork oak (5.75% of the regional territory), while in
 232 Corsica the area covered by cork oak woodlands is around 64,500 ha (7.40% of the regional territory) (Table
 233 2). On the other hand, *Quercus suber* L. woodlands encompass only about 1.91% (approximately 61,400 ha) of
 234 Catalonia (Table 2), and are only present in the northeastern zone of this region (Barcelona and Girona
 235 provinces).

236 **Table 2.** Summary of the *Quercus suber* L. woodlands extension (in hectares) for the study areas. The relative
 237 percentage of provincial and regional area is reported under parenthesis. The surface area is rounded to the
 238 nearest ten.

Cagliari	Sassari	Nuoro	Oristano	SARDINIA
9,030	80,430	35,880	13,090	138,430
(1.19%)	(10.46%)	(6.20%)	(4.33%)	(5.75%)
Corse du Sud	Haute-Corse			CORSICA
46,570	17,930			64,500
(10.95%)	(4.01%)			(7.40%)
Barcelona	Girona	Tarragona	Lleida	CATALONIA
7,910	53,480	-	-	61,390
(1.02%)	(9.06%)			(1.91%)

239 In Sardinia, the northern province of Sassari accounts for about 60% of the Sardinian cork oaks woodlands,
 240 while in Corsica *Quercus suber* L. woodlands are mainly present in Corse du Sud (46,570 ha) (Table 2).
 241 Catalonian *Quercus suber* L. woodlands are widespread in the province of Girona, which accounts to about
 242 87% of the overall regional cork oak area (Table 2). Considering the distribution of cork oaks at the provincial
 243 level, Corse du Sud, Sassari and Girona show the highest presence of *Quercus suber* L. woodlands, which cover
 244 more than 9% of the provincial area. Cork oak woodlands account for 4-6% of the provincial territory in Nuoro,
 245 Oristano, and Haute-Corse, while their incidence in the other provinces is very limited (Cagliari and Barcelona)
 246 or null (Tarragona and Lleida) (Table 2).

247 Concerning the analysis of the cork oak stands at different elevation classes, the study areas show
 248 statistically significant differences (Table 3). *Quercus suber* L. woodlands are located at elevations much above

249 the average provincial elevation in Cagliari and Oristano (376 m vs. 227 m a.s.l. and 350 m vs. 210 m a.s.l.,
 250 respectively), while in Corsica and Catalonia they grow in areas below or largely below the average provincial
 251 elevation. In Corsica, more than 80% of *Quercus suber* L. woodlands is located in the 0-250 m elevation class
 252 (Table 3). Also in Catalonia the presence of cork oaks is most important at 0-250 m elevation class (about 60%
 253 of the total area), while only 33% of the Catalonia cork oak woodlands is found at 250-500 m altitude. The
 254 presence of cork oaks at elevations above 500 m is less than 5% of the regional and provincial areas of Catalonia
 255 and Corsica (Table 3). In Sardinia, about 50% of the *Quercus suber* L. stands are located at 250-500 m altitude
 256 class, with a relevant presence also at the 500-750 m elevation class (26% vs. less than 3% for Catalonia and
 257 Corsica). In the provinces of Nuoro and Cagliari, cork oaks woodlands at elevations above 500 m represent
 258 about 55% and 40% of the provincial cork oak area, respectively. There are no significant statistical differences
 259 in average elevation of cork oak woodlands between Catalonian provinces, as well as for Cagliari, Oristano,
 260 and Sassari provinces (Table 3).

261 **Table 3.** Average terrain elevation (ELEV), average elevation of cork oak woodlands (ELEV QS), and
 262 percentage distribution of *Quercus suber* L. woodlands at different terrain elevation classes (0-250; 250-500;
 263 500-750; >750 m a.s.l.) in the provinces of the study areas. The Kruskal–Wallis test ($p = 0.05$) was performed
 264 to evaluate statistical differences in average elevation and average elevation of cork oak woodlands among
 265 provinces. Different letters in the same column indicate significant differences among provinces at $p = 0.05$.

PROVINCE	ELEV (m)	ELEV QS (m)	0-250	250-500	500-750	>750
Cagliari	227.1f	376.3b	24.08	36.54	38.84	0.54
Nuoro	526.1b	504.4a	12.32	32.57	41.37	13.74
Oristano	210.9g	350.0b	14.08	75.49	10.36	0.07
Sassari	325.7e	394.7b	22.39	55.33	20.00	2.28
Corse du Sud	560.4a	159.4e	87.51	12.13	0.36	0.00
Haute-Corse	592.3a	201.7d	69.07	28.30	2.48	0.15
Barcelona	518.1c	251.2c	68.78	29.14	2.07	0.00
Girona	517.2d	243.6c	60.08	34.05	5.16	0.71


266 The average climatic conditions of *Quercus suber* L. woodlands at the provincial level are provided in
 267 Table 4. Overall, the climate of these areas is quite similar, particularly at the annual time lag (i.e.: average
 268 temperatures range approximately from 14.9 °C in Oristano to 14.1 °C in Nuoro). When taking into account
 269 only the June-September period, the ranges of some climatic variables in the *Quercus suber* L. woodlands show
 270 to be larger, particularly the average accumulated precipitation (i.e.: from 76.4 mm in Cagliari to 232.9 mm in
 271 Barcelona) and the average maximum temperature (from 26.1 °C in Corse du Sud to 28.9 °C in Oristano) (Table
 272 4). The maps of average maximum temperatures and accumulated precipitation observed in the study areas
 273 during the period June-September are presented in the Suppl. Fig. 4 and 5.

274 **Table 4.** Average mean temperature (TAVG, in °C), minimum temperature (Tm, in °C), maximum temperature
 275 (TM, in °C), and accumulated precipitation (PP, in mm), at the annual level and for the period June-September,
 276 in the areas covered by *Quercus suber* L. woodlands. The results are summarized at the provincial level.

277 Climate data were obtained from the 1-km resolution “WorldClim version 2” data (Fick and Hijmans 2017).
 278 The Kruskal–Wallis test ($p = 0.05$) was performed to evaluate statistical differences in climate conditions of
 279 cork oak woodlands among provinces. Different letters in the same column indicate significant differences at p
 280 = 0.05.

PROVINCE	YEAR				JUNE-SEPT			
	<i>T</i>	<i>Tm</i>	<i>TM</i>	<i>PP</i>	<i>T</i>	<i>Tm</i>	<i>TM</i>	<i>PP</i>
Cagliari	14.8b	10.6e	19.0b	624.2e	21.6b	16.5d	28.6b	76.4f
Nuoro	14.1e	10.3f	17.9g	716.4b	21.1d	16.4e	27.8c	90.4d
Oristano	14.9a	10.8d	19.1a	728.0b	21.8a	16.7c	28.9a	83.9e
Sassari	14.5c	10.8d	18.3d	703.5c	21.3c	16.7c	27.8c	88.1d
Corse du Sud	14.9a	11.8a	18.1e	612.7e	21.0e	17.4b	26.1f	91.8d
Haute-Corse	14.5d	11.0c	18.0f	663.6d	20.7f	16.7c	26.3e	113.1c
Barcelona	14.9ab	11.4b	18.4c	761.4a	21.6b	17.8a	26.9d	232.9a
Girona	14.7bc	11.0c	18.4cd	726.7b	21.3c	17.3b	26.9d	214.1b

281 The variations in population density in the municipalities where the presence of *Quercus suber* L.
 282 woodlands is above 5% of the municipality area are reported in Fig. 2, which is summarized at the regional
 283 level. Overall, the population in Corsica and Catalonia increased between 1990 and 2010 by 22% and 39%,
 284 respectively, with the highest variation observed in the province of Barcelona (+48%). On the contrary, in the
 285 same period we observed a decrease of about 5% of the inhabitants in the main cork oak areas of Sardinia,
 286 particularly in the inner lands. In addition, while in Catalonia and Corsica the percent change in population from
 287 1990 to 2010 was quite uniform at the provincial level, Sardinia showed uneven changes: the province of Sassari
 288 exhibited slight changes and average variations close to zero, while the decrease in population in the main cork
 289 oak areas was over 11% in Cagliari and Oristano, and close to 10% in Nuoro.



290

291 **Fig. 2.** Map of the percent change in population between 1990 and 2010 in the study areas. The histograms
 292 report the number of inhabitants in the municipalities characterized by a presence of *Quercus suber* L.
 293 woodlands above 5% of the municipality area. The population data refer to census data of 1990, 2000, and
 294 2010.

295 As for changes in main land uses between 1990 and 2012 in the municipalities characterized by a presence
 296 of *Quercus suber* L. woodlands above 5% of the municipality area (Fig. 3), we observed an overall increase in
 297 the surface areas covered by forests (particularly relevant in Catalonia) and artificial surfaces. Scrubs and
 298 herbaceous vegetation decreased in the study areas, with particular relevance in Catalonia and Sardinia.
 299 Concerning the other land uses, the variations were more complex and did not follow a common trend (Fig. 3).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65





Fig. 3. Map of the main land covers of the study areas according to Corine Land Cover 2012 (CLC2012). The histograms report the percent variations between CLC2012 and CLC1990 of the main land covers for the municipalities characterized by a presence of *Quercus suber* L. woodlands above 5% of the municipality area. A.S. = artificial surfaces; A.L. = arable land; H.A. = heterogeneous agricultural areas; P.C. = permanent crops; S.&H. = scrubs and herbs association; F. = forests; O. = other land covers.

3.2. Evaluating the wildfire trends in cork oak areas

In the study period 2003-2015, the total cork oak woodland area burned in the regions under investigation was around 23,500 ha (Fig. 1). Sardinia accounted for about 66% of the total cork oak area burned. In this island, on average about 1,200 ha of *Quercus suber* L. stands (about 0.87% of the regional cork oak area) were burned each year (Fig. 4). In the 13-years study period, the total Sardinia cork oak area burned was about 15,500 ha. The worst seasons for *Quercus suber* L. woodlands were observed in 2007 and 2009, with about 2.79% (\approx 3,850 ha) and 3.23% (\approx 4,450 ha), respectively, of the total cork oak area burned. In Corsica (Fig. 4), only 3.42% (\approx 2,200 ha) of the regional area covered by *Quercus suber* L. stands was burned in the study period. This means that, on average, the annual cork oak area burned was about 165 ha, that is 0.26% of the total Corsica cork oak area. Furthermore, the annual percentage of *Quercus suber* L. woodlands burned over the total area covered by cork oaks was always less than 1%. In Catalonia, about 9.38% (\approx 5,700 ha) of the area covered by *Quercus suber* L. woodlands was burned, with an annual average of 0.72% (about 450 ha) (Fig. 4). Nearly 7.5% (\approx 4,400 ha) of the Catalonia *Quercus suber* L. woodland area was burned in 2012 and was almost entirely

319 concentrated in the province of Girona (Fig. 4). Overall, the combination of wildland fire and cork oak woodland
 320 shapefiles in Sardinia, Corsica and Catalonia revealed that the *Quercus suber* L. areas which burned more than
 321 once in the period 2003-2015 were limited and ranged from 0 to less than 1% of the total cork oak area burned
 322 at the provincial level.



323
 324 **Fig. 4.** *Quercus suber* L. woodlands area affected by wildland fires vs. regional cork oak area for Catalonia,
 325 Corsica and Sardinia, for the study period 2003-2015.

326 A small number (< 2%) of large fire events (> 100 ha) accounted for most of the total cork oak area burned
 327 at the regional scale. The role played by large fires in *Quercus suber* L. woodlands burned was very significant
 328 in Catalonia (about 99% of the total cork oak burned determined by large events), while in Corsica and Sardinia
 329 they accounted for about 79% and 72%, respectively, of the total *Quercus suber* L. area burned (Table 5). This
 330 analysis highlighted some relevant differences also at the provincial level. For instance, the most relevant
 331 variation in the role played by the different fire size class was observed in Sardinian provinces. In fact, while in
 332 Sassari and Nuoro provinces the large fires (> 100 ha) accounted for more than 75% of the total cork oak area
 333 burned, in southern Sardinia these events were responsible for about 31% of the *Quercus suber* L. burned, and
 334 the highest contribution to cork oak area burned was due to the fire size class 10-100 ha (Table 5). Corsica also
 335 presented marked differences: large events accounted for about 88% of the total *Quercus suber* L. woodlands
 336 burned in Haute Corse, and for about 64% in Corse du Sud. In Catalonia, in both Girona and Barcelona
 337 provinces, the fire size class >100 ha caused more than 98% of the total cork oak area burned. A single forest
 338 fire event (Suppl. Fig. 6a) played a key role in determining the cork oak losses in Catalonia and particularly in

339 Girona province. Specifically, on July 2012, the largest Catalonian wildland fire (La Jonquera, about 10,500
 340 ha) of the period 2003-2015 burned approximately 4,100 ha of *Quercus suber* L. woodlands, corresponding to
 341 about 94% of the total cork oak area burned in that year, and to about 72% of the total cork oak area burned in
 342 the region over the 13 years of reference. Another example of the significant role played by large events is
 343 represented by the wildland fire of Nuoro (Suppl. Fig. 6b) in Sardinia. This event (about 9,000 ha), which was
 344 the largest Sardinian wildland fire of the summer season 2007, burned approximately 2,600 ha of *Quercus suber*
 345 L. woodlands, which represent about 67% and 17% of the total cork oak area burned in 2007 and during the
 346 study period, respectively.

347 **Table 5.** Percentage of *Quercus suber* L. woodlands burned by wildland fire size class (≤ 1 ha; 1-10 ha; 10-
 348 100 ha; > 100 ha), for the period 2003-2015, at the regional and provincial levels.

Province / Region	Percentage of QS burned by wildfire size class				TOTAL
	≤ 1 ha	1-10 ha	10-100 ha	>100 ha	
Cagliari	2.75	18.17	48.21	30.86	100.00
Nuoro	0.49	7.36	15.15	77.01	100.00
Oristano	0.78	14.56	20.80	63.85	100.00
Sassari	0.61	8.21	17.13	74.05	100.00
Sardinia	0.67	8.94	17.94	72.45	100.00
Corse du Sud	1.17	5.13	29.36	64.34	100.00
Haute Corse	0.84	4.76	6.08	88.31	100.00
Corsica	0.97	4.91	15.41	78.71	100.00
Barcelona	0.00	0.00	0.00	100.00	100.00
Girona	0.00	0.00	1.32	98.68	100.00
Catalonia	0.00	0.00	1.28	98.72	100.00

349 In Catalonia and Sardinia, *Quercus suber* L. woodlands were mostly affected by fires in July (more than
 350 65% of the total cork oak area), followed by August (with about 13% and 19% of the total cork oak area burned
 351 in Corsica and Sardinia, respectively) (Fig. 5). The cork oak area burned in July was about 10,550 ha in Sardinia
 352 and 4,200 ha in Catalonia (concentrated in the province of Girona). However, in Sardinia, relevant differences
 353 in the monthly distribution of the cork oak burned were observed: in the northern provinces of Sassari and
 354 Nuoro, about 90% of the cork oak stands burned in July and August, while in the other two provinces the
 355 wildfires occurred in these months accounted for about 75% of the total *Quercus suber* L. area burned. Also in
 356 Corsica, July was the month with the highest percentage (about 39%) of *Quercus suber* L. woodlands burned,
 357 but a relevant percentage (about 22%) of the cork oak woodlands burned was also observed in August and
 358 October (Fig. 5). The Corsican provinces presented large differences in the monthly distribution of the cork oak
 359 burned: in Corse du Sud, about 73% of the total cork oak area burned was concentrated in July, while in Haute
 360 Corse the peak (about 87%) was observed in August-October.




Fig. 5. Percentage of the monthly distribution of the *Quercus suber* L. woodlands area affected by fires in Catalonia, Corsica, and Sardinia, for the study period 2003-2015.

The relation between the wildland fire ignition locations and the distance to cork oak stands was analyzed considering the reference period 2003-2015 and the previous 13-year timeframe (1990-2002). Fig. 6 highlights the spatial variations in large fire event (> 100 ha) location and ignition point densities between the two periods. On the whole, the fire ignition point density dropped from 1990-2002 to 2003-2015 for most of the study areas, the main exception being south-western Sardinia (Fig. 6), which showed more areas with peaks above 8 fire ignitions km⁻² in the period 2003-2015. Moreover, large fire (>100 ha) number and area burned decreased considerably in the study areas moving from the first to the second period (Fig. 6). For instance, the number of large fires dropped to 289 events (vs. 541 in the period 1990-2002) in Sardinia, to 47 (vs. 113) in Corsica, and to 75 (vs. 116) in Catalonia. The analysis of the percentage of fire ignition locations as function of the distance from cork oak woodlands and provinces for both 1990-2002 and 2003-2015 revealed that from the first to the second period, wildland fires ignited in the vicinity (< 1 km) of cork oak woodlands areas tended to decrease, except in the province of Girona (Suppl. Table 1). In the period 2003-2015, the likelihood of fire ignitions at the distance class 0-1 km from *Quercus suber* L. woodlands was over 40% of the total ignitions in the provinces of Girona, Nuoro, and Sassari, while Cagliari and Barcelona provinces showed values below 15%. The province of Barcelona presented the largest distances between fire ignitions and cork oak areas (Suppl. Table 1).




Fig. 6. Maps of the historic ignition point density (in # / km²) of the periods 1990-2002 (left) and 2003-2015 (right) for the study areas. The ignition point density was calculated using the Point Density tool of ArcGIS (ESRI) and a radius search distance of 5,000 m. The black triangles indicate the ignition point location of the large wildland fire events (>100 ha) that affected the study areas for the above periods.

3.3. Analyzing the relationships between wildfire occurrence in cork oak areas and environmental and socio-economic factors

The driving variables that mostly affected wildland fire ignitions inside and nearby (1-km buffer) cork oak areas at the regional scale were identified by a set of logistic regression models (Table 6). The Catalonia model was not statistically significant according to the Wald test, while the Sardinia and Corsica models resulted highly statistically significant ($p = 0.01$). However, the Corsica and Sardinia models exhibited a significant lack of fit according to the Hosmer & Lemeshow test (Table 6). Overall, the models showed values of correct classification between 55% and 73%, with the highest values observed in Corsica. C-statistic values range from 0.73 (Sardinia model) to 0.79 (Corsica and Catalonia models), and thus indicated an overall good predictive accuracy of the logistic models (Table 6). In the study areas, the most significant predictive variable that explained the spatial variation of wildland fire ignitions inside or nearby cork oaks was summer precipitation. In more detail, for both Sardinia and Corsica, the most significant predictive variables of the logistic models were land use classes (Anthropic and Agricultural Areas), population density, elevation and summer precipitation (Table 6). As far as climatic conditions are concerned, we also found a major effect of summer maximum temperatures in Sardinia and summer average temperatures in Corsica. Slope played a significant role for the Sardinia model. In Catalonia, the spatial variation of fire ignitions inside or nearby cork oak areas was mainly explained by climatic conditions, namely summer precipitation and annual precipitation (Table 6).

Table 6. Significance of the variables that affected wildland fire ignitions inside and nearby (1-km buffer) cork oak areas as provided by logistic regression models, considering the study period 1990-2015. The values of the coefficient estimates, the Wald Chi-Square values and the significance of the Chi-Square statistics are reported for each predictive variable. Regarding land uses, the main land use classes are reported if significant. ELEV = elevation; SLO = slope; TMAXS = average summer maximum temperatures; TMS = average summer mean temperatures; PRECS = average summer precipitation; PREC = average annual precipitation; POP =

407 population density; LU = land uses. WCF = Wald test coefficient; HLCF = Hosmer & Lemeshow test
 408 coefficient; CORR = percentage of correct estimates; c = c-statistic value. *** = $p < 0.01$; ** = $p < 0.05$; n.s.
 409 = not significant

		Sardinia	Corsica	Catalonia
Predictive variables	ELEV	-0.005 (190.7)***	0.03 (43.04)***	n.s.
	SLO	0.06 (10.32)**	n.s.	n.s.
	TMAXS	-1.22 (470.5)***	n.s.	n.s.
	TMS	n.s.	7.58 (43.04)***	n.s.
	PRECS	0.01 (5.65)***	3.62 (-0.03)***	-0.03 (3.00)***
	PREC	n.s.	n.s.	-0.11 (3.91)**
	POP	0.006 (83.41)***	-0.005 (9.95)***	n.s.
	LU	Agricultural -0.01*** Anthropic -0.33 (30.6)***	Agricultural -2.29*** Anthropic 4.11 (10.46)***	n.s.
Model performance parameters	WCF	862***	122***	11 n.s.
	HLCF	35.95***	33.20***	16.91 n.s.
	CORR	67.2	73.0	55.4
	c	0.73	0.79	0.79

410

411 4. Discussion and Conclusions

412 The main goal of this work was to analyze the recent wildland fire dynamics on cork oak areas in Sardinia
 413 (Italy), Corsica (France), and Catalonia (Spain). We aimed to provide answers to the following research
 414 questions: (1) What are the main geographic, environmental and socio-economic characteristics of the areas
 415 covered by *Quercus suber* L. woodlands? (2) To what extent have wildfires affected cork oaks? (3) Do
 416 environmental and socio-economic factors explain the historical wildfire incidence in *Quercus suber* L.
 417 woodlands?

418 The study areas encompass about 265,000 ha of *Quercus suber* L. woodlands, about 18% of the European
 419 cork oak surface area. Our study showed relevant differences in the spatial distribution of cork oaks in the
 420 regions under investigation. The presence of cork oak stands is mainly concentrated in the provincial areas of
 421 Corse du Sud, Sassari and Girona: these provinces account for more than 50% of the regional *Quercus suber* L.
 422 woodlands. Previous studies (Dettori and Filigheddu 2003; Pereira 2007; FAO 2013) have pointed out that the
 423 differences in cork oak extent within provinces and regions depend upon several interrelated factors, among
 424 which the most relevant are climatic conditions, income and jobs related to the cork industry, pressure of
 425 pastoral and agricultural activities, and urban sprawl. Our work showed that the climatic conditions in the cork
 426 oak woodlands of the study areas were relatively similar, but for summer precipitation and summer average
 427 maximum temperatures. This could explain the high presence of cork oak woodlands at elevation classes > 500
 428 m a.s.l. in Sardinia, while in Corsica and Catalonia *Quercus suber* L. areas are largely concentrated at the lowest

429 elevation classes (< 250 m a.s.l.). Differences in climatic conditions would also explain why Sardinian cork
1 430 oaks are located at higher elevations than in Corsica and Catalonia. Drought conditions and the increasing extent
2
3 431 and role played by agricultural expansion could further explain the limited presence of cork oaks in southern
4
5 432 Sardinia, or their absence in Tarragona and Lleida provinces.
6
7

8 433 The variation in population density between 1990 and 2010 assumed distinct patterns across regions.
9
10 434 While at the regional level most Corsican and Catalan municipalities presented, save from a few localized
11
12 435 inner areas, a general increase in population density, Sardinia showed a large decline of population density in
13
14 436 municipalities far from coastlands and main towns. The population density decline observed in the cork oak
15
16 437 areas of Sardinia can be related to a combination of socio-economic issues, including unemployment,
17
18 438 abandonment of rural and forest activities, ageing population, low birthrates, distance to main services and
19
20 439 towns, and low incomes, which led to emigration of young people towards coastal areas and towns, or even
21
22 440 outside, as observed in other Mediterranean areas (Romero-Calcerrada and Perry 2004; Figueredo and Raschi
23
24 441 2011; Kerckhof *et al.* 2016; Forleo *et al.* 2017). For land use variation from 1990 to 2012, we observed an
25
26 442 increase in forests and artificial surfaces and a reduction in shrublands in the municipalities covered by large
27
28 443 extensions of cork oak woodlands for Sardinia, Corsica and Catalonia. The highest increase in forest areas
29
30 444 (about 15%) in the vicinity of cork oak polygons was observed in Catalonia, followed by Sardinia and Corsica
31
32 445 (about 2-4%). These variations suggest that cork oaks or other forest types are gradually occupying areas
33
34 446 previously covered by shrublands and marginal rangelands, with a potential increase in fuel load and continuity,
35
36 447 particularly in the zones where population density is decreasing and vegetation succession is leading to scrub
37
38 448 encroachment and new forest development (Arianotsou 2001; Scozzafava and De Sanctis 2006; Moreira *et al.*
39
40 449 2011). In addition, the increased presence of artificial surfaces in the municipalities covered by large areas of
41
42 450 cork oak woodlands indicates an expansion of urban interfaces in the wildlands.
43
44

45 451 We found that in the study areas approximately 23,500 ha of cork oak stands (about 9% of the total
46
47 452 *Quercus suber* L. woodlands) were affected by wildland fires in the period 2003-2015. Sardinia accounted for
48
49 453 about 66% of the cork oak woodland area burned. Evident interannual variability in the cork oak area burned
50
51 454 was observed. Overall, the percentage of cork oak woodlands burned at least once ranged from 3.42% in Corsica
52
53 455 to 11.30% in Sardinia for the study period 2003-2015. The highest cork oak area burned observed in Sardinia
54
55 456 could be partially explained by the higher presence of herbaceous fuels and dehesas, which can favor the quick
56
57 457 spread of large wildland fires under strong wind conditions (Salis *et al.* 2018). The role played by environmental
58
59 458 factors on burned area in the Mediterranean Basin has been explored in previous works. There is evidence that
60
61 459 a small number of large fires spreading with extreme weather conditions is responsible for the majority of the
62
63
64
65

460 area burned during a fire season, and that containment activities have minor effectiveness on fire growth during
1
2 461 these events (Pereira *et al.* 2005; Koutsias *et al.* 2012; San Miguel-Ayanz *et al.* 2013; Alcasena *et al.* 2016b;
3
4 462 Salis *et al.* 2016a). Large wildland fires often exhibit extreme behavior (i.e., high intensity, crown fire activity
5
6 463 and spotting fire emission) and can produce damaging impacts on human life, assets and activities as well as
7
8 464 severe disturbances to forest ecosystems. Our work confirms that wildland fires greater than 100 ha, which
9
10 465 accounted for 1-2% of the total number of wildland fires, were associated to most of the area burned in *Quercus*
11
12 466 *suber* L. woodlands in the study areas, except for southern Sardinia. For instance, in the study period, about
13
14 467 70% of the cork oak woodlands in Catalonia burned in a single fire event (La Jonquera, final size about 10,000
15
16 468 ha), and more than 90% of the La Jonquera fire area burned at high or very high severity (severicat.ctfc.cat).
17
18 469 Similar findings were observed in Sardinia during the 2007 and 2009 fire seasons when extreme weather
19
20 470 conditions and simultaneous wildland fires concentrated in a few days overwhelmed fire suppression
21
22 471 capabilities and burned large portions of cork oak areas in the provinces of Nuoro and Sassari.

23
24 472 The present study also reveals that both the number of fire events and the area burned by wildland fires
25
26 473 bigger than 100 ha, which are the primary drivers of fire regime in cork oak landscapes in the study areas,
27
28 474 decreased considerably from 1990-2002 to 2003-2015, and that overall the distance between fire ignitions and
29
30 475 cork oak woodlands areas increased. Furthermore, we observed an overall reduction of fire ignitions from 1990-
31
32 476 2002 to 2003-2015 for almost the entire study areas, the main exception being in southern and western Sardinia.
33
34 477 This is in line with previous studies that analyzed wildland fire trends in the Mediterranean Basin (Turco *et al.*
35
36 478 2016; Jimenez-Ruano *et al.* 2017; Curt and Frejaville 2018). The fact that large wildland fires impacting cork
37
38 479 oak areas dropped in recent years can be explained by the improvements in fire monitoring and suppression
39
40 480 capabilities, by the reduction in fire ignitions related to agro-pastoral causes, and by a general shift of both
41
42 481 people and fire ignitions towards coastal and main residential areas (Viedma *et al.* 2006; Chas-Amil *et al.* 2013;
43
44 482 Salis *et al.* 2014; Curt and Frejaville 2018). Some studies emphasized that fire exclusion policies can
45
46 483 substantially increase the vulnerability of many dry forests to uncharacteristically extensive and severe
47
48 484 perturbations, particularly from wildfire and insects/disease, due to their emphasis on short-term outcomes
49
50 485 (reducing annual area burned and fire ignitions) vs. long-term goals (promoting more resilient landscapes and
51
52 486 forests) (Piñol *et al.* 2005, 2007; Calkin *et al.* 2015; Stephens *et al.* 2016; Camia *et al.* 2017; Curt and Frejaville
53
54 487 2018; Thompson *et al.* 2018). These issues pose severe threats for fire management in future years in the study
55
56 488 areas, since the reduction in forest management activities, in conjunction with the limited occurrence of small-
57
58 489 size and low severity fires in cork oak areas, and the decrease in population in inner areas such as seen in
59
60 490 Sardinia, could promote the accumulation of unmanaged high-load fuels, the expansion of ladder fuels, and the
61
62
63
64
65

491 increase in fuel connectivity. This can lead to more fire-prone landscapes in the Mediterranean area, to a
1 492 potential increase in the likelihood of extreme behavior and severe wildland fire events, thus able to spread
2 493 further and increase in size, and to more limitations for effective fire suppression of aerial and terrestrial forces
3 494 (Moreno *et al.* 1998; Pausas and Vallejo 1999; Pausas and Fernandez-Munoz 2012; Salis *et al.* 2013; Ferreira-
4 495 Leite *et al.* 2016). Therefore, it is likely that, in future years, wildland fires affecting forest areas will start
5 496 increasingly far from the woodlands and will impact the forest stands after spreading for long distances, thus
6 497 resulting in an overall increase of the so-called mega-fires, as dramatically observed in the European summer
7 498 season 2017 (San-Miguel-Ayaz *et al.* 2013; Viegas *et al.* 2017). High-intensity wildland fires could ultimately
8 499 trigger substantial changes on cork oak forest structures from old-growth forests with a commercial interest to
9 500 post-fire shrubby coppices, which require long-time and expensive restoration efforts. Consequently, specific
10 501 management strategies including fuel treatments should be planned and carried out by policy makers and land
11 502 managers to reduce potential losses from future wildland fires in the Mediterranean Basin, in order to preserve
12 503 important natural ecosystem heritages like the cork oak forests. A number of works highlighted that surface fuel
13 504 reduction, silvicultural treatments, and better-planned land mosaics could significantly decrease fire risk in cork
14 505 oak areas (Baeza *et al.* 2006; Santana *et al.* 2011; Catry *et al.* 2012b; Schaffhauser *et al.* 2012; Curt *et al.* 2013;
15 506 Salis *et al.* 2016b, 2018). This brings to light the need for integrating large fire spread analysis and fire risk
16 507 transmission with the design of appropriate wildfire management strategies, particularly for protecting highly
17 508 valued areas or habitats of endangered species (Ager *et al.* 2014a, 2017; Alcasena *et al.* 2017, 2018a; Thompson
18 509 *et al.* 2017; Salis *et al.* 2016b, 2018). Prevention and protection strategies may also involve prescribed burning
19 510 action and may allow the spread of low-intensity wildland fires when weather conditions are mild, in agreement
20 511 with the proactive concept of integrated fire management (Molina *et al.* 2010; Fernandes *et al.* 2013; Montiel
21 512 Molina and Galiana-Martín 2016; Alcasena *et al.* 2018b).

45 513 We observed that the highest cork oak area burned was concentrated during the months of July and August,
46 514 which in Catalonia and Sardinia accounted for almost 80% of the total *Quercus suber* L. area affected by fires.
47 515 Summer months are typically the hottest and driest of the year in the Mediterranean Basin (Flohn and Fantechi
48 516 1984). This confirms the key role of drought conditions, and particularly of dead and live fuel moisture, in the
49 517 dynamics of fire regimes in Mediterranean Europe (Piñol *et al.* 1998; Moreira *et al.* 2011; Turco *et al.* 2017).
50 518 From this point of view, the logistic regression analysis performed in the study areas demonstrated that climatic
51 519 factors, namely summer and annual precipitation, and maximum and average summer temperatures, were
52 520 significantly related to wildland fire ignitions nearby cork oak woodlands. Furthermore, summer precipitation
53 521 was a key driving factor in determining wildfires nearby cork oaks for the three study areas. The fact that
54
55
56
57
58
59
60
61
62
63
64
65

522 wildfires occurring in dry years tend to burn larger areas than in wet years was also observed in other European
1
2 523 fire-prone areas, as for instance in the Valencia region of Spain (Pausas 2004), in Portugal (Viegas and Viegas
3
4 524 1994), and in Greece (Dimitrakopoulos *et al.* 2011; Xystrakis *et al.* 2014). Moreover, our study indicates that,
5
6 525 while in Catalonia fire ignitions in the vicinity of cork oaks are mainly explained by summer and annual
7
8 526 precipitation, Corsica and Sardinia showed more complex patterns: in these regions, elevation and land use play
9
10 527 a significant role, together with population density and some climate variables. These results support the fact
11
12 528 that, even if wildfire ignitions in Mediterranean areas are mostly determined by anthropic factors, a significant
13
14 529 component of the variability in wildfire occurrence is explained by climatic conditions (Pausas 2004, Pereira *et*
15
16 530 *al.* 2013). For these reasons, predicted climate change is likely to affect future fire dynamics in *Quercus suber*
17
18 531 L. woodlands. With a higher frequency of extreme weather conditions, fire-prone landscapes can be
19
20 532 increasingly vulnerable and able to sustain high-intensity and uncontrollable fire events, which can negatively
21
22 533 affect cork oak postfire recovery (Fernandes and Rego 1996; Baeza *et al.* 2006; Pausas and Paula 2012;
23
24 534 Schaffhauser *et al.* 2012). In addition, under the predicted changes in climate conditions, cork oak ecosystems
25
26 535 could be potentially replaced by more xeric species and pyrophytic shrublands and could shift northward or to
27
28 536 higher elevations (Costa and Madeira 2011; Costa *et al.* 2014; Ibáñez *et al.* 2017; Varela 2017).

30
31 537 In conclusion, the cork oak woodlands of the study areas represent an important portion of the European
32
33 538 *Quercus suber* L. ecosystems and a significant ecological, cultural, and economic heritage for rural
34
35 539 communities. In the study period 2003-2015, about 9% of the cork oak woodlands of the study areas have been
36
37 540 affected by wildland fires. Though this represents a relatively small percentage of the total *Quercus suber* L.
38
39 541 surface area, cork oaks burned can lead to relevant losses of ecosystem services and income, as well as to the
40
41 542 death of monumental and ancient trees, which are key components of cultural landscapes and agro-pastoral
42
43 543 areas in Sardinia, Corsica and Catalonia. Wildfires nearby cork oak forests were primary linked to the
44
45 544 occurrence of dry summers, and the most of the *Quercus suber* L. area burned was related to a few large fire
46
47 545 events associated to extreme weather conditions. It is thus likely that the future changes in climate will further
48
49 546 increase wildfire risk for Mediterranean forests, unless significant efforts are made to promote appropriate forest
50
51 547 management and planning strategies. Socio-economic changes can even foster the occurrence of high-intensity
52
53 548 wildland fires, due to the magnitude of land use changes and the abandonment of forest and rural activities.

549 **Acknowledgments**

55
56
57
58
59 550 This work was supported by the Sardinia Region under the project "Multifunzionalità delle foreste a quercia da
60
61 551 sughero in Sardegna" (Multifunctionality of cork oak forests in Sardinia) funded within L.R. n. 7/2007
62
63
64
65

552 "Promozione della ricerca scientifica e della innovazione tecnologica in Sardegna, / Tender no. 2", and by the
 553 "Proterina-2" Project funded by the Italia Francia-Marittimo Programme. We would like to thank the Sardinia
 554 Forest Service, the Office Nationale de Foret, the Generalitat de Catalunya and the Area of Defense against
 555 Forest Fires (ADIF) of the Spanish Ministry of Agriculture, Food and the Environment (MAGRAMA) for
 556 providing fire data and information for Sardinia, Corsica and Catalonia. We also thank the anonymous reviewers
 557 whose comments helped improve this manuscript.

558 **References**

- 559 Acácio V, Holmgren M, Rego F, Moreira F, Mohren GM (2009). Are Drought and Wildfires Turning
 560 Mediterranean Cork Oak Forests into Persistent Shrublands? *Agroforestry Systems* 76, 389–400
- 561 Acha A, Newing HS (2015) Cork Oak Landscapes, Promised or Compromised Lands? A Case Study of a
 562 Traditional Cultural Landscape in Southern Spain. *Human Ecology* 4(43), 601-611
- 563 Ager AA, Day M, Finney MA, Vance-Borland K, Vaillant NM (2014a) Analyzing the transmission of
 564 wildfire exposure on a fire-prone landscape in Oregon, USA. *Forest Ecology and Management* 334, 377-390
- 565 Ager AA, Evers CR, Day MA, Preisler HK, Barros AMG, Nielsen-Pincus M (2017) Network analysis of
 566 wildfire transmission and implications for risk governance. *PLoS ONE* 12(3), e0172867
- 567 Ager AA, Preisler H, Arca B, Spano D, Salis M (2014b) Wildfire risk estimation in the Mediterranean
 568 area. *Environmetrics* 25(6), 384–396
- 569 Alcasena F, Ager AA, Bailey JD, Pineda N, Vega-Garcia C (2018a) Towards a comprehensive wildfire
 570 management strategy for Mediterranean areas: Framework development and implementation in Catalonia,
 571 Spain. *Journal of Environmental Management* (doi: 10.1016/j.jenvman.2018.10.027)
- 572 Alcasena F, Salis M, Nauslar NJ, Aguinaga AE, Vega Garcia C (2016a) Quantifying economic losses from
 573 wildfires in black pine afforestations of northern Spain. *Forest Policy and Economics* 73, 153-167
- 574 Alcasena FJ, Ager AA, Salis M, Day MA, Vega-García C (2018b) Optimizing prescribed fire allocation
 575 for managing wildfire risk in central Catalonia. *Science of the Total Environment* 621, 872-885
- 576 Alcasena FJ, Salis M, Ager AA, Castell R, Vega-García C (2017) Assessing wildland fire risk transmission
 577 to communities in Northern Spain. *Forests* 8(2), 30 (doi:10.3390/f8020030)
- 578 Alcasena FJ, Salis M, Vega-Garcia C (2016b) A fire modeling approach to assess wildfire exposure in
 579 central Navarra, Spain. *European Journal of Forest Research* 135, 87-107
- 580 APCOR (2016) APCOR's Cork Yearbook 2016. [http://www.apcor.pt/wp-](http://www.apcor.pt/wp-content/uploads/2016/09/Boletim-estatistico-2016.pdf)
 581 [content/uploads/2016/09/Boletim-estatistico-2016.pdf](http://www.apcor.pt/wp-content/uploads/2016/09/Boletim-estatistico-2016.pdf)

- 582 Arianoutsou M (2001) Landscape changes in Mediterranean ecosystems of Greece: implications for fire
1 and biodiversity issues. *Journal of Mediterranean Ecology* 2, 165-178.
2 583
- 3
4 584 Aronson J, Pereira JS, Pausas JG (2009) Cork Oak Woodlands on the Edge: Ecology, Adaptive
5 Management, and Restoration. Washington, DC. Island Press
6 585
- 7
8 586 Baeza J, Santana VM, Pausas JG, Vallejo R (2011) Successional trends in standing dead biomass in
9 Mediterranean basin species. *Journal of Vegetation Science* 22, 467-474
10 587
- 11
12 588 Baeza MJ, Raventós J, Escarré A, Vallejo VR (2006) Fire risk and vegetation structural dynamics in
13 Mediterranean shrubland. *Plant Ecology* 187, 189-201
14 589
- 15
16 590 Bamber D (1975) The area above the ordinal dominance graph and the area below the receiver operating
17 characteristic graph. *Journal of Mathematical Psychology* 12, 387-415
18 591
- 19
20 592 Barberis A, Dettori S, Filigheddu MR (2003) Management problems in Mediterranean cork oak forests:
21 post-fire recovery. *Journal of Arid Environments* 54, 565-569
22 593
- 23
24 594 Bedia J, Herrera S, Camia A, Moreno JM, Gutierrez JM (2014) Forest fire danger projections in the
25 Mediterranean using ENSEMBLES regional climate change scenarios. *Climatic Change* 122(1-2), 185-199
26 595
- 27
28 596 Bergot M, Cloppet E, Pérarnaud V, Déqué M, Desprez-Loustau ML (2004) Simulation of potential range
29 expansion of oak disease caused by *Phytophthora cinnamomi* under climate change. *Global Change Biology*
30 597
31 10, 1539-1552
32 598
- 33
34 599 Bonet A, Pausas JG (2007) Old field dynamics on the dry side of the Mediterranean Basin: patterns and
35 processes in semiarid SE Spain. In: Cramer, V.A., Hobbs, R.J. (Eds.), *Old Fields Dynamics and Restoration of*
36 600
37 *Abandoned Farmland*. Island Press, Washington, DC, pp. 247-264
38 601
- 39
40 602 Bugalho MN, Caldeira MC, Pereira JS, Aronson J, Pausas JG (2011) Mediterranean cork oak savannas
41 require human use to sustain biodiversity and ecosystem services. *Frontiers in Ecology and the Environment* 9,
42 603
43 278-286
44 604
- 45
46 605 Calkin DE, Thompson MP, Finney MA (2015) Negative consequences of positive feedbacks in US
47 wildfire management. *Forest Ecosystems* 2: 9
48 606
- 49
50 607 Camia A, Liberta G, San-Miguel-Ayán J (2017) Modeling the impacts of climate change on forest fire
51 danger in Europe. JRC Technical Report doi:10.2760/768481
52 608
- 53
54 609 Catry FX, Moreira F, Cardillo E, Pausas JG (2012a) Post-fire management of cork oak forests. In: Moreira
55 F, Arianoutsou M, Corona P, De las Heras J (Eds.), *Post-fire Management and Restoration of European Forests*.
56 610
57 *Managing Forest Ecosystems*, vol. 24. Springer, pp. 195-222
58 611
59
60
61
62
63
64
65

- 612 Catry FX, Moreira F, Duarte I, Acacio V (2009) Factors affecting post-fire crown regeneration of cork
1 oak (*Quercus suber*) trees. *European Journal of Forest Research* 128, 231–240
2
- 613
3
- 614 Catry FX, Moreira F, Pausas JG, Fernandes PM, Rego F, Cardillo E, Curt T (2012b) Cork oak vulnerability
4
5
615 to fire: the role of bark harvesting, tree characteristics and abiotic factors. *PLoS ONE* 7(6), e39810
6
7
- 616 Chas-Amil ML, Touza J, García-Martínez E (2013) Forest fires in the wildland–urban interface: a spatial
8
9
617 analysis of forest fragmentation and human impacts. *Applied Geography* 43, 127–137
10
11
- 618 Corona P, Quatrini V, Schirru M, Dettori S, Puletti N (2018) Towards the economic valuation of ecosystem
12
13
619 production from cork oak forests in Sardinia (Italy). *iForest* 11(5), 660-667
14
15
- 620 Costa A, Madeira M (2011) Assessment of Mediterranean evergreen oak (*Quercus suber* L.) woodlands
16
17
621 loss. Consequence of climate changes effects? A case study at south-western Portugal. Proceedings of the
18
19
622 Global Conference on Global Warming 2011. Lisbon, Portugal, 11-14 July
20
21
- 623 Costa A, Madeira M, Plieninger T (2014) Cork oak woodlands patchiness: a signature of imminent
22
23
624 deforestation? *Applied Geography* 54, 18
24
25
- 625 Curt T, Borgniet L, Bouillon C (2013) Wildfire frequency varies with the size and shape of fuel types in
26
27
626 south-eastern France: implications for environmental management. *Journal of Environmental Management*
28
29
627 117, 150–161
30
31
- 628 Curt T, Frejaville T (2018) Wildfire Policy in Mediterranean France: How Far is it Efficient and
32
33
629 Sustainable? *Risk Analysis* 38(3), 472-488
34
35
- 630 Curt T, Pausas JG (2010). Are changes in fire regime threatening cork oak-shrubland mosaics? IUFRO-
36
37
631 IALE International Conference “Landscapes Forest and Global Change: New Frontiers in Management,
38
39
632 Conservation and Restoration”. p. 60-69. Bragança, Portugal. September 2010
40
41
- 633 Dettori S, Filigheddu MR (2003) La Sughericoltura. In: “L’Arboricoltura da Legno: un’attività produttiva
42
43
634 al servizio dell’ambiente” (editor G.F. Minotta). Bologna, 247 pp
44
45
- 635 Dettori S, Filigheddu MR (2016) Il sughero in enologia: analisi della filiera nazionale. *L’Italia Forestale*
46
47
636 *e Montana* 71(6), 331-343
48
49
- 637 Dettori S, Filigheddu MR, Deplano G, Escamilla Molgora J, Ruiu M, Sedda L (2018) Employing a spatio-
50
51
638 temporal contingency table for the analysis of cork oak cover change in the Sa Serra region of Sardinia.
52
53
639 *Scientific Reports* 8, 16946, doi:10.1038/s41598-018-35319-1
54
55
- 640 DG Environment (2007) Interpretation manual of European Union habitats – EUR 27. 142 p.
56
57
641 (http://ec.europa.eu/environment/nature/legislation/habitatsdirective/docs/2007_07_im.pdf)
58
59
60
61
62
63
64
65

- 642 Dimitrakopoulos AP, Vlahou M, Anagnostopoulou CG, Mitsopoulos ID (2011) Impact of drought on
1 wildland fires in Greece: implications of climatic change? *Climatic Change* 109, 331–347
2
3
4 644 EEA (2007) European forest types. Categories and types for sustainable forest management reporting and
5
6 645 policy. European Environment Agency (EEA) technical report no 9/2006, 2nd edn, Copenhagen
7
8 646 EEA (2017a) Copernicus Land Monitoring Service – Corine Land Cover ([https://www.eea.europa.eu/data-](https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-corine)
9
10 647 [and-maps/data/copernicus-land-monitoring-service-corine](https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-corine))
11
12 648 EEA (2017b) Copernicus Land Monitoring Service - EU-DEM ([https://www.eea.europa.eu/data-and-](https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-eu-dem)
13
14 649 [maps/data/copernicus-land-monitoring-service-eu-dem](https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-eu-dem))
15
16 650 FAO (2013) State of Mediterranean Forests. 174 p. ISBN 978-92-5-107538-8
17
18 651 Fernandes P, Davies GM, Ascoli D, Fernández C, Moreira F, Rigolot E, Stoof CR, Vega JA, Molina DM
19
20 652 (2013) Prescribed burning in southern Europe: developing fire management in a dynamic landscape. *Frontiers*
21
22 653 *in Ecology and the Environment* 11, e4–e14
23
24 654 Fernandes P, Rego FC (1996) Changes in fuel structure and fire behaviour with heathlands aging in
25
26 655 Northern Portugal. In: 13th Fire and Forest Meteorology Conference. Lorne, Australia, pp. 433–436
27
28 656 Ferreira-Leite F, Bento-Gonçalves A, Vieira A, Nunes A, Lourenço L (2016) Incidence and recurrence of
29
30 657 large forest fires in mainland Portugal. *Natural Hazards* 84(2), 1035–1053
31
32 658 Fick SE, Hijmans RJ (2017) Worldclim 2: New 1-km spatial resolution climate surfaces for global land
33
34 659 areas. *International Journal of Climatology* 37, 4302–4315
35
36 660 Figueiredo E, Raschi A (2011) Un’ immensa campagna avvolta dal verde. Reinventing rural areas in Italy
37
38 661 through tourism promotional images. *European Countryside* 3(1), 1–20
39
40 662 Flohn and Fantechi (1984) The Climate of Europe: Past, Present and Future. Springer Publisher, 357 pp.
41
42 663 ISBN: 978-90-277-1745-0
43
44 664 Forleo MB, Giaccio V, Giannelli A, Mastronardi L, Palmieri N (2017) Socio-Economic Drivers, Land
45
46 665 Cover Changes and the Dynamics of Rural Settlements: Mt. Matese Area (Italy). *European Countryside* 9(3),
47
48 666 435–457
49
50 667 Hosmer DW, Lemeshow S (2000) Applied Logistic Regression (2nd Edition). New York, NY: John Wiley
51
52 668 & Sons; DOI:10.1002/0471722146
53
54 669 Ibáñez B, Gómez-Aparicio L, Ávila JM, Pérez-Ramos IM, Marañón T (2017) Effects of *Quercus suber*
55
56 670 decline on woody plant regeneration: potential implications for successional dynamics in Mediterranean forests.
57
58 671 *Ecosystems* 20, 630
59
60
61
62
63
64
65


- 672 Jiménez-Ruano A, Rodrigues Mimbbrero M, de la Riva Fernández J (2017) Exploring spatial–temporal
1 dynamics of fire regime features in mainland Spain. *Natural Hazards and Earth System Sciences* 17, 1697–1711
2 673
- 3
4 674 Kerckhof A, Spalevic V, Van Eetvelde V, Nyssen J (2016) Factors of land abandonment in mountainous
5
6 675 Mediterranean areas: the case of Montenegrin settlements. *SpringerPlus* 5:485
7
- 8 676 Koutsias N, Xanthopoulos G, Founda D, Xystrakis F, Nioti F, Pleniou M, Mallinis G (2012) On the
9
10 677 relationships between forest fires and weather conditions in Greece from long-term 20 national observations
11
12 678 (1894–2010). *International Journal of Wildland Fire* 22(4), 493–507
13
- 14 679 Kovats RS, Valentini R, Bouwer LM, Georgopoulou E, Jacob D, Martin E, Rounsevell M, Soussana J-F
15
16 680 (2014) Europe. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects.*
17
18 681 *Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
19
20 682 *Change* [Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ, Bilir TE, Chatterjee M, Ebi KL, Estrada
21
22 683 YO, Genova RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR, White LL (eds.)]. Cambridge
23
24 684 University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1267–1326
25
- 26 685 Levin N, Tessler N, Smith A, McAlpine C (2016) The Human and Physical Determinants of Wildfires and
27
28 686 Burnt Areas in Israel. *Environmental Management* 58, 549–562
29
- 30 687 Lozano O, Salis M, Ager AA, Arca B, Alcasena FJ, Monteiro A, Finney MA, Del Giudice L, Scoccimarro
31
32 688 E, Spano D (2017) Assessing climate change impacts on wildfire exposure in Mediterranean areas. *Risk*
33
34 689 *Analysis* 37(10), 1898–1916
35
- 36 690 Luciano P, Lentini A, Cao OV (2005) La lutte aux lépidoptères défoliateurs des subéraies dans la Province
37
38 691 de Sassari. *Industria Grafica Poddighe s.r.l, Sassari*
39
- 40 692 Luciano P, Roversi PF (2001) Fillofagi delle querce in Italia. *Industria Grafica Poddighe s.r.l, Sassari, p.*
41
42 693 161
43
- 44 694 Lung T, Lavallo C, Hiederer R, Dosio A, Bouwer LM (2013) A multi-hazard regional level impact
45
46 695 assessment for Europe combining indicators of climatic and non-climatic change. *Global Environmental*
47
48 696 *Change* 23(2), 522–536
49
- 50 697 Madrigal J, Fernández-Migueláñez I, Hernando C, Guijarro M, Vega-Nieva DJ, Tolosana E (2016) Does
51
52 698 forest biomass harvesting for energy reduce fire hazard in the Mediterranean basin? A case study in the Caroig
53
54 699 massif (Eastern Spain). *European Journal of Forest Research* 136, 13–26
55
- 56 700 Molina DM, Castellnou M, García D, Salgueiro A (2010) Improving fire management success through
57
58 701 fire behavior specialists. In *Towards Integrated Fire Management – Outcomes of the European Project Fire*
59
60
61
62
63
64
65


- 702 Paradox - EFI Research Report nr. 23. "Fire Paradox", Project no. FP6-018505, European Commission, p. 105-
1 119. ISBN 970-952-5453-48-5. http://www.efi.int/files/attachments/publications/efi_rr23.pdf
2 703
- 3
4 704 Montiel Molina C, Galiana-Martín L (2016) Fire scenarios in Spain: A territorial approach to proactive
5
6 705 fire management in the context of global change. *Forests* 7, 273
- 7
8 706 Moreira F, Duarte I, Catry F, Acacio V (2007) Cork extraction as a key factor determining post-fire cork
9
10 707 oak survival in a mountain region of southern Portugal. *Forest Ecology and Management* 253, 30–37
- 11
12 708 Moreira F, Viedma O, Arianoutsou M, Curt T, Koutsias N, Rigolot E, Barbati A, Corona P, Vaz P,
13
14 709 Xanthopoulos G, Mouillot F, Bilgili E (2011) Landscape–wildfire interactions in southern Europe: implications
15
16 710 for landscape management. *Journal of Environmental Management* 92(10), 2389–2402
- 17
18 711 Moreno G, Franca A, Pinto Correia MT, Godinho S (2014) Multifunctionality and dynamics of
19
20 712 silvopastoral systems. In: "Forage resources and ecosystem services provided by Mountain and Mediterranean
21
22 713 grasslands and rangelands", Ed.: R Baumont, P Carrère, M Jouven, G Lombardi, A López-Francos, B Martin,
23
24 714 A Peeters, C Porqueddu, Options Méditerranéennes, Series A: Mediterranean Seminars 109, 421–436
- 25
26 715 Moreno JM, Vazquez A, Velez R (1998) Recent history of forest fires in Spain. In: Moreno JM (ed.) Large
27
28 716 forest fires. Backhuys Publishers, Leiden, pp 159–185
- 29
30 717 Moriondo M, Good P, Durao R, Bindi M, Giannakopoulos C, Corte-Real J (2006) Potential impact of
31
32 718 climate change on fire risk in the Mediterranean area. *Climate Research* 31, 85–95
- 33
34 719 Moritz MA, Batllori E, Bradstock RA, Gill AM, Handmer J, Hessburg PF, Leonard J, McCaffrey S, Odion
35
36 720 DC, Schoennagel T, Syphard AD (2014) Learning to coexist with wildfire. *Nature* 515, 58–66
- 37
38 721 Oliveira S, Pereira JMC, San-Miguel-Ayanz J, Lourenço L (2014) Exploring the spatial patterns of fire
39
40 722 density in Southern Europe using geographically weighted regression. *Applied Geography* 51, 143–157.
- 41
42 723 Oliveira V, Lauw A, Pereira H (2016) Sensitivity of cork growth to drought events: insights from a 24-
43
44 724 year chronology. *Climatic Change* 137, 261–274
- 45
46 725 Oubrahim H, Boulmane M, Bakker MR, Augusto L, Halim M (2015). Carbon storage in degraded cork
47
48 726 oak (*Quercus suber*) forests on flat lowlands in Morocco. *iForest* 9, 125–137
- 49
50 727 Pausas J, Vallejo VR (1999) The role of fire in European Mediterranean ecosystems. In: Chuvieco E (ed.)
51
52 728 Remote sensing of large wildfires in the European Mediterranean basin. Springer, Berlin, pp 3–16
- 53
54 729 Pausas JG (1997) Resprouting of *Quercus suber* in NE Spain after fire. *Journal of Vegetation Science* 8,
55
56 730 703–706
- 57
58 731 Pausas JG, Paula S (2012) Fuel shapes the fire-climate relationship: evidence from Mediterranean
59
60 732 ecosystems. *Global Ecology and Biogeography* 21, 1074–1082
- 61
62
63
64
65

- 733 Pausas JG, Pereira JS, Aronson J (2009) The tree. In: Aronson J, Pereira JS, Pausas JG, editors. Cork oak
1 woodlands on the edge: ecology, adaptive management, and restoration. Washington DC: Island Press. 11–23.
2 734
- 3
4 735 Plieninger TD, van der Horst D, Schleyer C, Bieling C (2014) Sustaining ecosystem services in cultural
5
6 736 landscapes. *Ecology and Society* 19(2), 59
7
- 8 737 Pereira H (2007) Cork: biology, production and uses. Amsterdam: Elsevier Publishing. 336 pp
9
- 10 738 Pereira MG, Calado TJ, DaCamara CC, Calheiros T (2013) Effects of regional climate change on rural
11
12 739 fires in Portugal. *Climate Research* 57, 187-200
13
- 14 740 Pereira MG, Trigo RM, da Camara CC, Pereira JMC, Leite SM (2005) Synoptic patterns associated with
15
16 741 large summer forest fires in Portugal. *Agriculture and Forest Meteorology* 129, 11–25
17
- 18 742 Piñol J, Beven K, Viegas D (2005) Modelling the effect of fire-exclusion and prescribed fire on wildfire
19
20 743 size in Mediterranean ecosystems. *Ecological Modelling* 183, 397-409
21
- 22 744 Piñol J, Castellnou M, Beven KJ. Conditioning uncertainty in ecological models: Assessing the impact of
23
24 745 fire management strategies. *Ecological Modelling* 207(1), 34–44
25
- 26 746 Rodrigues M, Jiménez-Ruano A, Peña-Angulo D, de la Riva J (2018) A comprehensive spatial-temporal
27
28 747 analysis of driving factors of human-caused wildfires in Spain using Geographically Weighted Logistic
29
30 748 Regression. *Journal of Environmental Management* 225, 177-192
31
- 32 749 Romero-Calcerrada R, Novillo J, Millington DA, Gomez-Jimenez I (2008) GIS analysis of spatial patterns
33
34 750 of human-caused wildfire ignition risk in the SW of Madrid (Central Spain). *Landscape Ecology* 23(3), 341–
35
36 751 354
37
- 38 752 Romero-Calcerrada R, Perry GLW (2004) The role of land abandonment in landscape dynamics in the
39
40 753 SPA ‘Encinares del río Alberche y Cofio, Central Spain, 1984–1999. *Landscape and Urban Planning* 66, 217-
41
42 754 232
43
- 44 755 Salis M, Ager AA, Finney MA, Arca B, Spano D (2014) Analyzing spatiotemporal changes in wildfire
45
46 756 regime and exposure across a Mediterranean fire-prone area. *Natural Hazards* 71(3), 1389–1418
47
- 48 757 Salis M, Arca B, Alcasena F, Arianoutsou M, Bacciu V, Duce P, Duguy B, Koutsias N, Mallinis G,
49
50 758 Mitsopoulos I, Moreno JM, Pérez JR, Rodriguez I, Xystrakis F, Zavala G, Spano D (2016a) Predicting wildfire
51
52 759 spread and behavior in Mediterranean landscapes. *International Journal of Wildland Fire* 25(10), 1015-1032
53
- 54 760 Salis M, Del Giudice L, Arca B, Ager AA, Alcasena-Urdirroz F, Lozano O, Bacciu V, Spano D, Duce P
55
56 761 (2018) Modeling the effects of different fuel treatment mosaics on wildfire spread and behavior in a
57
58 762 Mediterranean agro-pastoral area. *Journal of Environmental Management* 212, 490-505
59
60
61
62
63
64
65

- 763 Salis M, Laconi M, Ager AA, Alcasena FJ, Arca B, Lozano OM, Oliveira AS, Spano D (2016b) Evaluating
1 alternative fuel treatment strategies to reduce wildfire losses in a Mediterranean area. *Forest Ecology and*
2 764 *Management* 368, 207-221
3
4 765
5
6 766 San-Miguel-Ayanz J, Durrant T, Boca R, Libertà G, Boccacci F, Di Leo M, López Pérez J, Schulte E
7
8 767 (2016) Forest Fires in Europe, Middle East and North Africa 2015. EUR 28158 EN; doi:10.2788/914
9
10 768 (http://effis.jrc.ec.europa.eu/media/cms_page_media/40/Forest_fires_in_Europe_Middle_east_and_North_Afr
11
12 769 [ica_2015_final_pdf_JkX4Yl3.pdf](http://effis.jrc.ec.europa.eu/media/cms_page_media/40/Forest_fires_in_Europe_Middle_east_and_North_Africa_2015_final_pdf_JkX4Yl3.pdf))
13
14 770 San-Miguel-Ayanz J, Moreno JM, Camia A (2013) Analysis of large fires in European Mediterranean
15
16 771 landscapes: lessons learned and perspectives. *Forest Ecology and Management* 294, 11–22
17
18 772 Santana J, Porto M, Reino L, Beja P (2011) Long-term understory recovery after mechanical fuel reduction
19
20 773 in Mediterranean cork oak forests. *Forest Ecology and Management* 261, 447–459
21
22 774 Schaffhauser A, Curt T, Véla E, Tatoni T (2012) Fire recurrence effects on the abundance of plants
23
24 775 grouped by traits in *Quercus suber* L. woodlands and maquis. *Forest Ecology and Management* 282, 157–166
25
26 776 Scozzafava S, De Sanctis A (2006) Exploring the effects of land abandonment on habitat structures and
27
28 777 on habitat suitability for three passerine species in a highland area of central Italy. *Landscape and Urban*
29
30 778 *Planning* 75, 23-33
31
32 779 Silva JS, Catry F (2006) Forest fires in cork oak (*Quercus suber* L.) stands in Portugal. *International*
33
34 780 *Journal of Environmental Studies* 63, 235–257
35
36 781 Stephens SL, Collins BM, Biber E, Fulé PZ (2016) US federal fire and forest policy: emphasizing
37
38 782 resilience in dry forests. *Ecosphere* 7, e01584
39
40 783 Thompson MP, MacGregor DG, Dunn CJ, Calkin DE, Phipps J (2018) Rethinking the Wildland Fire
41
42 784 Management System. *Journal of Forestry* 116, 382-390
43
44 785 Thompson MP, Rodríguez y Silva F, Calkin DE, Hand MS (2017) A review of challenges to determining
45
46 786 and demonstrating efficiency of large fire management. *International Journal of Wildland Fire* 26, 562-573
47
48 787 Turco M, Bedia J, Di Liberto F, Fiorucci P, von Hardenberg J, Koutsias N, Llasat MC, Xystrakis F,
49
50 788 Provenzale A (2016) Decreasing Fires in Mediterranean Europe. PLoS ONE 11(3): e0150663.
51
52 789 <https://doi.org/10.1371/journal.pone.0150663>
53
54 790 Turco M, von Hardenberg J, Agha Kouchak A, Llasat MC, Provenzale A, Trigo RM (2017). On the key
55
56 791 role of droughts in the dynamics of summer fires in Mediterranean Europe. *Scientific Reports* 7, 81
57
58
59
60
61
62
63
64
65

- 792 Urbieta IR, Zavala MA, Marañón T (2008) Human and non-human determinants of forest composition in
 1 southern Spain: evidence of shifts towards cork oak dominance as a result of management over the past century.
 2 793
 3
 4 794 *Journal of Biogeography* 35, 1688–1700
- 5
 6 795 Varela MC (2017) From genetic improvement to global change: what advantages and threats for the
 7
 8 796 production, adaptation and evolution of cork oak? Proceedings of the International Congress on Cork Oak Trees
 9
 10 797 and Woodlands (ed. Dettori S., Fligheddu M.R., Cillara M.), ISBN 978-88-907678-0-7. Sassari (Italy), 25-26
 11
 12 798 May
- 13
 14 799 Vega-Garcia C, Woodard T, Adamowicz WL, Lee B (1995) A logit model for predicting the daily
 15
 16 800 occurrence of human caused forest fires. *International Journal of Wildland Fire* 5, 101–111.
- 17
 18 801 Vericat P, Piqué M, Koua O, Pla M (2010). Mapa de formacions forestals pures i mixtes de Catalunya a
 19
 20 802 partir del Mapa Forestal de España 1:50.000 digitalizado. Centre Tecnològic Forestal de Catalunya, Solsona.
- 21
 22 803 Viedma O, Moreno JM, Rieiro I (2006) Interactions between land use/land cover change, forest fires and
 23
 24 804 landscape structure in Sierra de Gredos (Central Spain). *Environmental Conservation* 33, 212–222
- 25
 26 805 Viegas DX, Figueiredo Almeida M, Ribeiro LM, Raposo J, Viegas MT, Oliveira R, Alves D, Pinto C,
 27
 28 806 Jorge H, Rodrigues A, Lucas D, Lopes S, Silva LF (2017) O complexo de incendios de Pedroogao Grande e
 29
 30 807 concelhos limítrofes, iniciado a 17 de junho de 2017.
 31
 32 808 <https://www.portugal.gov.pt/pt/gc21/comunicacao/documento?i¼4o-complexo-de-incendios-de-pedrogao->
 33
 34 809 [grande-econcelhos-limitrofes-iniciado-a-17-de-junho-de-2017](https://www.portugal.gov.pt/pt/gc21/comunicacao/documento?i¼4o-complexo-de-incendios-de-pedrogao-grande-econcelhos-limitrofes-iniciado-a-17-de-junho-de-2017).
- 35
 36 810 Viegas DX, Viegas MT (1994) A relationship between rainfall and burned area for Portugal. *International*
 37
 38 811 *Journal of Wildland Fire* 4, 11–16
- 39
 40 812 Vogiatzakis IN, Careddu MB (2003) Mapping the distribution and extent of *Quercus suber* habitats in
 41
 42 813 Sardinia: a literature review and a proposed methodology. *Geographical paper* 171.
- 43
 44 814 Xystrakis F, Kallimanis AS, Dimopoulos P, Halley JM, Koutsias N (2014) Precipitation dominates fire
 45
 46 815 occurrence in Greece (1900–2010): its dual role in fuel build-up and dryness. *Natural Hazards and Earth System*
 47
 48 816 *Sciences* 14, 21–32
- 49
 50 817 http://ec.europa.eu/environment/nature/natura2000/index_en.htm
- 51
 52 818 <http://www.sardegnaegeoportale.it/>
- 53
 54 819 <https://inventaire-forestier.ign.fr/>
- 55
 56 820 severicat.ctfc.cat
- 57
 58
 59
 60
 61
 62
 63
 64
 65








Figure 4










Figure 5












Supplementary Fig. 1 - Percentage of the annual provincial and regional areas affected by wildland fires in the period 2003-2015.




Supplementary Fig. 2 – Percent total area burned per month at the provincial level for the study areas. The wildland fire data refer to the period 2003-2015.




*Supplementary Fig. 3 - Map of the municipalities characterized by a presence of *Quercus suber* L. woodlands above 5% of the municipality area*



Supplementary Fig. 4 - Map of the average maximum temperatures of the months June-September for the study areas. Climate data from the 1-km resolution "WorldClim version 2" (Fick and Hijmans 2017)





Supplementary Fig. 5 - Map of the average cumulated precipitation of the months June-September for the study areas. Climate data from the 1-km resolution "WorldClim version 2" (Fick and Hijmans 2017)






Supplementary Fig. 6 – The role played by large wildland fire events on *Quercus suber* L. woodlands area burned. Red areas indicate the fire perimeters of La Jonquera (about 10,500 ha, on the left), which affected the province of Girona on July 2012, and Nuoro (about 9,000 ha, on the right), which affected the province of Nuoro on July 2007. As showed in the manuscript, large wildland fires substantially contributed to the total cork oak area burned in the study areas.




Table 1. Percentage of fire ignition locations as a function of the distance from cork oak woodlands and provinces for the study period 2003-2015 and for the previous 13-years in the study areas.


Province (Region) / Distance	2003-2015			1990-2002		
	0-1 km	1-10 km	>10 km	0-1 km	1-10 km	>10 km
Cagliari (Sardinia)	13.45	79.16	7.39	18.46	75.45	6.10
Nuoro (Sardinia)	42.68	56.64	0.68	43.99	52.73	3.29
Oristano (Sardinia)	23.68	75.77	0.54	35.33	64.05	0.62
Sassari (Sardinia)	50.51	46.17	3.32	56.10	41.74	2.16
Corse du Sud (Corsica)	31.60	52.53	15.87	31.66	52.45	15.89
Haute-Corse (Corsica)	33.71	53.05	13.23	35.42	52.87	11.72
Barcelona (Catalonia)	2.67	20.65	76.68	3.41	18.86	77.73
Girona (Catalonia)	43.51	37.08	19.42	39.87	34.73	25.39




Municipalities with relevant presence of *Quercus suber* L.


-  Municipality Boundaries
-  Provinces
-  *Quercus suber* L.






**Cumulated Rain (mm)
(June-September)**






 **Quercus suber**


 **La Jonquera wildfire (22 July 2012)**

3
 Km



 **Quercus suber**

 **Nuoro wildfire (23 July 2007)**

 **3 Km**