

Comparison of post-lethality thermal treatment conditions on the reduction of *Listeria monocytogenes* and sensory properties of vacuum packed ricotta salata cheese

Questa è la versione Post print del seguente articolo:

Original

Comparison of post-lethality thermal treatment conditions on the reduction of *Listeria monocytogenes* and sensory properties of vacuum packed ricotta salata cheese / Spanu, Carlo; Scarano, Christian; Spanu, V. .; Pala, C.; Di Salvo, R.; Piga, C.; Buschettu, L.; Casti, D.; Lamon, S.; Cossu, F.; Ibba, M.; DE SANTIS, Enrico Pietro Luigi. - In: FOOD CONTROL. - ISSN 0956-7135. - 50:(2015), pp. 740-747. [10.1016/j.foodcont.2014.10.022]

Availability:

This version is available at: 11388/45956 since: 2022-05-25T15:15:25Z

Publisher:

Published

DOI:10.1016/j.foodcont.2014.10.022

Terms of use:

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

Publisher copyright

note finali coverpage

(Article begins on next page)

1 **Comparison of post-lethality thermal treatment conditions on the reduction of *Listeria monocytogenes***
2 **and sensory properties of vacuum packed ricotta salata cheese**

3 C. Spanu*¹, C. Scarano¹, V. Spanu¹, C. Pala¹, R. Di Salvo², C. Piga², L. Buschetti³, D. Casti¹, S. Lamon¹, F.
4 Cossu¹, M. Ibba¹, E. P. L. De Santis¹

5

6 ¹*Department of Veterinary Medicine, University of Sassari, Via Vienna 2, 07100, Sassari, Italy*

7 ²*Agris, Dipartimento per la Ricerca nelle Produzioni Animali, Loc. Bonassai, S.S. 291 Sassari-Fertilia – Km.*
8 *18,600, Italy*

9 ³*Cooperativa Allevatori Ovini Formaggi Soc. Coop. Agricola, Loc. "Perda Lada" Fenosu, 09170, Oristano,*
10 *Italy*

11

12 *Corresponding author. Tel.: +39 079 229454; fax.: +39 079 229458. *E-Mail address:* cspanu@uniss.it (C.
13 Spanu); Via Vienna 2, 07100, Sassari, Italy.

14

15 **Abstract**

16 Ricotta salata is a whey protein cheese produced in Sardinia that in the last decades has been linked to
17 several recalls and in 2012 to a severe human listeriosis outbreak. Contamination of ricotta salata with *L.*
18 *monocytogenes* mainly occurs during post-process handling and generally origins from the processing
19 environment. The application of water bath heat treatment in vacuum packed ricotta salata is a possible
20 strategy to control *L. monocytogenes* superficial contamination. The objective of the present study was to
21 select a heat treatment able to inactivate *L. monocytogenes* count of at least 5 log. Nine temperature time
22 combinations, 75 °C, 85 °C and 90 °C applied for 15 min, 25 min and 40 min each were tested in ricotta
23 wheels artificially contaminated with a mixture of 5 *L. monocytogenes* strains. Inactivation was assessed
24 respectively one day and 30 days after heat treatment. The efficacy of treatments was evaluated based on the
25 reduction in *L. monocytogenes* counts, on the impact on sensory properties and on the cost of the treatment.
26 Two out of nine treatment combinations, i.e. 85 °C for 40 min and 90 °C for 40 min, were effective in
27 reducing *L. monocytogenes* contamination level of 5 log. No significant difference was observed in sensory

28 properties after the heat treatments. Therefore both combinations are eligible to conduct a successive study
29 aimed to extend the shelf-life of ricotta salata up to several months.

30 Keywords: *Listeria monocytogenes*, whey cheese, post-lethality treatment, sensory properties.

31

32 1. Introduction

33 Ricotta salata is a traditional whey protein cheese obtained in Sardinia (Italy) through the heat coagulation of
34 the whey remaining after the production of sheep's milk cheeses. The main phases of production technology
35 of ricotta salata are described as follows. Traditionally is manufactured using the whey remaining after the
36 production of hard sheep's milk cheese, usually Pecorino Romano PDO (protected denomination of origin),
37 which is stored in a silo at 45 °C until use. The whey is filtered and preheated at 60-70 °C using a plate heat
38 exchanger. The whey is then transferred in large open kettles with approximately 1,200-1,500 liters capacity,
39 added with 1% by weight of sodium chloride and heated to temperature above 80 °C for 30 minutes. As a
40 result of heating, curd start floating on the top of liquid, this is collected using perforated scoops and
41 transferred into plastic molds. The so called ricotta "*Toscanella*" is formed into cylindrical shapes and
42 pressed to enhance drainage. The curd is salted either by dry-salting (5% w/v) or by brine-salting and dried
43 for about 10 days in cold rooms at 10-12 °C. The manufacturing process result in cheese wheels weighing
44 approximately 3 kg with a pH of 6.1-6.9, a_w of 0.940-0.970, moisture of 50-60% (< 50% if intended for
45 grating), fat of 28-33% and protein of 14-23% (Spanu, Scarano, Spanu, Penna, Viridis, & De Santis, 2012;
46 Spanu, Spanu, Pala, Viridis, Scarano, & De Santis, 2013). The final product is individually packed in vacuum
47 bags and stored at refrigeration temperature with a set shelf-life which differs from three weeks up to several
48 months, depending on the food business operators. Packaging of ricotta salata depends on the final use of the
49 product, being ricotta salata wheels vacuum packed as a whole in shrinking bags if intended to be consumed
50 grated, for mixing with other cheeses or as an ingredient, or cut into wedges before packaging if consumed
51 plain. No preservatives are used for shelf life extension. In recent years contamination of ricotta salata with
52 *Listeria monocytogenes* leded voluntary recalls by international companies importing the product from
53 Sardinia. In 2008 the European Commission documented a case of *L. monocytogenes* infection associated
54 with the consumption of ricotta salata cheese (RASFF, 2008). The most recent and severe episode occurred

55 in the USA where a multistate outbreak of listeriosis linked to ricotta salata imported from Italy caused 20
56 hospitalization and 4 deaths (CDC, 2012). Heat treatments, such as thermization and pasteurization, applied
57 to milk during cheese making and to whey during ricotta production inactivate *Listeria* cells to levels of
58 approximately 3 to 6 log₁₀ cfu (Buazzi, Johnson, & Marth, 1992; Casadei, Esteves de Matos, Harrison, &
59 Gaze, 1998; ICMSF, 1996; Villani, Pepe, Mauriello, Moschetti, Sannino, & Coppola, 1996). Contamination
60 of whey cheeses with *L. monocytogenes* origins from the processing environment and is localized almost
61 exclusively on the rind, with a reported prevalence in ricotta salata of approximately 20% (Pintado &
62 Malcata, 2000; Lioliou, Litopoulou-Tzanetaki, Tzanetakis, & Robinson, 2001; Ibba, Cossu, Spanu, Viridis,
63 Spanu, Scarano, & De Santis, 2013; Spanu, Scarano, Ibba, Spanu & De Santis, 2015). The intrinsic
64 properties of ricotta salata support the growth of *L. monocytogenes*, once onto the product, to level as high as
65 7.0 log₁₀ cfu g⁻¹ of rind, potentially harmful to human health (Spanu, Scarano, Spanu, Penna, Viridis, & De
66 Santis, 2012). Ricotta salata produced in Sardinia is mainly exported in North America and in other
67 European countries. However, international health authorities accept different health risk for *L.*
68 *monocytogenes*, leading to an absence in 25 g recommended by FDA and 10² cfu g⁻¹ criteria at the time of
69 consumption set by European Commission (EC) Regulation No. 2073/2005. Even with a strict application of
70 good hygienic practices during production, superficial contamination of ricotta salata could not be totally
71 avoided, but only reduced (Tompkin, Scott, Bernard, Sveum, & Gombas, 1999). Therefore, the application
72 of alternative control strategies should be applied if the product is exposed to environmental contamination
73 after the lethality treatment (e.g., cooking) and before packaging (FSIS, 2014). In order to reduce *L.*
74 *monocytogenes* contamination in ready to eat food, a number of post-package decontamination methods have
75 been proposed, such as thermal pasteurization, irradiation and high-pressure. The efficacy of these
76 decontamination technologies in different ready to eat products have been reviewed (Zhu, Du, Cordray, &
77 Ahn, 2005). The final choice of the treatment to apply in ready to eat food stays on the food business
78 operator based on scientific evidences on the efficacy, but is certainly cost-oriented. Heat post-lethality
79 treatments (i.e. hot water bath and steam pasteurization) are widely used in the food industry due to their
80 effectiveness in reducing the load of pathogenic microorganisms (Arnoldi, 2002; Orta-Ramirez & Smith,
81 2002). The effectiveness of a thermal treatment is influenced by several factors such as temperature-time

82 ratio, food composition, size and weight of the product and microorganism characteristics (Doyle, Mazzotta,
83 Wang, Wiseman, & Scott, 2001; Ray, 2004; Sofos, 2002; Yen, Sofos, & Schmidt, 1991). Hence, the
84 validation of a post-lethality treatment should be designed around the product, taking into account the
85 formulation, packaging and the expected storage and use conditions. Therefore, results obtained on a specific
86 product cannot be extended on another product, even if similar. Previous research demonstrated the efficacy
87 of the immersion of vacuum packed ricotta salata wheels in water bath at 85 °C for 90 min in reducing *L.*
88 *monocytogenes* counts of 6 log₁₀ cfu g⁻¹ of rind (Spanu, Spanu, Pala, Viridis, Scarano, & De Santis, 2013).
89 However, the effect on sensory characteristics of such treatment was not investigated. Any technological
90 interventions that negatively affect the sensory quality of a product become useless for a commercial
91 purpose. For this reason sensory evaluation play a non negligible role in this type of investigation.
92 The objective of the present study was to compare 9 different temperature-time conditions for the superficial
93 treatment of whole ricotta salata wheels. The efficacy will be evaluated taking into account the extent of
94 reduction and survival of artificially inoculated *L. monocytogenes* and the impact on sensory properties. The
95 results will be used to select the temperature-time ratio to perform a further study aimed to extend the shelf-
96 life of ricotta salata up to 180 days.

97

98 **2. Materials and methods**

99

100 *2.1. Ricotta salata samples*

101 A total of 465 vacuum packed ricotta salata wheels were provided by a local cheese-making plant using
102 sheep milk. Samples were randomly selected from 3 different batches (155 ricotta wheels for each batch) and
103 stored in a cold room at 4±2 °C until the experiment was performed. Immediately after their arrival samples
104 were labeled according to their use for the experiment. Experimental Units (EUs) were defined ricotta salata
105 wheels artificially contaminated with *L. monocytogenes* and successively submitted to heat treatment.
106 Positive Controls (PCs) were defined ricotta salata wheels artificially contaminated with *L. monocytogenes*.
107 Blank Samples (BLs) were defined the units not inoculated and used to evaluate the level of natural
108 contamination of ricotta salata with *L. monocytogenes*. Sensory Units (SEs) were defined the not inoculated

109 samples used to evaluate sensory properties either after heat treatment (SEt) or as control with no treatment
110 (SEc). Composition Units (CUs) were defined not inoculated samples used for the determination of intrinsic
111 properties (pH and a_w) and composition (moisture, fat and proteins) after heat treatments.

112

113 2.2. Artificial inoculation

114 The Technical Guidance document prepared by the EU Community Reference Laboratory (CRL) for *L.*
115 *monocytogenes* (Beaufort, Cornu, Bergis, Lardeux, & Lombard, 2014) was used for the experiment
116 designing. A mixture of 5 *L. monocytogenes* strains was used to artificially contaminate EUs and PCs ricotta
117 salata wheels. Of the strains that composed the inoculum one was the reference strain ATTC 19111 (serovar
118 1/2a) obtained from American Type Culture Collection (Manassas, VA, USA), while the other four were
119 wild-type strains (respectively serotypes 1/2a, 1/2b, 1/2c and 4b), previously recovered from the cheese-
120 making plant environments or from ricotta salata. The wild-type strains were selected in order to be
121 representative of the main serotypes associated with foodborne listeriosis. All the strains were stored at -80
122 °C in Brain Heart Infusion (BHI) broth (Oxoid, Basingstoke, UK) with glycerol (15% v/v). The inoculum
123 level was aimed to demonstrate a reduction in *L. monocytogenes* level, or Performance Criterion (PC), of 5
124 \log_{10} cfu g^{-1} of rind, considered to suffice to attain a Food Safety Objective (FSO) of 10^2 cfu g^{-1} throughout
125 the entire storage period under refrigeration. Previous experiments were conducted to standardize the
126 preparation of inoculum according to the indications contained in the Guidelines for conducting *Listeria*
127 *monocytogenes* challenge testing of foods (Scott, Swanson, Frier, Pruett jr., Sveum, Hall, Smoot, & Brown,
128 2005). In order to prepare cells in the same physiological state (late exponential or early stationary phase)
129 each strain was separately inoculated into tubes containing BHI broth and cultured overnight at 30 °C in a
130 shaking water bath (100 rev min^{-1}). To adapt cultures at refrigeration temperatures, cells were then
131 subcultured into 10 mL of BHI and incubated at 4 ± 2 °C for approximately 15 days. A “mixed working
132 culture” was obtained by transferring equal volumes of each individual culture into a sterile flask. The
133 concentration was adjusted to ca. 10^7 cfu mL^{-1} using sterile saline solution (0.85% NaCl). Plate count on
134 Trypticase Soy Agar (TSA, Biolife, Milan, Italy) was used to confirm concentrations. The whole surface of
135 ricotta salata wheels was evenly sprayed with 2 mL of *L. monocytogenes* mixed culture using an atomizer. A

136 holding period of 15 min at room temperature was allowed to inoculated samples in order to let the
137 suspension attach, after which ricotta salata were individually vacuum packed in shrink bags (Criovac Cook-
138 In HT-3000, Sealedair Ltd., St Neots, UK) and stored at refrigeration temperature until further use.

139

140 2.3. Heat treatment and experimental design

141 The experiment was conducted in three independent trials, one for each batch, conducted one month apart.
142 Heat treatment was performed by immersion of vacuum packed ricotta salata wheels in hot water bath. Nine
143 different temperature- time conditions were tested: 75 °C, 85 °C and 90 °C applied for 15 min, 25 min and
144 40 min each. The number and the types of ricotta salata samples used for each treatment condition are
145 reported in table 1. Immediately after the heat treatment ricotta salata wheels were immersed in a tank
146 containing iced water for approximately 2 hours and then stored at 4 ± 2 °C until analysis. The analysis points
147 or testing times (T) were: the day of inoculum and heat treatment, defined as T_0 ; 24 hours after heat
148 treatment, defined as T_1 and 30 days after heat treatment defined as T_{30} . T_1 was performed the day
149 subsequent the heat treatment to avoid false negative caused by the presence of sub-lethally injured *L.*
150 *monocytogenes* cells that may survive the heat treatment but are not immediately culturable. The PCs were
151 analysed at T_0 , 6 hours after inoculation to assess if the level of contamination was effectively $10^5 \log_{10} \text{cfu g}^{-1}$.
152 The BLs were also examined at T_0 , to account for eventual natural contamination of ricotta salata with *L.*
153 *monocytogenes*. The EUs and CUs were submitted to heat treatment and analysed at T_1 and T_{30} . Part of SE
154 units were treated (SEt) and part, used as negative controls (SEc), were no treated (Table 1).

155 The sampling plan with sample units, testing times and related analysis is summarized in table 2. The
156 effective temperature obtained on ricotta salata surface during each heat treatment was monitored using an
157 additional ricotta salata wheel where a data logger (KT 20T, Kimo, Montpon Ménéstérol, France) was placed
158 1.5 cm below the surface and the temperature recorder during the treatment.

159

160 2.4. Microbiological analysis

161 Detection and enumeration of *L. monocytogenes* (ISO 11290-1:1996/Amd 1:2004; ISO 11290-2:1998/Amd
162 1:2004) and enumeration of aerobic mesophilic bacteria (ISO 4833:2003) were conducted on 25 g of ricotta

163 salata aseptically collected cutting the rind up to 2 cm. In order to detect the presence of sublethally injured
164 cells that may survive in the product but may not be cultured on selective media, on heat treated samples the
165 enumeration of *L. monocytogenes* was also conducted using the Thin Agar Layer (TAL) method. The TAL
166 method consists in the overlay of a nonselective agar medium onto agar plates containing a selective medium
167 that combines the ability to enumerate and to differentiate heat injured cells (Kang and Fung, 1999; Wu and
168 Fung, 2001). From each positive sample, 5 suspected colonies of *L. monocytogenes* were submitted to
169 phenotypic identification. Multiplex PCR was carried out to confirm identification and to separate the major
170 serovars (1/2a, 1/2b, 1/2c and 4b) into distinct serogroups (Doumith, Buchrieser, Glaser, Jacquet, & Martin,
171 2004). The *prs* gene, specific for *Listeria* spp. was used as internal amplification control. A selection of the
172 strains recovered from EUs at T₃₀ was submitted to pulsed-field gel electrophoresis (PFGE) to confirm that
173 the strains recovered were the same that were inoculated. From each of the 3 replicate were selected up to
174 five strains for each temperature-time combination. In order to capture as much variability as possible a
175 preliminary screening of isolates to submit to PFGE was conducted based on the serogroups. PFGE was
176 carried out using the protocol proposed by Graves & Swaminathan (2001). The obtained restriction profiles
177 were analysed by visual examination to distinguish inoculated strains among each other and from strains
178 originating by natural contamination.

179

180 2.5. Physico-chemical properties and composition

181 Intrinsic properties and chemical composition of ricotta salata were determined to account for possible
182 interaction with *L. monocytogenes* survival and growth. PH and a_w were measured using pH meter GLP22
183 (Crison Instruments SA, Barcelona, Spain) and water activity meter Aqualab 4TE (Decagon, Pullman, WA,
184 USA), respectively. Near infrared transmittance (NIT) compositional analyzer (FOSS, Eden Prairie, MN,
185 USA) was used for the analysis of fat, moisture, protein and total solids. Differences in intrinsic properties
186 and composition ($\bar{x} \pm SD$) of ricotta salata cheese between the different temperature-time combinations used
187 for the heat treatment and over time (T₁ and T₃₀) were compared using Fisher's least significant difference
188 (LSD) test. Statistical analysis was performed with Statgraphics Centurion XVI software (StatPoint
189 Technologies, Warrenton, VA, USA).

190

191 *2.6. Sensory analysis*

192 The "Difference from control test" was applied to highlight sensory differences between heat treated samples
193 (SEt) and the negative control (SEc). This test is very helpful to determine difference between one or more
194 samples against the control and, if the difference is significant, to measure its size (Meilgaard, Civille, &
195 Carr, 1999). On the other hand this test can cause a sensory fatigue when many samples have to be taken into
196 account because, during each session, the control sample as reference and as blind sample must be served. In
197 order to avoid the sensory fatigue only five out nine temperature-time combination were evaluated: 75 °C
198 and 90 °C treated for 15 min and 40 min, and 85 °C treated for 25 min. Thirty judges (14 females and 16
199 males, aged 25-50 years) specialized in dairy products, previously selected for their sensitivity and after
200 attending a course of 60 hours in sensory analysis (ISO 8586-1: 1993), evaluated the samples against an
201 untreated control on a numerical category scale (0 = no difference and 9 = very large difference). Ricotta
202 salata samples were kept at 4-6 °C until sensory assessments. Before analysis the ricotta samples were
203 portioned extracting two opposing slices. The slices were further portioned into parallelepiped pieces (5 x 1.5
204 x 1.5 cm) and served, at room temperature, in odorless plastic containers marked with a random three-digit
205 number (Meilgaard, Civille, & Carr, 1999). Judges were also provided by a tray containing an unsalted
206 cracker and a glass of water. The evaluation was carried out in a randomized and balanced order (Macfie,
207 Bratchell, Greenhoff, & Vallis, 1989). Statistical analysis was performed with Statgraphics Centurion XVI
208 software (StatPoint Technologies) by the one-way ANOVA (factor: samples) and the Fisher's LSD.

209

210 *2.7. Validation of heat treatment*

211 The experiment was conducted in three independent trials for each of the 9 temperature-time combinations.
212 Samples used in each trial belonged to three different production batches (batch A, B and C). Analyses were
213 conducted at two different sampling times (T_1 and T_{30}). For each heat treatment combination and sampling
214 time were analyzed three samples. To account for a margin of safety, the effectiveness of heat treatment, i.e.
215 performance standard (Δ) was considered in the worst conditions, i.e. the minimum level of $\log_{10} \text{cfu g}^{-1}$
216 reduction in *L. monocytogenes* counts. Reduction obtained as consequence of heat treatments was calculated

217 independently for each batch, by computing the minimum difference between the concentration (\log_{10} cfu g⁻¹) before the treatment (T₀) and after the treatment (T₁ and T₃₀) observed in the triplicate samples.

219

220 2.8. Statistical analysis

221 Mean mesophilic bacteria counts (\log_{10} cfu g⁻¹), intrinsic properties ($\bar{x} \pm \text{SD}$) and composition (% \pm SD) between the different temperature-time combinations at T₁ and T₃₀ were compared using Fisher's least significant difference (LSD) test. All statistical analyses were performed with Statgraphics Centurion XVI software (StatPoint Technologies, Warrenton, VA, USA).

225

226 3. Results

227 3.1. *L. monocytogenes* contamination and background microflora

228 Natural contamination of ricotta salata rind with *L. monocytogenes* occurred in 6 out of 18 BLs (30.0%) all originating from the first batch. Enumeration of *L. monocytogenes* was possible in five BLs, showing a level of contamination of $2.68 \pm 0.51 \log_{10}$ cfu g⁻¹ ($\bar{x} \pm \text{SD}$). The mean level of artificial contamination expressed as \log_{10} cfu g⁻¹ ($\bar{x} \pm \text{SD}$) obtained on the rind of PCs units at T₀ was 4.82 ± 0.43 , 5.5 ± 0.04 and 5.36 ± 0.09 in the first, second and third replicate, respectively. Enumeration of aerobic mesophilic bacteria was conducted on 18 BLs, 9 PCs and 162 EUs. In BLs aerobic mesophilic population ($\bar{x} \pm \text{SD} \log_{10}$ cfu g⁻¹) was 7.41 ± 0.47 , 7.45 ± 1.04 , 7.83 ± 1.00 while in PCs was 7.21 ± 0.31 , 7.78 ± 0.68 and 8.83 ± 0.11 in the first, second and third batch respectively. The mean \log_{10} cfu g⁻¹ reduction in total bacterial counts observed at T₁ ranged between ca. 1.0 and 3.0 for ricotta treated at 75 °C, 4.0 and 5.5 for treatment at 85 °C and from 5 to 6 for 90 °C. After 30 days of refrigerated storage the microbial population increased of less than 1.0 \log_{10} cfu g⁻¹ in ricotta salata treated at 75 °C and of ca. 1.0 and 2.0 \log_{10} cfu g⁻¹ in samples treated respectively at 85 °C and 90 °C. Pair-wise comparison of aerobic mesophilic bacteria counts between ricotta salata samples submitted to the 9 treatment combinations and between samples analysed at T₁ and T₃₀ are reported in table 3.

241

242 3.2. Inactivation and survival of *L. monocytogenes*

243 The lethal effect was evaluated on 9 EUs (3 for each replicate) for each temperature-time combination at T_1
244 and T_{30} . The minimum differences in *L. monocytogenes* counts between T_1 and T_0 (Δ_1) and T_{30} and T_0 (Δ_{30})
245 for each of the 9 treatments are reported in table 4. At T_1 three out of nine combinations, i.e. 85 °C for 40
246 min, 90 °C for 25 min and 90 °C for 40 min, were effective either with the enumeration and detection
247 methods. However, the 90 °C for 25 min combination showed the survival of *L. monocytogenes* after storage
248 at 4 °C for 30 days. The complete description of the effect of each treatment at T_1 and T_{30} is reported in table
249 5 and figure 1.

250 Overall, 334 strains were confirmed as *L. monocytogenes* by molecular identification. Twenty-seven strains
251 (90.0%) isolated from BLs were serogroup 1/2a, while 3 (10.0%) were serogroup 1/2c. Of the 49 strains
252 isolated from PCs, 17 (34.7%) were serogroup 1/2a, 15 (30.6%) serogroup 1/2b, 11 (22.4%) serogroup 1/2c
253 and 6 (12.2%) serogroup 4b. From EUs at T_1 were isolated 95 strains which were grouped as follows: 32
254 (33.7%) serogroup 1/2a, 12 (12.6%) serogroup 1/2b, 16 (16.8%) serogroup 1/2c and 35 (36.8%) serogroup
255 4b. From EUs at T_{30} were isolated 160 strains which were grouped as follows: 52 (32.5%) serogroup 1/2a, 23
256 (14.4%) serogroup 1/2b, 41 (25.6%) serogroup 1/2c and 44 (27.5%) serogroup 4b. Of *L. monocytogenes*
257 recovered from EUs at T_{30} were submitted to PFGE 19, 13 and 30 strains from batch A, B and C
258 respectively. Strains recovered showed the same PFGE profile of the inoculated strains and belonged
259 exclusively to the wild type: 18 (29.0%) were 1/2a, 12 (19.4%) were 1/2b, 13 (21.0%) were 1/2c and 19
260 (30.6%) were 4b.

261

262 3.3. Temperature monitoring

263 The initial temperature of ricotta salata rind before the immersion in hot water was 6.0 ± 0.8 °C. Figure 2
264 shows the temperature profile recorded on the rind of ricotta salata during water bath heat treatments for the
265 three temperatures.

266

267 3.4. Sensory features

268 The results of sensory analysis (average values \pm SD) are shown in table 7. The blind control allowed
269 estimating the *placebo* effect, produced by asking to find a difference when in fact no differences exist.

270 Statistic inferences are estimates by comparing the samples and the blind control. All the heat treated
271 samples are significantly different from the untreated one (blind control) with the exception of that treated at
272 lowest temperature-time combination (75 °C for 15 min). However the difference size goes from 1.3 to 1.8
273 points that converted in the verbal scale correspond to “slight/moderate difference”. The heat treated samples
274 were not different between each other, meaning that heating from 75 to 90 °C for a time ranging between 15
275 and 40 min do not significantly change the sensory characteristics of ricotta salata.

276 3.5. Ricotta salata composition

277 Physico-chemical characteristics were determined on untreated samples (18 BLs) and heat treated samples
278 (162 CUs). Intrinsic properties values ($\bar{x} \pm SD$) for untreated samples were 6.42 ± 0.09 for pH and 0.963 ± 0.01
279 for a_w . Composition values ($\% \pm SD$) were respectively of 55.35 ± 2.09 for moisture, 21.75 ± 2.42 for fat,
280 14.55 ± 1.37 for proteins and 4.56 ± 1.38 for salt. Differences in intrinsic properties and composition between
281 the 9 treatment combinations at T_1 and T_{30} are reported in table 6. PH and a_w were always within limits for *L.*
282 *monocytogenes* growth and no significant difference was observed between values of ricotta salata submitted
283 to heat treatment with different temperature-time conditions ($P > 0.05$).

284

285 4. Discussion

286 In the last decades ricotta salata has been associated with several recalls due to *L. monocytogenes*
287 contamination and more recently even with foodborne listeriosis outbreaks. Contamination of ricotta salata
288 with *L. monocytogenes* mainly originates from food processing environment and is localized almost exclusively
289 on the rind (Pilo, Marongiu, Corgiolu, Viridis, Scarano, & De Santis, 2007). Whole ricotta salata wheels are
290 generally intended to be consumed grated including the rind. Post-process control strategies are needed in
291 order to comply with international health authorities limits. Hot water bath treatment in vacuum packed
292 ricotta salata has been previously evaluated, demonstrating to be an effective and economic method to
293 inactivate surface contamination of ricotta salata cheese (Spanu, Spanu, Pala, Viridis, Scarano, & De Santis,
294 2013). However, optimization of the process was needed in order to account for the level of reduction in *L.*
295 *monocytogenes* counts (\log_{10} cfu g^{-1}), changes in sensory properties and cost of the treatment. In the present
296 study was compared the listericidal effect of nine temperature-time combinations for the treatment of

297 artificially contaminated whole ricotta salata wheels. The recovery of *L. monocytogenes* from not inoculated
298 blank samples, confirms that natural contamination of ricotta salata is not a rare finding. The prevalence of
299 contamination of ricotta salata produced in Sardinia is estimated around 20% (Ibba, Cossu, Spanu, Viridis,
300 Spanu, Scarano, & De Santis, 2013; Spanu, Scarano, Ibba, Spanu & De Santis, 2015). This level can result,
301 when ricotta is stored at refrigeration temperatures for up to two months, in concentration of the pathogens of
302 approximately $10^7 \log_{10} \text{ cfu g}^{-1}$, potentially harmful to human health (Spanu, Scarano, Spanu, Penna, Viridis,
303 & De Santis, 2012). The present study was aimed to validate the temperature-time combinations able to
304 reduce *L. monocytogenes* concentration of $5 \log \text{ cfu g}^{-1}$, considered sufficient to comply with the food safety
305 objective of $<100 \text{ cfu g}^{-1}$ for the products placed on the market during their shelf-life (EC, 2005).
306 Enumeration of *L. monocytogenes* in positive control units confirmed that the desired level of contamination
307 of 10^5 cfu g^{-1} was obtained. Out of nine temperature-time combinations only two, $85 \text{ }^\circ\text{C}$ for 40 min and 90
308 $^\circ\text{C}$ for 40 min, showed to be effective in reducing *L. monocytogenes* to undetectable levels either at T_0 and
309 T_{30} . These combinations allowed to reach respectively $56.3 \pm 1.5 \text{ }^\circ\text{C}$ and $57.7 \pm 1.4 \text{ }^\circ\text{C}$ on ricotta surface,
310 effective in killing *L. monocytogenes*. On one hand, little or no efficacy was observed for treatment
311 conducted at $75 \text{ }^\circ\text{C}$, regardless of the time of application, while on the other hand no efficacy was
312 demonstrated for treatment conducted for 15 min, regardless of the temperature used. Despite an initial
313 inactivation of up to 5.0 log at T_0 as consequence of treatments performed at $75 \text{ }^\circ\text{C}$ (i.e. 25 and 40 min), *L.*
314 *monocytogenes* was still culturable with the detection method. The survival and successive growth during
315 storage at refrigeration temperature for 30 days resulted in counts as high as ca. 7 log. Failure of heat
316 treatment at 75°C to inactivate *L. monocytogenes* could be explained with the difference between water
317 temperature and the maximum temperature obtained on ricotta rind ($47.5 \text{ }^\circ\text{C}$). Treatment conducted at $85 \text{ }^\circ\text{C}$
318 for 25 min resulted in a temperature on ricotta rind of $49.8 \pm 1.5 \text{ }^\circ\text{C}$ which allowed *L. monocytogenes* survival
319 to concentrations of up to 1 log at day zero. As consequence the microorganism grew during the successive
320 storage to level as high as 5 log. Although *L. monocytogenes* was not countable with the enumeration method
321 after 30 days in samples treated at $90 \text{ }^\circ\text{C}$ for 25 minutes, it was still detectable with the qualitative method.
322 No guarantee can be provided that the pathogen will not growth to levels potentially dangerous to human
323 during ricotta salata shelf-life. The highest temperature detected on ricotta salata rind after 25 min was 52.5

324 °C for the treatment at 90 °C. This could explain the presence of heat injured cells, which recovered after the
325 subsequent storage at refrigeration temperature for 30 days. Strains capable to survive and growth after heat
326 treatment belonged to the wild type, suggesting that they are characterized by a greater resistance as
327 compared with reference strains. As far as the gap between the temperature of water during treatments and
328 the temperature recorded on the ricotta salata rind it should be noted that temperatures were detected 1.5 cm
329 below the ricotta surface, which may underestimate the effective temperature reached on the interface
330 between packaged ricotta and water. The heat transfer is a function of the thermal properties of foods, which
331 depend, among other factors, by chemical composition and temperature. However, due to the complexity of
332 heat transfer calculations, specific experiments should be conducted in order to define the specific thermal
333 properties of ricotta salata. Changes in sensory properties of heat treated ricotta salata were observed with
334 respect of untreated samples, but no differences were among treatments. This indicates the feasibility of
335 using more protective treatments with no negative implication for ricotta salata sensory profile.

336

337 **5. Conclusion**

338 Contamination of Ricotta salata with *L. monocytogenes* can effectively be controlled by the application of
339 water bath heat treatment applied after packaging of the product. Treatments performed at 85 °C for 40 min
340 or 90 °C for 40 min can be effectively used to obtain a reduction of 5 log of the pathogen. No significant
341 difference was observed in the sensory properties between the treatments. Although treatments at 85 °C for
342 40 min might gather food processors favors as compared to 90 °C for 40 min, the latter may provide a greater
343 safety of the product when is stored for periods of time longer than 30 days. Treatments applied for 40 min,
344 either at 85 °C and 90 °C, are eligible as combination of choice to be used in a further study to assess the
345 efficacy as post-lethality treatment aimed to extend ricotta salata shelf-life.

346

347 **Acknowledgements**

348 This work was funded by “ Programma di Sviluppo Rurale Sardegna 2007-2013 Misura 124 Cooperazione
349 per lo Sviluppo di Nuovi Prodotti, Processi e Tecnologie nei Settori Agricolo Alimentare e in quello
350 Forestale –project ID: H78F13000050007. The authors are grateful to all the members of the joint dairy

351 industry consortium “Associazione Temporanea di Scopo - Aziende casearie Riunite” for their cooperation
352 in the research.

353

354 **References**

- 355 Arnoldi, A. (2002). Thermal processing and nutritional quality. In: Henry, C.J.K., Chapman, C. (Eds.),
356 The Nutrition Handbook for Food Processing. Woodhead, Cambridge, England, pp. 265-286.
- 357 Beaufort, A., Cornu, M., Bergis, H., Lardeux, A.L., Lombard, B. (2014). EURL Lm Technical guidance
358 document for conducting shelf-life studies on *Listeria monocytogenes* in ready-to eat foods. Version 3.
359 Community Reference Laboratory for *Listeria monocytogenes*. Available at:
360 [http://www.fsai.ie/uploadedFiles/EURL%20Lm_Technical%20Guidance%20Document%20Lm%20shelf-life%20studies_V3_2014-06-06%20\(2\).pdf](http://www.fsai.ie/uploadedFiles/EURL%20Lm_Technical%20Guidance%20Document%20Lm%20shelf-life%20studies_V3_2014-06-06%20(2).pdf) Accessed October 2 2014.
- 361
- 362 Buazzi, M. M., Johnson, M. E., & Marth, E. H. (1992). Fate of *Listeria monocytogenes* during the
363 manufacture of mozzarella cheese. *Journal of Food Protection*, 55, 80-83.
- 364 Casadei, M. A., Esteves de Matos, R., Harrison, S. T., & Gaze, J. E. (1998). Heat resistance of *Listeria*
365 *monocytogenes* in dairy products as affected by the growth medium. *Journal of Applied Microbiology*,
366 84(2), 234-239.
- 367 CDC. (2012). Multistate Outbreak of Listeriosis Linked to Imported Frescolina Marte Brand Ricotta
368 Salata Cheese (Final Update). Available at: <http://www.cdc.gov/listeria/outbreaks/cheese-09-12/>. Accessed 10 May 2014
- 369
- 370 Doumith, M., Buchrieser, C., Glaser, P., Jacquet, C., & Martin, P. (2004). Differentiation of the major
371 *Listeria monocytogenes* serovars by multiplex PCR. *Journal of Clinical Microbiology*, 42, 3819-3822.
- 372 Doyle, M. E., Mazzotta, A. S., Wang, T., Wiseman, D. W., & Scott, V. N. (2001). Heat Resistance of
373 *Listeria monocytogenes*. *Journal of Food Protection*, 64(3), 410-429.
- 374 EC. (2005). Commission Regulation (EC) No. 2073/2005 of 15 November 2005 on microbiological
375 criteria for foodstuffs. [http://eur-](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2005:338:0001:0026:EN:PDF)
376 [lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2005:338:0001:0026:EN:PDF](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2005:338:0001:0026:EN:PDF) Accessed 28.06.14.

377 Food Safety and Inspection Service. (2014). Compliance Guideline: Controlling *Listeria monocytogenes*
378 in post-lethality exposed ready-to-eat meat and poultry products. United States Department of
379 Agriculture. Available at: [http://www.fsis.usda.gov/wps/wcm/connect/d3373299-50e6-47d6-a577-](http://www.fsis.usda.gov/wps/wcm/connect/d3373299-50e6-47d6-a577-e74a1e549fde/Controlling_LM_RTE_Guideline_0912?MOD=AJPERES)
380 [e74a1e549fde/Controlling_LM_RTE_Guideline_0912?MOD=AJPERES](http://www.fsis.usda.gov/wps/wcm/connect/d3373299-50e6-47d6-a577-e74a1e549fde/Controlling_LM_RTE_Guideline_0912?MOD=AJPERES). Accessed 20 March 2014.

381 Graves, L. M., & Swaminathan, B. (2001). PulseNet standardized protocol for subtyping *Listeria*
382 *monocytogenes* by macrorestriction and pulsed-field gel electrophoresis. *International Journal of Food*
383 *Microbiology*, 65, 55-62.

384 Ibba, M., Cossu, F., Spanu, V., Viridis, S., Spanu, C., Scarano, C., & De Santis, E. P. L. (2013). *Listeria*
385 *monocytogenes* contamination in dairy plants: evaluation of *Listeria monocytogenes* environmental
386 contamination in two cheese-making plants using sheeps milk. *Italian Journal of Food Safety*, 2, 109-
387 112.

388 ICMSF. (1996). Microorganisms in Foods 5: Microbiological Specification of Food Pathogens.
389 International Commission on Microbiological Specifications for Foods (ICMSF). London: Blackie
390 Academic & Professional, (Chapter 8).

391 ISO 8586-1:1993. Sensory analysis-General Guidance for The Selection, Training and Monitoring of
392 Assessors. Part 1: Selected assessors. International Organization for Standardization, Geneva,
393 Switzerland.

394 ISO 11290-1:1996/Amd 1:2004. Microbiology of food and animal feeding stuffs-Horizontal method for
395 the detection and enumeration of *Listeria monocytogenes*. Part 1: Detection method (1st ed.),
396 International Organization for Standardization, Geneva, Switzerland.

397 ISO 11290-2:1998/Amd 1:2004. Microbiology of food and animal feeding stuffs-Horizontal method for
398 the detection and enumeration of *Listeria monocytogenes*. Part 2: Enumeration method (1st ed.),
399 International Organization for Standardization, Geneva, Switzerland.

400 ISO 4833:2003. Microbiology of food and animal feeding stuffs-Horizontal method for the enumeration
401 of microorganisms: Colony-count technique at 30 °C. International Organization for Standardization,
402 Geneva, Switzerland.

- 403 Kang, D. H., Fung, D. Y. (1999). Thin agar layer method for recovery of heat-injured *Listeria*
404 *monocytogenes*. *Journal of Food Protection*, 62(11), 1346-1349.
- 405 Lioliou, K., Litopoulou-Tzanetaki, E., Tzanetakis, N., & Robinson, R. K. (2001). Changes in the
406 microflora of manouri, a Greek whey cheese, during storage. *International Journal of Dairy Technology*,
407 54(3), 100-106.
- 408 Macfie, H. J. H., Bratchell, N., Greenhoff, K., & Vallis, L. V. (1989). Designs to balance the effect of
409 order of presentation and first– order carry–over effects in hall tests. *Journal of Sensory Studies*, 4, 2,
410 129-148.
- 411 Meilgaard, M., Civille, G. V., & Carr, B. T. (1999). *Sensory Evaluation Techniques*, pp. 86-91 CRC
412 Press, Boca Raton, 1991.
- 413 Orta-Ramirez, A., & Smith, D. M. (2002). Thermal inactivation of pathogens and verification of
414 adequate cooking in meat and poultry products. *Advances in Food and Nutrition Research*, 44, 147-194.
- 415 Pilo, A. L., Marongiu, P., Corgiolu, G., Viridis, S., Scarano, C., & De Santis, E. P. L. (2007). *Listeria*
416 *monocytogenes* contamination sources in sheep cheese processing plants and strains virulence genes
417 typing. *In proceeding of: 5th IDF International Symposium On The Challenge to Sheep and Goats Milk*
418 *Sectors*, Alghero, Italy.
- 419 Pintado, M. E., & Malcata, F. X. (2000). Optimization of modified atmosphere packaging with respect to
420 physicochemical characteristics of Requeijão. *Food Research International*, 33, 821-832.
- 421 RASFF. (2008). The Rapid Alert System for Food and Feed Annual Report.
422 http://ec.europa.eu/food/food/rapidalert/report2008_en.pdf (accessed 10.03.2014).
- 423 Ray, B. (2004). *Fundamental food microbiology*. Boca Raton: CRC Press LLC.
- 424 Scott, V. N., Swanson, K., Frier, T. A., Pruett jr., W. P., Sveum, W. H., Hall, P.A., Smoot, L. A., Brown,
425 D. G. (2005). Guidelines for conducting *Listeria monocytogenes* challenge testing of foods. *Food*
426 *Protection Trends*, 25(11), 818-825.
- 427 Sofos, J. N. (2002). Approaches to pre-harvest food safety assurance. In F. J. M. Smulders & J. D.
428 Collins (Eds.), *Food safety assurance and veterinary public health: Vol. 1* (pp. 23-48). Wageningen,
429 Netherlands: Wageningen Academic Publishers.

- 430 Spanu, C., Scarano, C., Spanu, V., Penna, C., Viridis, S., & De Santis, E. P. L. (2012). *Listeria*
431 *monocytogenes* growth potential in Ricotta salata cheese. *International Dairy Journal*, 24, 120-122.
- 432 Spanu, C., Spanu, V., Pala, C., Viridis, S., Scarano, C., & De Santis, E. P. L. (2013). Evaluation of a post-
433 lethality treatment against *Listeria monocytogenes* on Ricotta salata cheese. *Food Control*, 30(1), 200-
434 205.
- 435 Spanu, C., Scarano, C., Ibba, M., Spanu, V., & De Santis, E. P. L. (2015). Occurrence and traceability of
436 *Listeria monocytogenes* strains isolated from sheep's milk cheese-making plants environment. *Food*
437 *Control*, 47, 318-325.
- 438 Tompkin, R. B., Scott, V. N., Bernard, D. T., Sveum, W. H. & Gombas, K. S. (1999). Guidelines to
439 prevent post-processing contamination from *Listeria monocytogenes*. *Dairy, Food and Environmental*
440 *Sanitation*, 19, 551-562.
- 441 Villani, F., Pepe, O., Mauriello, G., Moschetti, G., Sannino, L., & Coppola, S. (1996). Behavior of
442 *Listeria monocytogenes* during the traditional manufacture of water-buffalo Mozzarella cheese. *Letters*
443 *in Applied Microbiology*, 22, 357-360.
- 444 Wu, V. C. H., Fung, D. Y. C. (2001). Evaluation of thin agar layer method for recovery of heat-injured
445 foodborne pathogens. *Journal of Food Science*, 66(4), 580-583.
- 446 Yen, L. C., Sofos, J. N., & Schmidt, G. R. (1991). Effect of meat curing ingredients on thermal
447 destruction of *Listeria monocytogenes* in ground pork. *Journal of Food Protection*, 54, 408-412.
- 448 Zhu, M., Du, M., Cordray, J., & Ahn, D.U. (2005). Control of *Listeria monocytogenes* contamination in
449 ready-to-eat meat products. *Comprehensive Reviews in Food Science and Food Safety*, 4(2), 34-42.

1 Table 1. Temperature-time combinations used for water bath heat treatment of ricotta salata

Temperature-time condition		Ricotta salata samples					Total
Temperature	Minutes	BL ¹	PC ²	EUs ³	CUs ⁴	SEs ⁵	
No treatment	-	18	9	-	-	42	69
75 °C	15	-	-	18	18	12	48
	25	-	-	18	18	3	39
	40	-	-	18	18	12	48
85 °C	15	-	-	18	18	3	39
	25	-	-	18	18	12	48
	40	-	-	18	18	3	39
90 °C	15	-	-	18	18	12	48
	25	-	-	18	18	3	39
	40	-	-	18	18	12	48
Total		18	9	162	162	114	465

2 ¹BLs (Blank Samples): not inoculated units; ²PCs (Positive Controls): samples inoculated with *L.*

3 *monocytogenes*; ³EUs (Experimental Units): samples inoculated with *L. monocytogenes* and successively

4 heat treated; ⁴CUs (Composition Units): heat treated units for physico-chemical analysis; ⁵SEs (Sensory

5 Units): samples used to assess the effect of heat treatment on sensory properties.

6

7

8

9

10

11

12

13

14

15 Table 2. Number of ricotta salata wheels and analysis performed at each sampling time.

Analysis	Test units	Sampling time			Total
		T ₀ ^a	T ₁ ^b	T ₃₀ ^c	
Detection and enumeration of <i>L. monocytogenes</i> and aerobic mesophilic bacteria	BLs ¹	18	-	-	-
	PCs ²	9	-	-	9
	EUs ³	-	81	81	162
	CUs ⁴	-	81	81	162
Intrinsic properties and composition	BLs ¹	18	-	-	-
	CUs ⁴	-	81	81	162
Sensory analysis	SEs ⁵				38

16 Superscript letters are referred to the time between inoculation and analysis: ^a = day of inoculum and heat
 17 treatment; ^b = 24 hours after heat treatment; ^c = 30 days after heat treatment. Superscript numbers are referred
 18 to test units: ¹BLs (Blank Samples): not inoculated units; ²PCs (Positive Controls): samples inoculated with
 19 *L. monocytogenes*; ³EUs (Experimental Units): samples inoculated with *L. monocytogenes* and successively
 20 heat treated; ⁴CUs (Composition Units): heat treated units for physico-chemical analysis; ⁵SEs (Sensory
 21 Units): samples used to assess the effect of heat treatment on sensory properties.

22

23

24

25

26

27

28

29

30

31

32 Table 3. Comparison of aerobic mesophilic bacteria counts (\log_{10} cfu g^{-1} ; $\bar{x} \pm SD$) of heat treated ricotta
 33 salata with 9 different temperature-time combinations analyzed 24 h after the treatment (T_1) and after storage
 34 at refrigeration temperature for 30 days (T_{30}).

Treatment		Aerobic mesophilic bacteria			
Temperature	Minutes	+ve/n	T_1	+ve/n	T_{30}
75 °C	15	9/9	6.94± 0.62 ^A	9/9	7.06±0.80 ^A
	25	9/9	5.69± 0.55 ^B	9/9	6.20±1.36 ^A
	40	9/9	5.12± 0.42 ^B	7/9	6.50±1.28 ^A
85 °C	15	9/9	3.90± 0.85 ^C	9/9	4.88±0.88 ^B
	25	9/9	3.71± 0.94 ^C	5/9	4.58±1.14 ^B
	40	9/9	2.45± 1.97 ^D	4/9	4.14±1.78 ^{BC}
90 °C	15	9/9	2.02± 1.22 ^D	6/9	4.15±0.82 ^{BC}
	25	9/9	1.68± 1.29 ^D	3/9	2.90±0.78 ^C
	40	9/9	2.01± 1.12 ^D	3/9	3.67±1.62 ^{BC}

35 Means in the same column on the same testing time (T_1 or T_{30}) with different capital letter are significantly
 36 different ($P<0.05$).

37

38

39

40

41

42

43

44

45

46

47

48 Table 4. *Listeria monocytogenes* reduction (Δ)^a on ricotta salata rind after water bath heat treatment

Temperature	Batch	15 min		25 min		40 min	
		ΔT_1	ΔT_{30}	ΔT_1	ΔT_{30}	ΔT_1	ΔT_{30}
75 °C	A	-0.23	4.18	-4.54	2.26	-3.24	2.57
	B	-2.59	0.48	-5.45	-5.45	-5.45	-5.45
	C	-1.78	-1.36	-5.27	-1.73	-5.27	-5.27
85 °C	A	-4.54	-4.54	-4.54	-4.54	-4.54	-4.54
	B	-5.45	-1.51	-5.45	-0.22	-5.45	-5.45
	C	-2.15	-2.67	-3.97	-5.27	-5.27	-5.27
90 °C	A	-4.54	-4.54	-4.54	-4.54	-4.54	-4.54
	B	-5.45	-5.45	-5.45	-5.45	-5.45	-5.45
	C	-3.97	-2.13	-5.27	-5.27	-5.27	-5.27

49 ^aValues are the difference between concentration (\log_{10} cfu g^{-1}) the day of artificial inoculation (T_0) and 24
50 hours (ΔT_1) and 30 days (ΔT_{30}) after treatment. For each batch and for each temperature-time combination
51 values are the minimum difference between the initial contamination level and the maximum count after the
52 treatment in the triplicate samples.

53 Table 5. Enumeration and detection of *L. monocytogenes* in ricotta salata artificially contaminated and heat treated with different temperature-time combinations
 54 and relative compliance with Regulation CE limits evaluated 24 h (T₁) and 30 days (T₃₀) after the heat treatment.

Treatment		<i>L.monocytogenes</i>						
Temperature	Minutes	Batch	T ₁		T ₃₀			
			Enumeration	Detection in 25 g	Enumeration	Detection in 25 g		
			+ve/n	log ₁₀ cfu/g	+ve /n	+ve/n	log ₁₀ cfu/g	+ve /n
75 °C	15	A	2/3	3.67±0.89	3/3 ¹	3/3	8.20±0.48 ²	3/3
		B	1/3	2.86±0.00	1/3 ¹	1/3	5.93±0.00 ²	3/3
		C	1/3	3.49±0.00	3/3 ¹	3/3	3.47± 0.48 ²	3/3
	25	A	0/3	0.00±0.00	1/3 ¹	2/3	6.39±0.58 ²	3/3
		B	0/3	0.00±0.00	1/3 ¹	0/3	0.00±0.00 ²	0/3
		C	0/3	0.00±0.00	1/3 ¹	1/3	3.54±0.00 ²	2/3
	40	A	2/3	1.30±0.00	3/3 ¹	3/3	6.87±0.34 ²	3/3
		B	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
		C	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
85 °C	15	A	0/3	0.00±0.00	0/3	0/3	0.00±0.00	1/3
		B	0/3	0.00±0.00	0/3	2/3	2.82±1.58 ²	2/3
		C	1/3	3.12± 0.00	2/3 ¹	1/3	2.60 ± 0.00 ²	3/3
	25	A	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
		B	0/3	0.00±0.00	0/3	1/3	5.23±0.00 ²	1/3
		C	1/3	1.30± 0.00	1/3 ¹	0/3	0.00±0.00	1/3
	40	A	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
		B	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
		C	0/3	0.0±0.00	0/3	0/3	0.00±0.00	0/3
90 °C	15	A	0/3	0.00±0.00	0/3	0/3	0.00±0.00	2/3
		B	0/3	0.00±0.00	0/3	0/3	0.00±0.00	1/3
		C	1/3	1.30± 0.00	3/3 ¹	2/3	2.92±0.22 ²	2/3
	25	A	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
		B	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
		C	0/3	0.00±0.00	0/3	0/3	0.00±0.00	2/3
	40	A	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
		B	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3
		C	0/3	0.00±0.00	0/3	0/3	0.00±0.00	0/3

55 Compliance are intended as follows: ¹not compliant with the Regulation CE 2073/2005 detection limits before the food has left the immediate control of the food
56 business operator (T_1); ²not compliant with the Regulation CE 2073/2005 enumeration limits for the products placed on the market during their shelf-life (T_{30}).

57

ACCEPTED MANUSCRIPT

58 Table 6. Intrinsic properties ($\bar{x} \pm SD$) and composition ($\% \pm SD$) of ricotta salata submitted to 9 different heat treatment combinations and analyzed 24 h (T_1) and
 59 30 days (T_{30}) after storage at refrigeration temperature.

Treatment		pH		a_w		Moisture %		Fat %		Proteins %		NaCl %	
Temperature	Min	T_1	T_{30}	T_1	T_{30}	T_1	T_{30}	T_1	T_{30}	T_1	T_{30}	T_1	T_{30}
75 °C	15	6.41±0.12 ^A	6.32±0.06 ^A	0.959±0.01 ^A	0.952±0.01 ^A	54.96±1.44 ^A	54.59±1.67 ^A	20.97±1.88 ^A	20.87±2.29 ^A	15.19±3.38 ^{AB}	14.75±1.35 ^{AB}	5.02±0.84 ^A	5.50±1.33 ^A
	25	6.42±0.11 ^A	6.36±0.05 ^A	0.954±0.01 ^A	0.951±0.01 ^A	54.90±2.08 ^A	54.94±1.36 ^A	21.73±2.01 ^{AB}	22.70±3.01 ^A	15.56±2.63 ^B	14.05±1.09 ^A	5.04±0.72 ^A	5.57±0.79 ^A
	40	6.41±0.13 ^A	6.35±0.07 ^A	0.950±0.01 ^A	0.654±0.01 ^A	55.09±2.23 ^A	53.53±1.93 ^A	20.90±3.11 ^A	22.71±2.32 ^A	14.44±1.29 ^{ABC}	14.56±0.93 ^{AB}	5.18±0.30 ^A	5.20±0.74 ^{AB}
85 °C	15	6.41±0.12 ^A	6.36±0.09 ^A	0.952±0.01 ^A	0.953±0.01 ^A	54.53±2.04 ^A	54.17±2.16 ^A	21.31±2.26 ^{AB}	21.65±2.52 ^A	14.20±2.32 ^{ABC}	15.01±1.54 ^{AB}	5.26±1.12 ^A	4.98±0.95 ^{AB}
	25	6.41±0.12 ^A	6.33±0.08 ^A	0.956±0.01 ^A	0.956±0.01 ^A	55.40±2.03 ^A	54.07±2.74 ^A	21.23±1.73 ^{AB}	22.29±3.13 ^A	13.97±1.04 ^{ABC}	15.37±1.12 ^{AB}	5.24±0.84 ^A	4.63±0.59 ^B
	40	6.38±0.10 ^A	6.33±0.11 ^A	0.953±0.01 ^A	0.956±0.01 ^A	54.72±1.81 ^A	53.91±2.12 ^A	21.65±2.21 ^{AB}	22.48±2.09 ^A	14.95±1.78 ^{ABC}	14.94±0.97 ^{AB}	5.16±0.76 ^A	5.09±0.76 ^{AB}
90 °C	15	6.40±0.11 ^A	6.33±0.09 ^A	0.953±0.01 ^A	0.954±0.01 ^A	54.30±1.64 ^A	53.61±1.62 ^A	23.07±0.88 ^B	22.75±2.15 ^A	13.33±0.56 ^C	14.48±1.50 ^{AB}	5.36±0.66 ^A	5.18±0.44 ^{AB}
	25	6.38±0.12 ^A	6.35±0.15 ^A	0.955±0.01 ^A	0.954±0.01 ^A	54.78±1.66 ^A	54.24±2.24 ^A	21.74±2.01 ^{AB}	22.52±2.36 ^A	14.38±2.00 ^{ABC}	14.33±0.82 ^{AB}	5.12±0.90 ^A	5.32±0.79 ^{AB}
	40	6.35±0.11 ^A	6.35±0.12 ^A	0.954±0.01 ^A	0.955±0.01 ^A	54.58±1.37 ^A	54.22±2.02 ^A	23.03±1.08 ^B	21.80±2.26 ^A	13.70±0.54 ^{AC}	15.50±2.66 ^B	5.19±0.87 ^A	5.03±0.93 ^{AB}


60 Each data point is the mean of three samples. For each parameter means in the same column on the same testing time (T_1 or T_{30}) with different capital letter are
 61 significantly different ($P < 0.05$).

62 Table 7. Mean values and standard deviations of sensory differences among the control (SEc) and the
63 samples heat treated (SEt).


Temperature-time condition	Differences
Blind samples	2.3 ^{a*} ± 1.5
75 °C x 15 min	3.4 ^{ab} ± 2.3
75 °C x 40 min	3.8 ^b ± 2.0
85 °C x 25 min	4.1 ^b ± 2.4
90 °C x 15 min	3.6 ^b ± 2.3
90 °C x 40 min	3.9 ^b ± 2.5

64 Mean values with different superscript letters are significantly different among samples.* ($P \leq 0.05$).

- 1 **Figure 1.** Reduction in *L. monocytogenes* counts (\log_{10} cfu g^{-1}) of artificially contaminated Ricotta salata wheels (T_0) analyzed 24 h (T_1) and 30
- 2 days (T_{30}) after waterbath heat treatment with 9 temperature-time combinations.



6 **Figure 2.** Temperatures recorded 1.5 cm below ricotta salata surface during water bath heat treatment



7

8 Each data point is the mean of temperatures recorder in the three replicates (batch A, B and C).

Highlights

1. Post-lethality treatment on *L. monocytogenes* was assessed in ricotta salata.
2. A reduction of $5 \log_{10}$ cfu g^{-1} in *L. monocytogenes* count was validated.
3. No effect of treatments on sensory properties was observed.