

Metazoan parasites in the head region of the bullet tuna *Auxis rochei* (Osteichthyes: Scombridae) from the western Mediterranean Sea

Questa è la versione Post print del seguente articolo:

*Original*

Metazoan parasites in the head region of the bullet tuna *Auxis rochei* (Osteichthyes: Scombridae) from the western Mediterranean Sea / Mele, S., Saber, S., Gómez Vives, M.j., Garippa, G., Alemany, F., Macías, D., Merella, P.. - In: JOURNAL OF HELMINTHOLOGY. - ISSN 0022-149X. - 89:(2015), pp. 734-739. [10.1017/S0022149X14000662]

*Availability:*

This version is available at: 11388/86362 since: 2022-05-27T10:30:13Z

*Publisher:*

*Published*

DOI:10.1017/S0022149X14000662

*Terms of use:*

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

*Publisher copyright*

note finali coverpage

(Article begins on next page)

Metazoan parasites in the head region of the bullet tuna *Auxis rochei* (Osteichthyes:  
Scombridae) from the western Mediterranean Sea

Salvatore Mele, Sámar Saber, María José Gómez-Vives, Giovanni Garippa, Francisco  
Alemany, David Macías, Paolo Merella

This is an Author's Original Manuscript of an article published by Cambridge University Press

in Journal of Helminthology on 12 September 2014, available online:

<http://dx.doi.org/10.1017/S0022149X14000662>

Metazoan parasites in the head region of the bullet tuna *Auxis rochei* (Osteichthyes:  
Scombridae) from the western Mediterranean Sea

Salvatore Mele<sup>1\*</sup>, Sámar Saber<sup>2</sup>, María José Gómez-Vives<sup>3</sup>, Giovanni Garippa<sup>1</sup>, Francisco  
Alemany<sup>4</sup>, David Macías<sup>3</sup>, Paolo Merella<sup>1</sup>

<sup>1</sup>Parassitologia e Malattie Parassitarie, Dipartimento di Medicina Veterinaria, Università di Sassari, via Vienna 2, 07100 Sassari, Italy; <sup>2</sup>Departamento de Biología Animal, Universidad de Málaga, Campus de Teatinos s/n, 29071 Málaga, Spain; <sup>3</sup>Centro Oceanográfico de Málaga, Instituto Español de Oceanografía, Puerto Pesquero s/n, 29640 Fuengirola, Spain; <sup>4</sup>Centre Oceanogràfic de les Balears, Instituto Español de Oceanografía, Moll de Ponent s/n, 07015 Palma, Spain

## Introduction

The bullet tuna *Auxis rochei* (Risso, 1810) (Osteichthyes: Scombridae) is a pelagic fish distributed in tropical and subtropical areas, including the Mediterranean Sea (Uchida, 1981). The limits of its distribution are not well known, mainly because this species is often confused with its congener, the frigate tuna *Auxis thazard* (Lacepède, 1800), another cosmopolitan fish (Di Natale *et al.*, 2009). In fact, the systematics of the genus *Auxis* is still controversial: some authors consider the species synonymous (Collignon, 1961; Nair *et al.*, 1970), while others recognise the existence of two distinct species (Yesaki & Arce, 1991; Collette & Aadland, 1996). Although at present the latter hypothesis is the one considered valid (Catanese *et al.*, 2008), recent genetic and morphometric studies showed that *A. rochei* is the only species distributed in the Mediterranean Sea and in the adjacent areas of the Atlantic Ocean (Orsi Relini *et al.*, 2008), suggesting that some occurrences of this species from this area could have been misidentified as *A. thazard*, even in scientific papers (Orsi Relini *et al.*, 2009).

*Auxis rochei*, which is the most abundant tuna in the Mediterranean Sea, represents an important component of the food web (Mostarda *et al.*, 2007) and is exploited by artisanal fisheries, representing 39% of the total tuna landings in the Mediterranean area (9829 t in 2010; FAO, 2011). The migrations of *A. rochei* in the Atlantic Ocean have been rarely studied; Grudtsev (1992) suggested that it may undertake a trophic migration along the north west coast of Africa, and Richards & Simmons (1971) suggested that spawning took place south of the Islands of Cape Verde. In addition, little is known about its life cycle and migrations in the Mediterranean Sea. Tortonese (1963) and Reglero *et al.* (2012) reported that *A. rochei* performs local migrations around the spawning areas in the neritic habitat of the mainland and islands, whereas Sabatés & Recasens (2001) proposed a spawning migration from the

Atlantic Ocean to the western Mediterranean Sea.

Parasites are useful tags to investigate the biology, ecology, migration and population structure of marine organisms (MacKenzie and Abaunza, 2014), and they have also been successfully used to clarify taxonomic relationships between hosts (Whittington, 2005). Although the parasites of the head region of representatives of the genus *Auxis* have been studied in several areas of the world, the confusion in the systematics of *A. rochei* and *A. thazard* makes specific information on their parasite fauna perplexing, e.g. in the Atlantic Ocean and Mediterranean Sea records of several parasites assigned to unidentified *Auxis* sp. (Dollfus, 1926; Palombi, 1949; Cressey & Cressey, 1980) could refer to *A. rochei*. The aim of this study is to describe the metazoan parasites of the head of *A. rochei* from the western Mediterranean Sea.

## Materials and methods

### *Collection and examination of fish*

A total of 63 *A. rochei* (32 males and 31 females), caught in the traditional trap fishery of La Azohía (37°32'59"N, 1°10'44"W, south east Spain, western Mediterranean Sea), were examined for parasites: 21 specimens were collected in May 2008 and 42 in May 2011. An additional group of nine specimens of bullet tuna (six males and three females), caught in the traditional trap fishery of Tarifa (36°00'59"N, 5°37'44"W, Strait of Gibraltar) in May 2008, were examined for comparative purposes. For each individual fish, the fork length to the nearest centimetre was recorded (FL = 34-44 cm).

The heads of fish were excised, stored individually in plastic bags and frozen at -20° C. Subsequently the samples were defrosted and examined for parasites according to Mele *et al.* (2012). The following literature was used for species identification: for monogeneans,

Palombi (1949), Fuentes Zambrano (1997), Mogrovejo & Santos (2002) and Mogrovejo *et al.* (2004); for didymozoids, Skrjabin (1955), Yamaguti (1970) and Pozdnyakov & Gibson (2008); for copepods, Vervoort (1962, 1965), Cressey & Cressey (1980), Boxshall & Halsey (2004) and Lin & Ho (2006).

### *Data analysis*

The prevalence of infection and mean abundance of each parasite species was calculated according to Bush *et al.* (1997). Confidence intervals of prevalence and mean abundance were assessed with the Sterne's exact method and the bias-corrected and accelerated Efron-Tibshirani bootstrap, respectively, using the free software Quantitative Parasitology 3.0 (Reiczigel & Rózsa, 2005).

Possible correlations between abundance of infection and host size were evaluated using the Spearman rank correlation coefficient, and its significance tested using the R software ("spearman" method, "cor.test" function, "stats" library of the R-software; R Development Core Team, 2014). Levels of infection of each parasite species were calculated for each of two size groups: (1), FL 33-37 cm; (2), FL 38-44 cm. Samples from south east Spain comprised four groups according to sampling year and host size: group (1) and (2) of 2008 ( $n = 11$  and  $22$ , respectively) and 2011 ( $n = 10$  and  $20$ , respectively). Samples from the Strait of Gibraltar of 2008 belonged only to the host size group (2) ( $n = 9$ ). Differences between the parasite infections of the five host groups and between host sexes were evaluated using the Fisher exact test for prevalence and the Welsh bootstrap t-test for mean abundance (Reiczigel & Rózsa, 2005).

Published and new data on the parasites of the head region of *Auxis* spp. worldwide were used to assess the dissimilarity between the parasite faunas of the head region of the

bullet and frigate tunas from four geographical regions: *A. thazard* from the Atlantic Ocean (data from: Vervoort, 1965; Fuentes Zambrano, 1997; Mogrovejo *et al.*, 2004; Chisholm & Whittington, 2007), *A. thazard* from the Indian Ocean (data from: Silas, 1962; Mogrovejo *et al.*, 2004; Chisholm & Whittington, 2007), *A. thazard* from the Pacific Ocean (data from: Silas, 1962; Yamaguti, 1970; Mogrovejo *et al.* 2004; Chisholm & Whittington, 2007), and *A. rochei* from the Mediterranean Sea (present results). Parasite fauna dissimilarity was evaluated with the Marczewski-Steinhaus distance (“cc” method, “betadiver” function, “vegan” library of the R-software) and the Bray-Curtis index (“-1” method, “betadiver” function, “vegan” library of the R-software) (Culurgioni *et al.*, 2014; Mele *et al.*, 2014).

## Results and discussion

This study is the first description of the parasite fauna of the head region of *A. rochei* from the western Mediterranean Sea. A total of seven parasite species were found in and on the hosts from south east Spain, with 72% being adult didymozoids belonging to two species, *Didymozoon auxis* Taschenberg, 1879 and Nematobothriinae gen. sp. (table 1). Up to 21% were monogeneans, including *Alloposeudaxine macrova* (Unnithan, 1957), *Churavera triangula* (Mamaev, 1967) and *Hexostoma auxisi* Palombi, 1943. The remaining 7% comprised two copepod species, *Caligus bonito* Wilson, 1905 and *Unicolax mycterobius* (Vervoort, 1965). Unidentified post-larval stages of didymozoids were found in the gill arch tissues. Only three parasite species were detected in *A. rochei* from the Strait of Gibraltar: *D. auxis* (86.7% of all specimens), *A. macrova* (6.7%) and *U. mycterobius* (6.6%). Only *D. auxis* and *H. auxisi* have been previously reported in *Auxis* sp. from the Mediterranean Sea (Dollfus, 1926; Palombi, 1949).

The unidentified Nematobothriinae gen. sp. has never been described before,

therefore it could be a specific parasite of *A. rochei* in the Mediterranean Sea, although the lack of previous records of this parasite could also be due to the difficulty of detection in the retrobulbar fat tissue. The other parasites have been reported in a wide range of hosts and regions: the congener *A. thazard* was found to harbour *C. triangula* in the Atlantic and Pacific Oceans and *H. auxisi* in the Atlantic Ocean only (Mogrovejo *et al.*, 2004); *A. macrova* infects several tunas such as *A. thazard*, *Euthynnus* spp., *Thunnus albacares* and *Katsuwonus pelamis* from the three oceans (Bussi eras & Baudin-Laurencin, 1973; Rohde *et al.*, 1980; Gibson *et al.*, 2005; Alves & Luque, 2006); *C. bonito* infects pelagic fish from the Mediterranean Sea (*Coryphaena hippurus*, *K. pelamis*; Carbonell *et al.*, 1999; Mele *et al.*, 2012), the north west Atlantic Ocean (*Thunnus thynnus*) and the tropical areas of the Atlantic, Indian and Pacific Oceans (*Euthynnus* spp., *K. pelamis* and *Sarda* spp.; Cressey & Cressey, 1980;  ktener & Trilles, 2009). In the Mediterranean Sea *A. macrova* seems to be restricted to *A. rochei*, since this monogenean was not found in 156 *E. alletteratus* and 35 *K. pelamis* caught within the same sampling program (Mele, 2013). The cross-infection of *A. macrova* among the Mediterranean tuna could be limited by the different life history and habitat of hosts (Reglero *et al.*, 2012; Rodr guez *et al.*, 2013).

No significant differences in prevalence and mean abundance were found ( $p > 0.05$ ) between host sexes and years of sampling; therefore the south east Spain data were pooled across sex and years, and the south east Spain hosts were only divided according to host size (table 1). The highest prevalence was that of *D. auxis* in the host size group (2) (71% in the south east Spain and 78% in the Strait of Gibraltar samples), being twice than in the host size group (1) (31% of the south east Spain samples). Significant differences between the mean abundance of three parasites were found: *A. macrova* had higher mean abundance in the host size group (2) from the south east Spain than in host size group (1); Nematobothriinae

gen. sp. had higher mean abundance in both host groups from the south east Spain than in that from the Strait of Gibraltar; and *C. bonito* had higher mean abundance in the host size group (2) from the south east Spain than in that from the Strait of Gibraltar (table 1). The higher prevalence of *D. auxis* and the higher mean abundance of *A. macrova* in the larger *A. rochei* than in the smaller ones could be due to the different origin of the fish. However, considering that *D. auxis* infects the definitive host through the food web, these differences could also indicate a change of the diet with size. In fact large fish (> 35 cm) can predate on fast-swimming prey including juvenile and adult fish and cephalopods (Mostarda *et al.*, 2007), which are among the intermediate hosts of didymozoids (Felizardo *et al.*, 2011). The difference of the mean abundance of Nematobothriinae gen. sp. and *C. bonito* according to geographical region could also be influenced by the small number of hosts sampled from the Strait of Gibraltar. The level of infection of Nematobothriinae gen. sp. did not change with host size (table 1). Considering that its geographical range is limited to the Mediterranean Sea and that the host cannot lose this tissue parasite with migration, the occurrence of Nematobothriinae gen. sp. only in the Mediterranean *A. rochei* could indicate that this host population is separated from that of *Auxis* spp. from other areas. However, the presence of this parasite could go unnoticed in routine parasitological analysis.

A negative relationship between the abundance of *U. mycterobius* and the host size ( $-0.27$ ,  $p = 0.003$ ) was found. This fact could be due to the increase of the distance between the narine walls in the larger hosts that will not offer an ideal microhabitat for the parasite to live wedged in the nasal cavities (Cressey & Cressey, 1980).

The parasite fauna of *A. thazard* from the Atlantic and Pacific Oceans has the highest richness, with nine and eight species, respectively. The poorest is the parasite fauna of *A. thazard* from the Indian Ocean, with only six species, although this could be due to the

scarcity of parasitological studies on this fish from this area. The parasite fauna of *A. rochei* from the Mediterranean Sea showed the closest similarity with that of *A. thazard* from the Atlantic Ocean, while the other groups were largely dissimilar (fig. 1). Considering the current debate on the host taxonomy and identification, the affinity between the parasite assemblages of the Mediterranean *A. rochei* and the Atlantic *A. thazard* could be compatible with the existence of a unique *Auxis* species throughout the Atlantic Ocean and the Mediterranean Sea (Orsi Relini *et al.*, 2009).

#### Acknowledgements

We wish to express our gratitude to the technicians of the laboratory of Mazarrón of the IEO *Centro Oceanográfico de Murcia*, Spain, to Enrique Majuelos for the assistance during the sampling in Tarifa, and to Rod A. Bray for the revision of the English style and grammar of the manuscript.

#### Financial support

This work was supported by the Spanish grants (GPM-3, GPM-4, PARATUN AGL2010-20892 and EC Data Collection Framework n. 199/2008) and the *Regione Autonoma della Sardegna* grant (S. M., Master and Back 2012-2013 PRR-MAB-A2013-17605).

## References

- Alves, D.R. & Luque, J.L. (2006) Ecologia das comunidades de metazoários parasitos de cinco espécies de escombrídeos (Perciformes: Scombridae) do litoral do estado do Rio de Janeiro, Brasil. *Revista Brasileira de Parasitologia Veterinária* 15, 167–181.
- Boxshall, G.A. & Halsey, S.H. (2004) *An introduction to copepod diversity*. 966 pp. London, The Ray Society.
- Bush, A.O., Lafferty, K.D., Lotz, J.M. & Shostak, A.W. (1997) Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology* 83, 575–583.
- Bussiéras, J. & Baudin-Laurencin, F. (1973) Les helminthes parasites des thons tropicaux. *Revue d'Élevage et de Médecine Vétérinaire des Pays Tropicaux* 26, 13–19.
- Carbonell, E., Massutí, E., Castro, J.J. & García, R.M. (1999) Parasitism of dolphinfishes, *Coryphaena hippurus* and *Coryphaena equiselis*, in the western Mediterranean (Balearic Islands) and central-eastern Atlantic (Canary Islands). *Scientia Marina* 63, 343–354.
- Catanese, G., Infante, C. & Manchado, M. (2008) Complete mitochondrial DNA sequences of the frigate tuna *Auxis thazard* and the bullet tuna *Auxis rochei*. *DNA Sequence* 19, 159–166.
- Chisholm, L.A. & Whittington, I.D. (2007) Review of the Capsalinae (Monogenea: Capsalidae). *Zootaxa* 1559, 1–30.
- Collette, B.B. & Aadland, C.R. (1996) Revision of the frigate tunas (Scombridae, *Auxis*), with descriptions of two new subspecies from the eastern Pacific. *Fishery Bulletin* 94, 423–441.
- Collignon, J. (1961) Le thazard ou melva dans l'Atlantique oriental. *Bulletin de l'Institut des Pêches Maritimes du Maroc* 7, 55–72.
- Culurgioni, J., Mele, S., Merella, P., Addis, P., Figus, V., Cau, A., Karakulak, F.S. & Garippa, G. (2014) Metazoan gill parasites of the Atlantic bluefin tuna *Thunnus thynnus* (Linnaeus) (Osteichthyes: Scombridae) from the Mediterranean and their possible use as biological tags. *Folia Parasitologica* 61, 148–156.
- Cressey, R. & Cressey, H.B. (1980) Parasitic copepods of mackerel and tuna-like fishes (Scombridae) of the world. *Smithsonian Contributions to Zoology* 311, 1–186.
- Di Natale, A., Srour, A., Hattour, A., Keskin, Ç., Idrissi, M. & Orsi Relini, L. (2009) Regional study on small tunas in the Mediterranean including the Black Sea. *Studies and Reviews FAO* 85.

- Dollfus, R.P. (1926) Sur l'état actuel de la classification des Didymozoonidae Monticelli, 1888 (Didymozoidae Franz Poche, 1907). *Annales de Parasitologie Humaine et Comparée* 4, 148–161.
- FAO (2011). *FishStatJ: Universal software for fishery statistical time series*. FAO Fisheries and Aquaculture Department, Statistics and Information Service, Rome, Italy.
- Felizardo, N.N., Justo, M.C., Knoff, M., Fonseca, M.C.G., Pinto, R.M. & Gomes, D.C. (2011) Juvenile didymozoids of the types, Torticaecum and Neotorticaecum (Didymozoidae: Digenea), from new marine fish hosts (Pisces: Teleostei) in the neotropical region of Brazil. *Journal of Helminthology* 85, 270–275.
- Fuentes Zambrano, J.L. (1997) *Neohexostoma mochimae* n. sp. y *Pseudochauhanea elegans* n. sp. (Monogenea) dos nuevas especies de parásitos de peces de la Bahía de Mochima, Venezuela. *Boletín del Instituto Oceanográfico de Venezuela* 36, 45–52.
- Gibson, D.I., Bray, R.A. & Harris, E.A. (2005) *Host-parasite database of the Natural History Museum*, London. URL
- Grudtsev, M.E. (1992) Particularités de répartition et caractéristique biologique de la melva *Auxis rochei* (Risso) dans les eaux du Sahara. *Collective Volume of Scientifics Papers ICCAT* 39, 284–288.
- Lin, C. & Ho, S. (2006) Four species of *Unicolax* Cressey and Cressey, 1980 (Copepoda: Bomolochidae) parasitic on marine fishes of Taiwan. *Zoological Studies* 45, 339–356.
- Mackenzie, K. & Abaunza, P. (2014) Parasites as biological tags. pp. 185–203 in Cadrin, S.X., Kerr, L.A. & Mariani, S. (Eds) *Stock identification methods*. 2nd edn. San Diego, Academic Press.
- Mele, S. (2013) *Gill metazoan parasites of tunas (Scombridae: Thunnini) from the western Mediterranean Sea: systematics, assemblages and use as biological tags*. PhD thesis, University of Sassari.
- Mele, S., Macías, D., Gómez, M.J., Garippa, G., Alemany, F. & Merella, P. (2012) Metazoan parasites on the gills of the skipjack tuna *Katsuwonus pelamis* (Osteichthyes: Scombridae) from the Alboran Sea (western Mediterranean Sea). *Diseases of Aquatic Organisms* 97, 219–225.
- Mele, S., Pennino, M.G., Piras, M.C., Bellido, J.M., Garippa, G. & Merella, P. (2014) Parasite of the head of *Scomber colias* (Osteichthyes: Scombridae) from the western Mediterranean Sea. *Acta Parasitologica* 59, 173–183.

- Mogrovejo, C.D. & Santos, C. (2002) *Caballerocotyla lenti* n. sp., a capsalid monogenean from *Auxis thazard* (Scombridae) from off the Southeastern Coast of Brazil. *Memórias do Instituto Oswaldo Cruz* 97, 1067–1071.
- Mostarda, E., Campo, D., Castriota, L., Esposito, V., Scarabello, M. P. & Andaloro, F. (2007) Feeding habits of the bullet tuna *Auxis rochei* in the southern Tyrrhenian Sea. *Journal of the Marine Biological Association of the United Kingdom* 87, 1007–1012.
- Nair, R.V., Virabhadra Rao, K. & Dorairaj, K. (1970) The tunas and tuna-like fishes of India. *Bulletin of the Central Marine Fisheries Research Institute* 23, 1–94.
- Öktener, A. & Trilles, J.P. (2009) Four parasitic copepods on marine fish (Teleostei and Chondrichthyes) from Turkey. *Acta Adriatica* 50, 121–128.
- Orsi Relini, L., Palandri, G., Garibaldi, F., Lanteri, L., Cilli, G., Ferrara, G. & Tinti, F. (2008) Towards a new taxonomical approach to Mediterranean small tuna of genus *Auxis*. *Biologia Marina Mediterranea* 15, 207–210.
- Orsi Relini, L., Palandri, G., Garibaldi, F., Lanteri, L. & Tinti, F. (2009) Between lumpers and splitters, which taxonomical approach to Mediterranean small tunas of genus *Auxis*? *Collective Volume of Scientific Papers ICCAT* 64, 2200–2210.
- Palombi, A. (1949) I trematodi d'Italia. Parte I. Trematodi monogenetici. *Archivio Zoologico Italiano* 34, 204–408.
- Pozdnyakov, S.E. & Gibson, D.I. (2008) Family Didymozoidae Monticelli, 1888. pp.631–734 in Bray, R. A., Gibson, D. I. & Jones, A. (Eds) *Keys to the Trematoda*, vol. 3. Wallingford, CABI Publishing and the Natural History Museum.
- R Development Core Team (2014) *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. Available from: [http://wwwR-project.org](http://www.R-project.org).
- Reglero, P., Ciannelli, L., Álvarez-Berastegui, D., Balbín, R., López-Jurado, J. L. & Alemany, F. (2012) Geographically and environmentally driven spawning distributions of tuna species in the western Mediterranean Sea. *Marine Ecology Progress Series* 463, 273–284.
- Reiczigel, J. & Rózsa, L. (2005) *Quantitative parasitology 3.0*, Budapest. Distributed by the authors. Available from: <http://www.zoologia.hu/qp/>.
- Richards, W.J. & Simmons, D.C. (1971) Distribution of tuna larvae (Pisces, Scombridae) in the northwestern Gulf of Guinea and off Sierra Leone. *Fishery Bulletin* 69, 555–568.

- Rodríguez, J.M., Álvarez, I., López-Jurado, J.L., García, A., Balbin, R., Álvarez-Berastegui, D., Torres, A.P. & Alemany, F. (2013) Environmental forcing and the larval fish community associated to the Atlantic bluefin tuna spawning habitat of the Balearic region (Western Mediterranean), in early summer 2005. *Deep-Sea Research* 77, 11–22.
- Rohde, K., Roubal, F. & Hewitt, G. C. (1980) Ectoparasitic Monogenea, Digenea and Copepoda from the gills of some fishes of New Caledonia and New Zealand. *New Zealand Journal of Marine and Freshwater Research* 14, 1–13.
- Sabatés, A & Recasens, L. (2001) Seasonal distribution and spawning of small tunas (*Auxis rochei* and *Sarda sarda*) in the northwestern Mediterranean. *Scientia Marina* 65, 95–100.
- Silas, E.G. (1962) Parasites of scombroid fishes, Part 1, Monogenetic trematodes, digenetic trematodes, and cestodes. *Proceedings of the Symposium on Scombroid Fishes, Part 3*, Marine Biological Association of India, 12–15 January 1962, Mandapam, India.
- Skrjabin, K.I. (1955) Suborder Didymozoata Skjrabin et Schulz, 1937. *Osnovy Trematodology* 11, 5–254.
- Tortonese, E. (1963) La popolazione Mediterranea di *Auxis* (Pisces Thunnidae) in rapporto alla sistematica del genere. *Annali del Museo Civico di Storia Naturale Giacomo Doria* 74, 140–155.
- Uchida, R.N. (1981) Synopsis of biological data on frigate tuna, *Auxis thazard*, and bullet tuna, *A. rochei*. *FAO Fisheries Synopsis* 124.
- Vervoort, W. (1962) A review of the genera and species of the Bomolochidae (Crustacea, Copepoda), including the description of some old and new species. *Zoologische Verhandelingen* 56, 1–111.
- Vervoort, W. (1965) Three new species of Bomolochidae (Copepoda, Cyclopoida) from tropical Atlantic tunnies. *Zoologische Verhandelingen* 76, 3–40.
- Whittington, I.D. (2005) Monogenea Monopisthocotylea (ectoparasitic flukes). pp. 63–71 in Rohde, K. (ed) *Marine parasitology*. Wallingford, CABI Publishing.
- Yamaguti, S. (1970) *Digenetic trematodes of Hawaiian fishes*. Tokyo, Keigaku Publishing Co.
- Yesaki, M. & Arce, F. (1991) A review of the *Auxis* fisheries of the Philippines and some aspects of the biology of frigate (*A. thazard*) and bullet (*A. rochei*) tunas in the Indo-Pacific region. *FAO Fisheries Technical Paper* 336, 409–439.

Table 1. Prevalence (%), mean abundance (MA), 95% confidence intervals (CI) and microhabitat of the metazoan parasites in the head region of *Auxis rochei* from south east Spain and the Strait of Gibraltar; FL – fork length, na – no data given, \* and ^- level of significant differences between pairs of samples given as  $p \leq 0.05$ .

Geographical region	South east Spain				Strait of Gibraltar		Microhabitat
	FL 33 - 37 cm		FL 38 - 44 cm		FL 38 - 44 cm		
Host size class							
Parasite	% (CI)	MA (CI)	% (CI)	MA (CI)	% (CI)	MA (CI)	
Monogenea							
<i>Alloposeudaxine macrova</i>	5(0-23)	0.1 (0-0.1)*	24 (13-39)	0.3 (0.1-0.4)*	11 (1-44)	0.1 (0.0-0.3)	Gill filaments
<i>Churavera triangula</i>	5(0-23)	0.1 (0-0.1)	7 (2-20)	0.1 (0.0-0.3)	0 (0-32)	0.0 (na)	Gill filaments
<i>Hexostoma auxisi</i>	14 (4-35)	0.9 (0.1-3.9)	2 (0-13)	0.1 (0.0-0.1)	0 (0-32)	0.0 (na)	Gill filaments
Didymozoidae							
<i>Didymozoon auxis</i>	33 (16-55)*^	1.4 (0.4-4.1)	71 (56-84)*	2.2 (1.4-3.3)	78 (44-96)^	1.4 (0.7-2.3)	Gill filaments
Nematobothriinae gen. sp.	24 (10-46)	0.3 (0.1-0.4)*	24 (13-39)	0.3 (0.2-0.6)^	0 (0-32)	0.0 (na)*^	Retrobulbar fat tissue
Copepoda							
<i>Caligus bonito</i>	5(0-23)	0.1 (0-0.1)	17 (8-31)	0.2 (0.1-0.3)*	0 (0-32)	0.0 (na)*	Gill chamber and mouth
<i>Unicolax mycterobius</i>	20 (7-47)	0.2 (0.0-0.4)	5 (0-16)	0.1 (0.0-0.1)	11 (1-44)	0.1 (0.0-0.3)	Nasal cavities

## Legend of the figure

Fig. 1. Cluster dendrograms (group-average linkage) of the parasites of the head of *Auxis rochei* from the western Mediterranean Sea (Ar-Med), and of *A. thazard* from the Atlantic Ocean (At-Atl), Indian Ocean (At-Ind) and Pacific Ocean (At-Pac), using Marczewski-Steinhaus (A) and Bray-Curtis (B) dissimilarity measures based on the presence/absence data with the cophenetic correlation coefficients having a similar value of 0.89. Source: Silas (1962); Vervoort (1965); Yamaguti (1970); Cressey & Cressey (1980); Muruges & Madhavi (1995); Fuentes Zambrano (1997); Mogrovejo *et al.* (2004); Chisholm & Whittington (2007); and present results.

