



Review

# Thyroid Hormone Diseases and Osteoporosis

Alessandro P. Delitala \*, Angelo Scuteri and Carlo Doria

Department of Clinical and Experimental Medicine, University of Sassari, 07100 Sassari, Italy; d341elefante@virgilio.it (A.S.); cdoria@uniss.it (C.D.)

\* Correspondence: aledelitala@tiscali.it; Tel.: +39-079228184

Received: 19 February 2020; Accepted: 3 April 2020; Published: 6 April 2020



**Abstract:** Thyroid hormones are essential for normal skeletal development and normal bone metabolism in adults but can have detrimental effects on bone structures in states of thyroid dysfunction. Untreated severe hyperthyroidism influences the degree of bone mass and increases the probability of high bone turnover osteoporosis. Subclinical hyperthyroidism, defined as low thyrotropin (TSH) and free hormones within the reference range, is a subtler disease, often asymptomatic, and the diagnosis is incidentally made during screening exams. However, more recent data suggest that this clinical condition may affect bone metabolism resulting in decreased bone mineral density (BMD) and increased risk of fracture, particularly in postmenopausal women. The main causes of exogenous subclinical hyperthyroidism are inappropriate replacement dose of thyroxin and TSH suppressive L-thyroxine doses in the therapy of benign thyroid nodules and thyroid carcinoma. Available data similarly suggest that a long-term TSH suppressive dose of thyroxin may decrease BMD and may induce an increased risk of fracture. These effects are particularly observed in postmenopausal women but are less evident in premenopausal women. Overt hypothyroidism is known to lower bone turnover by reducing both osteoclastic bone resorption and osteoblastic activity. These changes in bone metabolism would result in an increase in bone mineralization. At the moment, there are no clear data that demonstrate any relationship between BMD in adults and hypothyroidism. Despite these clinical evidences, the cellular and molecular actions of thyroid hormones on bone structures are not complete clear.

**Keywords:** thyroid dysfunctions; subclinical hypothyroidism; subclinical hyperthyroidism; osteoporosis; bone mass density

## 1. Introduction

Osteoporosis is by far the most common metabolic bone disease and is characterized by low bone mass with microarchitectural alteration of bone structure, leading to reduced bone strength, which predisposes to an increased fracture risk. The definition of osteoporosis is based on densitometric measurement of bone mass in the spine or hip and not on clinical criteria. The World Health Organization defines osteoporosis as a bone mineral density (BMD) 2.5 or more standard deviations (SDs) below that of a young adult at any site (T score), whereas osteopenia is defined when BMD is  $>1$  SD and  $<2.5$  SD lower than the young- adult mean end [1].

Decreased bone mass and increased fragility in age-related osteoporosis can occur because of failure to achieve optimal peak bone mass. Although genetic determinants are responsible for up to 85% of the variation in peak bone mass, other factors during childhood and adolescence can affect the ability to achieve optimal peak bone mass. These variables include nutrition, particularly calcium intake, delayed puberty, physical activity, a wide variety of intercurrent illnesses, and social factors such as low family income [2].

Peak bone mass is usually achieved at the end of the second decade. The progressive bone loss is due to an increased bone resorption, which is the major mechanism for increased bone fragility.

Any increase in the number of resorption sites produces decreased bone mass together with changes in skeletal microarchitecture, since the mechanism of resorption requires a much shorter time than that of bone formation in the bone remodeling cycle. Thus, the amount of bone formation decreases with age, leading to a more fragile skeleton. Although many variables may be involved in this process, decrease in bone formation may also be due to an age-related decline in skeletal growth factors [3]. Postmenopausal osteoporosis is a function of aging and estrogen deficiency. It is estimated that about 75% of bone loss during the first 15 years after menopause is attributed to estrogen deficiency rather than to aging [4]. The bone loss is more evident in vertebral bodies since the trabecular component is metabolically very active and decreases significantly when estrogen is deficient. The peak bone mass achieved at maturity and the age of menopause are strong predictors of an increased risk for fracture. Bisphosphonates are the cornerstone for the treatment of osteoporosis [5] and are commonly prescribed for its treatment, both in women and men [6]. More recently, other drugs (i.e., teriparatide and denosumab) are often prescribed and showed a safety profile [7], even when prescribed for therapy of non-unions [8]. However, a surgical approach sometimes necessary, in particular, in the case of vertebral compression fractures [9].

Thyroid hormones have multiple effects on body homeostasis [10,11] and metabolism [12], and their increased concentration may lead to different complications, mainly at the cardiovascular level [13]. Thyroid hormones are also essential for skeletal maturation and have an important physiological role in the maintenance of adult bone structure and strength. Although thyroid dysfunction has been known to represent a risk factor for bone disease, the role of thyroid hormone excess or deficiency in the pathogenesis of osteoporosis and risk factors of fractures has been underestimated, and the underlying mechanisms are still uncertain.

In this review, we summarize the systemic and local effects of thyroid hormone action and the bone consequence of long-term effects of thyroid dysfunction including the subclinical forms of these diseases.

## 2. Materials and Methods

We performed on PubMed a literature search for the articles published until January 2020 by using the search terms “subclinical hyperthyroidism”, “osteoporosis”, “bone mass density”, and “subclinical hypothyroidism”. Titles of interest were further reviewed by abstract.

## 3. Thyroid Hormone Physiology

The main thyroid hormones in humans are thyroxine (T4) and 3,5,3'-triiodo-L-thyronine (T3). The synthesis and secretion of these hormones are finely regulated by the thyroid-stimulating hormone (TSH) axis, whose activity is negatively regulated by thyroid hormones and cytokines [14]. TSH acts directly on the TSH receptor (TSH-R) expressed on the thyroid follicular cell basolateral membrane [15]. The nuclear genomic effect of thyroid hormones is mediated by the intracellular binding of T3 to nuclear receptor where it activates either thyroid hormone receptor  $\alpha$  (TR $\alpha$ ) or  $\beta$  (TR $\beta$ ). TRs act as a hormone-dependent transcription factor that mediates transcriptional repression in the unliganded state. T3 binding results in the dissociation of co-repressors and the recruitment of co-activators resulting in stimulation of gene transcription [16]. Both TR $\alpha$  and TR $\beta$  are widely expressed in a tissue-specific manner. TR $\beta$  is the main receptor expressed in the pituitary and hypothalamus where it mediates the negative feedback control of the hypothalamic–pituitary–thyroid axis [17], whereas TR $\alpha$  is expressed in higher concentration than TR $\beta$  in the skeleton, where it mediates T3 action on bone and cartilage [18]. The knockout of the thyroid hormone TR $\alpha$  gene results in delayed bone maturation, while the lack of all isoforms of TR $\beta$  has no effect on bone cells [18].

The relation between thyroid hormones and bone metabolism is well documented in children with hypothyroidism. In severe untreated hypothyroidism, delayed skeletal development, defective endochondral ossification, short stature, and epiphyseal dysgenesis are well known clinical features that can be reversed particularly when early and prompt replacement therapy is started.

Thyroid hormone is crucial for cartilage growth and differentiation and enhances the response to growth hormone.

In adults, overt hypothyroidism induces a low bone turnover with a prolonged bone remodeling cycle, which is caused by reduced osteoclastic bone resorption together with decreased osteoblastic activity. On the contrary, in states of hyperthyroidism, there is a high bone turnover with a shortened remodeling time. The discrepancy between bone formation and resorption results in a negative balance with a net loss of bone.

Nuclear T3 receptor has been found in osteoblastic cell lines [19] as well as in osteoclastoma-derived osteoclasts [20]. Moreover, T3 directly stimulates bone resorption *in vitro* [21] possibly through the involvement of interleukin 6, a potent stimulator of osteoclastic activity. An additional role in bone metabolism has been proposed for TSH, since the TSH receptor, although predominantly expressed in thyroid follicular cells, has been described in other tissues including osteoblasts and osteoclasts [22]. TSH has been suggested as a key negative regulator of bone turnover, with a direct effect on osteoblastic bone resorption due to decreased local production of tumor necrosis factor- $\alpha$  [23]. However, this hypothesis did not explain the increased risk of osteoporosis reported in patients with Basedow's disease. Indeed, these patients have increased levels of circulating TSH receptor autoantibodies that stimulate the TSH receptor. Thus, another study showed that a high dose of TSH was unable to affect the differentiation or function of both osteoblasts and osteoclasts *in vitro* [24]. These data would suggest that the skeletal abnormalities found in hypothyroidism are independent of systemic TSH. Studies on humans are limited to the effect of different TSH concentration on serum markers of bone metabolism. Interesting studies analyzed the effect of acute administration of recombinant human TSH (rhTSH) in patients without thyroid glands and with constant free thyroxine (FT4) level due to their L-thyroxine (L-T4) supply. This test is commonly used as a part of follow-up in patients with differentiated thyroid cancer and gives the opportunity to test *in vivo* the effect of increased concentration of TSH on bone turnover markers. One study reported an increase of N-terminal propeptide of type-I procollagen, an index of osteoblastic activity, and an increase of serum receptor activation for nuclear factor  $\kappa$ B ligand (RANKL) in postmenopausal women but not in premenopausal women, suggesting a possible role of estrogen on bone reactivity to TSH [25]. On the other hand, another study reported no effect of rhTSH on serum osteoprotegerin and RANKL [26].

#### 4. Bone Physiology

The skeleton is one of the largest organs in the body and has multiple physiological actions including the structural strength and integrity of the body, having an important role in the maintenance of normal serum level of calcium and phosphate. Bone consists of a mineralized matrix and a highly metabolic active cellular component. Cortical bone is a dense bone, which represents up to 80% of skeletal mass, and is present in the shafts of long bones. The bone remodeling cycle is initiated and orchestrated by osteocytes, which are embedded within mineralized bone and communicate via ramifications of dendritic processes. Bone remodeling may result from changes in mechanical load, structural damage, or exposure to systemic or paracrine factors. Hemopoietic cells of the monocyte/macrophage lineage differentiate to mature osteoclasts and resorb bone [27]. During the reversal phase, osteoblastic progenitors are recruited to the site of resorption, differentiate and synthesize osteoids, and mineralize the new bone matrix to repair the defect. Crosstalk between bone-forming osteoblasts and bone-resorbing osteoclasts regulates bone remodeling and maintains skeletal homeostasis [18].

Bone is formed by osteoblasts, which are derived from mesenchymal cells in the skeletal microenvironment; they are connected by gap junctions and secrete collagen and non-collagen proteins [28]. Mature osteoblasts may die by apoptosis or become osteocytes, which are embedded in the matrix or are transformed into flattened lining cells, which cover a large proportion of the bone surface. Although osteocytes may still synthesize collagen and other proteins, they have an important role in bone repair by providing active molecules for the initiation of bone remodeling at the site of

bone damage. This effect is probably mediated by mechanical forces that produce a fluid shear stress in the canaliculi between osteocytes, thereby inducing intracellular signals, which result in the secretion of active molecules. Most of the molecules, including hormonal factors, act on osteoblasts for initiating bone resorption. In particular, osteoblasts secrete RANKL (receptor activation for nuclear factor  $\kappa$ B ligand) and CSF1 (colony-stimulating factor 1), and other factors such as cytokines, prostaglandins, and growth factor that regulate bone resorption and are crucial for osteoclastogenesis [29]. The most important stimulator of osteoclastic formation is RANKL, a member of the Tumor Necrosis Factor (TNF) protein superfamily. All factors involved in osteoclast genesis stimulate RANKL by osteoblastic cell lines. Osteoclastic formation is inhibited by osteoprotegerin (OPG), which is a soluble receptor for RANKL, thereby preventing interaction of RANKL with its receptor RANK, a member of the TNF receptor family [30].

Resorption and formation of bone occurs throughout life and is a coordinated process that is regulated by basic multicellular units, that are activated to maintain skeletal strength and repair bone microdamage [31]. The bone remodeling cycle, which is influenced by systemic hormones and local factors, is characterized by activation, resorption, reversal, and formation steps, and this turnover is greater in cancellous than in cortical bone. In this cycle in young adults, the amount of new bone formed in the resorptive cavity is equal to the amount resorbed by osteoclasts. When the activity of osteoclasts is enhanced and/or that of osteoblasts is reduced, there is a net loss of bone with an increased risk of osteoporosis [32].

Although a direct action of T3 on osteoblasts and chondrocytes is recognized, the effect of T3 on osteoclasts is still unclear [33]. Thyroid hormones might have a direct effect on osteoclasts, or their action on bone resorption might be mediated by osteoblasts or other cell types.

The most widely used method for measuring bone mass is dual-energy X-ray absorptiometry (DXA). The technic provides accurate values for bone mineral content (BMC) and BMD in the lumbar spine, the proximal femur, the distal radius, and the whole body, with minimal radiation exposure. BMD is calculated from BMC and the area of bone scanned ( $\text{g}/\text{cm}^2$ ). Ultrasonography, particularly in the calcaneous, is a rapid procedure, which does not use X-rays and may predict fracture risk. Data of both techniques are reported in terms of T scores (i.e., standard deviations from the young adult values) or Z scores (i.e., standard deviation from the expected normal values). DXA scanners are also used for vertebral fracture assessment, which offer a high degree of accuracy in diagnosing fractures [34]. Other methods include quantitative computed tomography (QCT), which is likely more accurate than DXA.

## 5. Hyperthyroidism

The main causes of hyperthyroidism are Graves' disease, toxic multinodular goiter, and toxic adenoma (Table 1). Toxic multinodular goiter is the most frequent cause of spontaneous hyperthyroidism in areas with a low iodine intake and hyperthyroidism may also be precipitated by excess iodine intake from drugs or radiographic contrast agents. Overt thyroid disorders, defined as a suppression of TSH with increased FT4 and/or free triiodothyronine (FT3) are clinically evident and an early diagnosis is usually made. Due to its deleterious effect on the cardiovascular system [35], treatment of hyperthyroidism is always necessary to ensure a rapid relief of symptoms and to avoid long-term consequences. For these reasons, severe long-lasting hyperthyroidism is now rarely encountered in clinical practice. It is well known that overt hyperthyroidism has a detrimental effect on bone mass and fragility fractures due to a high bone turnover as documented by a shortened bone remodeling cycle, together with an increase in biochemical markers of bone resorption and bone formation [18].

**Table 1.** Causes of thyrotoxicosis.

<b>Hyperthyroidism (hormone overproduction)</b>
Graves' disease Toxic multinodular goiter, toxic adenoma Iodide-induced hyperthyroidism (Jod-Basedow effect) Amiodarone-associated hyperthyroidism due to iodide release TSH-secreting pituitary tumors
<b>Thyrotoxicosis (transient hormone excess)</b>
Autoimmune thyroiditis Subacute thyroiditis Drug-induced thyroiditis (amiodarone, lithium, interferon alfa, interleukin 2, thyrosine kinase inhibitors)
<b>Exogenous thyroid hormone</b>
Iatrogenic over-replacement Thyrotoxicosis factitia

The accelerated remodeling cycle causes an increased release of calcium into the systemic circulation [36]. High levels of calcium reduce parathyroid hormone secretion leading to an increased urinary calcium loss and a negative calcium balance. The conversion of vitamin D into its active form is also reduced by low parathormone thus reducing gastrointestinal calcium absorption and resultant fecal calcium losses.

Some studies demonstrated 12–20% reduction in the BMD of hyperthyroid subjects [37], while other authors found no difference in BMD between thyrotoxic patients and euthyroid subjects [38]. A meta-analysis of 25 studies reported by Vestergaard [39] showed that BMD was decreased in untreated patients with hyperthyroidism, with an increased risk of hip fracture, which increases significantly with age. This aspect is more evident in postmenopausal women with a three-to-fourfold increase in fracture [40]. Similar results have been reported by Jodar et al. both in postmenopausal women with Graves' disease and in those with toxic nodular goiter [41]. BMD was also evaluated in geriatric patients with toxic nodular goiter [42]. Both women and men had a BMD that was significantly decreased compared to the control. Longitudinal studies also evaluated whether bone loss could be reversed by treatment of hyperthyroidism. A study by Dhanwal et al. reported an early recovery at hip and lumbar spine after eight weeks of carbimazole therapy [43]. Another study reported that the increased level of OPG found in hyperthyroid, normalized after medical treatment, even in the presence of persistent abnormal bone structure [44]. Collectively, all the data demonstrate that the severity of hyperthyroidism appears to influence the degree of bone mass and increase the probability of osteoporosis. Moreover, the majority of studies in hyperthyroid postmenopausal women demonstrated a reduction in BMD. Furthermore, a prior history of hyperthyroidism is an independent risk factor for hip and vertebral fracture (relative risk 1.8) [45]. Data seem also to suggest that treatment of hyperthyroidism has a beneficial effect on bone metabolism, but studies are too scanty to draw firm conclusions.

## 6. Subclinical Hyperthyroidism

Subclinical disorders of the thyroid gland, defined as TSH outside the reference range of values and FT3 and FT4 within the normal range, are more subtle diseases and the diagnosis is incidentally made during screening exams. The availability of sensitive assay for TSH has allowed recognition of a syndrome in which usually there are no signs or symptoms of thyrotoxicosis, but the serum TSH is subnormal. Subclinical hyperthyroidism, defined as low TSH and free hormones within the reference range, has gained attention in the last years for its association with cardiovascular disease (CVD), in particular with atrial fibrillation in patients with a serum TSH concentration of 0.1 mU/L or less. The incidence of subclinical hyperthyroidism increases with age, especially in women, and is present in about of 15% of women over the age of 60 [46]. Serum TSH may also progress to overt hyperthyroidism,

particularly, in patients with autonomous thyroid nodules or multinodular goiter. A reduction in serum TSH can occur with euthyroid Graves' disease, subacute thyroiditis, autonomous adenoma, multinodular goiter, or administration of amounts of thyroid hormone greater than that required for metabolic needs. Exogenous hyperthyroidism is due to iatrogenic over-replacement with thyroid hormone supplementation. This condition is present when TSH suppressive therapy is prescribed in patients with differentiated carcinoma of thyroid or with benign uninodular or multinodular goiter.

## 7. Endogenous Subclinical Hyperthyroidism

The possibility that subclinical hyperthyroidism is involved in fracture risk has been investigated in a number of studies on bone turnover BMD measurement or fracture risk in pre- and postmenopausal women and in men [47], as reported in Table 2. No effect of endogenous subclinical hyperthyroidism on bone turnover was reported by Gurlek and Gedik in fifteen premenopausal women with a stable TSH suppression during a period ranging between 6 and 11 months [48]. Persistent increase in bone turnover in Graves' patients with subclinical hyperthyroidism was observed in other studies. However, no data on bone density were studied [49]. Women older than 65 years of age who have suppressed TSH have been reported to have a threefold increased risk for hip fractures and a fourfold increased risk for vertebral fracture [41]. Decreased BMD was observed in a small group of patients with subclinical hyperthyroidism with nodular goiter [50]. Similar data have been reported in pre- and postmenopausal women with subclinical hyperthyroidism caused by multinodular goiter. BMD and bone turnover markers were decreased and respectively higher in post- than in premenopausal ones [51]. A cross-sectional study on data collected in the third U.S. National Health and Nutrition Examination Survey (NHANES III) showed an association between low serum TSH and osteopenia and osteoporosis. Interestingly, a graded increase in BMD with increased serum TSH across the normal range was observed in healthy American women of both black and white races [52]. Low bone density together with high biochemical markers of bone turnover has been confirmed in additional studies [53–56], although no association between subclinical hyperthyroidism and BMD or hip fractures in older men and women was found in a large cohort study of community-dwelling individuals 65 years old or older enrolled in the Cardiovascular Health Study [57].

**Table 2.** Studies that analyzed the effect of endogenous subclinical hyperthyroidism on lumbar spine bone mass density and femoral bone mass density.

Study	n	Gender	Age (Years)	Lumbar Spine BMD (g/cm <sup>2</sup> )	Lumbar Spine T Score	Femur BMD (g/cm <sup>2</sup> )	Femur T Score	Comment
Lee [58]	96	F	13.8 ± 11.1	1.162 ± 0.160	−0.04 ± 1.43	0.959 ± 0.154	−1.12 ± 1.28	No association with BMD and T score
	1320	M	46.8 ± 10.5	1.200 ± 0.165	0.25 ± 1.38	0.984 ± 0.176	0.46 ± 1.13	No association with BMD and T score
Ding [59]	47	F	74 (67–81)	0.946 ± 0.157	NA	0.746 ± 0.112 #	NA	Reduced BMD at femoral neck compared to euthyroid
Rosario [60]	90	F	74 (65–82)	0.91 (0.80–1.21)	NA	1.05 (0.69–1.38) #	NA	No association with BMD and T score
Saler [61]	86	F	33.2 ± 9.5	0.920 ± 0.160	−0.63 ± 1.11	0.830 ± 0.140	−0.76 ± 1.11	No association with BMD and T score
	50	F	73.7 ± 6.8	1.040 ± 0.290	NA	0.700 ± 0.120	NA	No association with BMD and T score
Garin [57]	32	M	73.8 ± 6.6	1.140 ± 0.240	NA	0.950 ± 0.160	NA	No association with BMD and T score
	38	F	56.0 ± 4.4	0.520 ± 0.200	−0.72 ± 1.2	NA	NA	Reduced BMD and T score compared to euthyroid
Rosario [54]	48	F pre	52.9 (35–63)	1.17 (0.92–1.43)	NA	1.04 (0.80–1.29) #	NA	Reduced BMD at femoral neck compared to healthy control
		F post		0.97 (0.60–1.32)	NA	0.89 (0.69–1.13) #	NA	
Lee [55]	19	F	54.3 ± 7.1	0.890 ± 0.160	NA	0.750 ± 0.080 #	NA	Reduced BMD at femoral neck compared to euthyroid
	30	F pre	40.9 ± 7.3	NA	−0.14 ± 0.70 ¶	NA	−0.51 ± 0.81 ¶	Reduced Z score compared to euthyroid
Tauchmanova [51]	30	F post	57.7 ± 6.8	NA	−0.37 ± 0.93 ¶	NA	0.05 ± 0.61 ¶	Reduced Z score compared to euthyroid
	8	F	33.0 ± 5.0	NA	NA	0.921 ± 0.030 #	NA	No association with BMD compared to euthyroid
Gurlek [48]	15	F	28.6 ± 5.8	1.020 ± 0.100	NA	0.790 ± 0.080 #	NA	No association with BMD compared to euthyroid
Foldes [64]	13	F pre	NA	1.010 ± 0.170	NA	0.870 ± 0.150 #	NA	No association with BMD compared to euthyroid
	24	F post	NA	0.840 ± 0.150	NA	0.730 ± 0.120 #	NA	No association with BMD compared to euthyroid

Abbreviation: F: female; M: male; F pre: female premenopausal; F post: female postmenopausal; BMD: bone mineral density; NA: not applicable; # femur neck; Z score.

In the review paper on the skeletal consequences of thyrotoxicosis by Nicholls et al. [18], the authors concluded that endogenous subclinical hyperthyroidism may be associated with increased biochemical markers of bone turnover and a small reduction in BMD. An increased risk of fracture may be present, although these alterations are particularly found in postmenopausal but not in premenopausal women [18]. Two more recent meta-analysis studies on subclinical thyroid dysfunction and fracture risk gave additional support to the clinical effects of subclinical hypothyroidism on bone metabolism [65,66]. This clinical condition was associated with an increased risk of hip and other fractures, particularly in patients with TSH <0.10 mIU/L, but not with TSH between 0.1 and 0.4 mIU/L [67] and older than 65 years, when osteoporosis per se has a high prevalence. On the other hand, a recent study did not find any association of subclinical thyroid disease with bone turnover markers or hip fractures in older men [68], whereas a greater osteoporosis risk due to accelerated hip bone loss in subclinical hyperthyroidism has been reported in a systematic literature search in MEDLINE/EMBASE (1946–016) [69].

Despite some negative results, the available data, taken together support the concept that subclinical hypothyroidism may affect bone metabolism and may be associated with an increased risk for hip and other fractures. The effect of treatment of subclinical thyroid dysfunction on BMD gave conflicting results [70], but large randomized controlled trials are needed. Recent guidelines recommend treatment of subclinical hyperthyroidism in all patients older than 65 years with TSH persistently 0.1 mIU/L or lower and in those with symptoms or comorbidities [71]. The treatment in older subjects is aimed to reduce the incidence of cardiovascular disease. However, TSH measurement should be repeated at least after 4 weeks in order to discriminate transient (in particular transient thyroiditis) and persistent subclinical hyperthyroidism and avoid overtreatment. If a suppressed TSH is confirmed, the possible increased risk of fractures associated with subclinical hyperthyroidism might benefit from this treatment, although there are no randomized studies on this topic.

## 8. Exogenous Subclinical Hyperthyroidism

The main causes of exogenous subclinical hyperthyroidism are inappropriate replacement dose of thyroxine, TSH suppressive thyroxine doses in the therapy of benign thyroid nodules, and thyroid carcinoma. Differentiated thyroid carcinoma (DTC)—papillary and follicle carcinoma—is the most frequent endocrine malignancy [72]. After an initial total thyroidectomy followed by radioactive iodine ablation of thyroid remnant and/or metastatic lesion, patients with DTC are treated with doses of L-T4 sufficient to suppress the circulating TSH concentration because it inhibits further growth of any residual neoplastic tissue and prevents progression of thyroid cancer and relapse [73]. Despite increased relapse-free survival, long-term TSH suppression represents a state of chronic subclinical hyperthyroidism and may be associated with adverse side effects on bone metabolism, cardiac function, and an increased risk of atrial fibrillation, particularly in the elderly.

Early studies reported a negative effect on bone metabolism with a reduced bone mass in patients on prolonged L-T4 treatment with reduced serum TSH [74–76], as reported in Table 3. However, additional studies failed to confirm these data [77–79]. No effect of prolonged L-T4 treatment on bone mass was observed in premenopausal women with reduced serum TSH in a meta-analysis report, whereas postmenopausal women with subclinical hyperthyroidism due to TSH-suppressive doses of L-T4 had reduced bone mass [80]. In postmenopausal women on L-T4 therapy, bone turnover is related to serum TSH concentration the reduction of L-T4 dose is beneficial to BMD and bone turnover [81], whereas L-T4 doses able to suppress TSH without inducing subclinical hyperthyroidism did not induce adverse effects on BMD [82]. Lack of effect on BMD of long-term therapy with L-T4 TSH suppressive for DTC was also reported both in premenopausal and postmenopausal women [83]. Similar data were reported by Belaya et al. [53]. Studies in young patients with exogenous subclinical hyperthyroidism have been reported to have no effect on peak bone mass [84]. A recent cross-sectional study performed in patients treated with a supraphysiological dose of L-T4, initiated during childhood or adolescence for treatment of DTC during the attainment of peak bone mass also suggests that L-T4-mediated subclinical hyperthyroidism does not have significant negative effects on BMD and microarchitecture [85]. Overall,

the available evidences suggest that premenopausal women on chronic TSH suppressive treatment with L-T4 do not have adverse effect on BMD. On the contrary, postmenopausal women on TSH suppressive doses of L-T4 are at risk of bone loss, particularly, when osteopenia or osteoporosis are already present. Finally, a very recent meta-analysis study on the influence of TSH suppression on BMD in patients with DTC suggested a possible association between L-T4-mediated TSH suppression and the lower BMD in postmenopausal women, but not in premenopausal women and men [86].

**Table 3.** Studies that analyzed the effect of exogenous subclinical hyperthyroidism on lumbar spine bone mass density and femoral bone mass density.

Study	n	Gender	Age (Years)	Lumbar Spine BMD (g/cm <sup>2</sup> )	Lumbar Spine T Score	Femur BMD (g/cm <sup>2</sup> )	Femur T Score	Comment
Moon [87]	25	F pre	45.8 ± 3.1	NA	0.29 ± 0.99	NA	0.08 ± 0.86 #	No association with T score
	74	F post	61.4 ± 7.6	NA	−0.84 ± 1.22	NA	−0.97 ± 0.91 #	No association with Z score
Mendonça Monteiro de Barros [85]	17	F	27.4 ± 6.4	1.204 ± 0.140	0.15 ± 1.02	0.150	0.50 ± 1.15 †	No association with BMD and Z score
				NA	NA	1.058 ± 0.170 #	0.24 ± 1.13 #	No association with BMD and Z score
Eftekhari [88]	22	F pre	NA	1.080 ± 0.180	NA	NA	NA	No association with BMD
	33	F post	NA	0.980 ± 0.210	NA	NA	NA	No association with BMD
	11	M	NA	1.110 ± 0.210	NA	NA	NA	No association with BMD
Reverter [83]	44	F pre	39.0 ± 9.0	1.229 ± 0.167	NA	1.032 ± 0.124 #	NA	No association with BMD, T score and Z score
	44	F post	58.8 ± 9.0	1.094 ± 0.248	NA	0.927 ± 0.124 #	NA	No association with BMD, T score and Z score
Appetechia [82]	40	F pre	38.4 ± 3.8	1.030 ± 0.200	0.06 ± 1.15	1.080 ± 0.090	0.15 ± 0.70	No association with BMD, T score and Z score
	56	F post	49.7 ± 1.5	0.850 ± 0.200	−0.11 ± 0.80	0.900 ± 0.0700	−0.10 ± 0.75	No association with BMD, T score and Z score
	40	F pre	NA	1.080 ± 0.160 *	0.07 ± 0.80 *	0.940 ± 0.130 *	0.13 ± 0.80 *	No association with BMD, T score and Z score
Poomthavorn [84]	56	F post	NA	0.810 ± 0.600 *	−0.10 ± 0.70 *	0.800 ± 0.060 *	−0.09 ± 0.78 *	No association with BMD, T score and Z score
	18	F	22.4 ± 4.4	1.160 ± 0.400	NA	0.980 ± 0.250 #	NA	No association with BMD
Kim [79]	36	F post	53(67–65)	0.997 ± 0.149	NA	NA	NA	No association with BMD
			NA	0.986 ± 0.145 *	NA	NA	NA	No association with BMD
Nuzzo [89]	40	F	41.0 ± 1.6	1.070 ± 1.030	0.52 ± 0.24	0.840 ± 0.030 #	−0.03 ± 0.29 #	No association with BMD and Z score
Lecomte [90]	36	F	NA	0.970 ± 0.175	NA	NA	NA	No association with BMD

Abbreviation: F: female; M: male; F pre: female premenopausal; F post: female postmenopausal; BMD: bone mineral density; NA: not applicable; # femur neck; Z score; \* after follow-up.

### 9. Hypothyroidism

Overt hypothyroidism is defined as increased TSH together with T3 and T4 below the lower limit of the reference range. It impairs bone turnover by reducing both osteoclastic bone resorption and osteoblastic activity. Causes of overt hypothyroidism are reported in Table 4. These changes in bone metabolism would result in an increase in bone mineralization. Data on BMD and fracture risk in hypothyroidism are scanty and inconclusive. At the moment, there are no clear data that demonstrate any relationship between BMD in adults and hypothyroidism.

**Table 4.** Causes of hypothyroidism.

<b>Primary hypothyroidism</b>
Acquired
Hashimoto’s thyroiditis
Postablative thyroiditis due to surgery, 131 therapy, neck irradiation for non-thyroid malignancy
Drug induced (lithium, thionamide, amiodarone, iodide)
Drug-induced thyroid destruction (tyrosine kinase inhibitor).
Cytokine induced (interferon- $\gamma$ , interleukin-2)
Thyroid infiltration (hemochromatosis, amyloidosis, Riedel’s struma, sarcoidosis)
Iodine deficiency (endemic goiter)
Goitrogens in foodstuffs
Congenital (NIS or pendrin mutations, organification disorders, defects in thyroglobulin synthesis or processing)
Thyroid agenesis or dysplasia
TSH receptor defects
<b>Central hypothyroidism</b>
Pituitary or hypothalamic disorders
<b>Dopamine infusion</b>
Chronic severe illness
<b>Resistance to thyroid hormone</b>
<b>Transient hypothyroidism</b>
Post subacute thyroiditis

### 10. Subclinical Hypothyroidism

Subclinical hypothyroidism is by definition a laboratory diagnosis and is defined by an elevated TSH value in the presence of normal T4 and T3 serum concentration. The frequency of this clinical condition varies from 6.5 to 8.5 in different populations and tends to increase in older people [91]. Subclinical hypothyroidism is usually asymptomatic and does not have significant effects [92], apart from alteration in lipid metabolism [93–95]. The most common cause of subclinical hypothyroidism is autoimmune thyroiditis, diagnosed through the presence of circulating autoantibodies directed against antigens of the thyroid gland. No replacement therapy is needed until TSH is >10mIU/L [91]. Data on subclinical hypothyroidism effect on bone metabolism are rather scanty, as shown in Table 5. Reduced femoral BMD has been reported in subclinical hypothyroidism [55], whereas thyroid autoimmunity has been suggested to be a potential marker of higher fracture risk in these patients [96]. On the contrary, no association was found between subclinical hypothyroidism and hip fracture risk or BMD in men and women [57–59,61]. Thus, the available evidences cannot demonstrate the possible bone consequence of subclinical hypothyroidism in adults.

### 11. Treatment of Subclinical Thyroid Disorders

Very few studies analyzed the effect of the treatment of thyroid disorders on bone metabolism (Table 6). In one study, Faber et al. analyzed 28 postmenopausal women before and after 2 years of the treatment of subclinical hyperthyroidism with radioiodine. Authors showed that BMD at spin tended to increase after one year and the effect was stable [80] after 2 years. Similar results were found by other authors [81,97–99] while others failed to demonstrate a beneficial effect of the treatment of subclinical hyperthyroidism. No prospective studies have been done for patients with subclinical hypothyroidism.

**Table 5.** Studies that analyzed the effect of subclinical hypothyroidism on lumbar spine bone mass density and femoral bone mass density.

Study	n	Gender	Age (Years)	Lumbar Spine BMD (g/cm <sup>2</sup> )	Lumbar Spine T Score	Femur BMD (g/cm <sup>2</sup> )	Femur T Score	Comment
Lee [90]	1320	F	46.8–10.5	1.134 ± 0.157	−0.273–1.366	0.925–0.136	−0.404–1.129	Reduced risk of osteoporosis in postmenopausal women
	552	M	49.9–12.0	1.190 ± 0.157	0.086–1.310	1.000–0.146	0.462–1.125	No association with BMD and T score
Ding [91]	100	F	70 (67–75)	0.934 ± 0.161	NA	0.752–0.119 #	NA	No association with BMD and T score
	418	F	73.2 ± 5.6	0.920 ± 0.240	NA	0.630 ± 0.130	NA	No association with BMD and T score
Garin [57]	260	M	75.5 ± 6.1	1.160 ± 0.330	NA	0.760 ± 0.160	NA	No association with BMD and T score
	19	F	54.2 ± 7.0	0.930 ± 0.020	NA	0.740 ± 0.120 #	NA	Reduced BMD at femoral neck compared to euthyroid

Abbreviation: F: female; M: male; F pre: female premenopausal; F post: female postmenopausal; BMD: bone mineral density; NA: not applicable; # femur neck.

**Table 6.** Studies that analyzed the effect of the treatment of subclinical hyperthyroidism on lumbar spine bone mass density and femoral bone mass density.

Study	n	Gender	Age (Years)	Lumbar Spine BMD (g/cm <sup>2</sup> ) PRE-Treatment	Lumbar Spine BMD (g/cm <sup>2</sup> ) POST-Treatment	Femur BMD (g/cm <sup>2</sup> ) PRE-Treatment	Femur BMD (g/cm <sup>2</sup> ) POST-Treatment	Comment
Rosario [97]	36	29 F	NA	NA	+1.6% *	NA	+1.9% *	Increased BMD after treatment
Yonen [99]	20	19 F	36.1 ± 1.4	0.991 ± 0.046	0.998 ± 0.048	0.828 ± 0.038 #	0.826 ± 0.042 #	No benefit
Faber [100]	28	F	60 (52–68)	NA	+1.9% *	NA	NA	Increased BMD after treatment
Arata [98]	14	F	20–35	0.908 ± 0.115	1.103 ± 0.139	0.751 ± 0.075	0.905 ± 0.137	Increased BMD after treatment
	18	F post	NA	1.150 ± 0.210	1.170 ± 0.210	0.870 ± 0.130	0.900 ± 0.150	Increased BMD after treatment
Guo [81]	23	F post	NA	1.060 ± 0.0170	1.050 ± 0.170	0.850 ± 0.140	0.830 ± 0.140	Increased BMD after treatment

Data are presented as mean ± standard deviation or median (interquartile range). Abbreviations: F: female; M: male; F pre: female premenopausal; F post: female postmenopausal; BMD: bone mineral density; NA: not applicable; # femur neck; \* variation from baseline levels.

## 12. Conclusions

Current evidences do not allow to draw clear conclusions because most of the studies are limited by their small sample size. However, it is reasonable to hypothesize that subclinical hyperthyroidism may be associated with increased biochemical markers of bone turnover and a small reduction in BMD, but the effective risk of increased frequency of osteoporosis and frequency of fracture needs to be elucidated. The question whether treatment of subclinical hyperthyroidism might reduce the bone loss—or even increase BMD—remains unsolved due to the lack of clear evidences.

On the contrary, subclinical hypothyroidism seems to be unrelated to the bone metabolism, again for the low number of evidences currently available.

While the treatment of overt thyroid disease is mandatory due to its effect on the cardiovascular system, lipid metabolism, and metabolic consequences, the treatment of subclinical disorders are still under debate [89]. The decision of their treatment is based on different factors and the possible reduction of the BMD cannot be considered the main reason to start appropriate therapy.

**Author Contributions:** Writing—original draft preparation, A.P.D.; Writing—review and editing, A.S. and C.D. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Wu, Q.; Xiao, X.; Xu, Y. Evaluating the Performance of the WHO International Reference Standard for Osteoporosis Diagnosis in Postmenopausal Women of Varied Polygenic Score and Race. *J. Clin. Med.* **2020**, *9*, 499. [[CrossRef](#)] [[PubMed](#)]
2. Wu, Q.; Xu, Y.; Lin, G. Trends and Disparities in Self-Reported and Measured Osteoporosis among US Adults, 2007–2014. *J. Clin. Med.* **2019**, *8*, 2052. [[CrossRef](#)] [[PubMed](#)]
3. Sornay-Rendu, E.; Munoz, F.; Duboeuf, F.; Delmas, P.D. Rate of forearm bone loss is associated with an increased risk of fracture independently of bone mass in postmenopausal women: The OFELY study. *J. Bone Min. Res.* **2005**, *20*, 1929–1935. [[CrossRef](#)] [[PubMed](#)]
4. Richelson, L.S.; Wahner, H.W.; Melton, L.J., 3rd; Riggs, B.L. Relative contributions of aging and estrogen deficiency to postmenopausal bone loss. *N. Engl. J. Med.* **1984**, *311*, 1273–1275. [[CrossRef](#)] [[PubMed](#)]
5. Langdahl, B.L.; Andersen, J.D. Treatment of Osteoporosis: Unmet Needs and Emerging Solutions. *J. Bone Metab.* **2018**, *25*, 133–140. [[CrossRef](#)]
6. Doria, C.; Mosele, G.R.; Solla, F.; Maestretti, G.; Balsano, M.; Scarpa, R.M. Treatment of osteoporosis secondary to hypogonadism in prostate cancer patients: A prospective randomized multicenter international study with denosumab vs. alendronate. *Minerva Urol. Nefrol.* **2017**, *69*, 271–277.
7. Caggiari, G.; Leali, P.T.; Mosele, G.R.; Puddu, L.; Badessi, F.; Doria, C. Safety and effectiveness of teriparatide vs alendronate in postmenopausal osteoporosis: A prospective non randomized clinical study. *Clin. Cases Min. Bone Metab.* **2016**, *13*, 200–203. [[CrossRef](#)]
8. Ciurlia, E.; Leali, P.T.; Doria, C. Use of teriparatide off-label: Our experience and review of literature. *Clin. Cases Min. Bone Metab.* **2017**, *14*, 28–34. [[CrossRef](#)]
9. Leali, P.T.; Solla, F.; Maestretti, G.; Balsano, M.; Doria, C. Safety and efficacy of vertebroplasty in the treatment of osteoporotic vertebral compression fractures: A prospective multicenter international randomized controlled study. *Clin. Cases Min. Bone Metab.* **2016**, *13*, 234–236.
10. Delitala, A.P.; Manzocco, M.; Sinibaldi, F.G.; Fanciulli, G. Thyroid function in elderly people: The role of subclinical thyroid disorders in cognitive function and mood alterations. *Int. J. Clin. Pract.* **2018**, *72*, e13254. [[CrossRef](#)]
11. Delitala, A.P.; Terracciano, A.; Fiorillo, E.; Orru, V.; Schlessinger, D.; Cucca, F. Depressive symptoms, thyroid hormone and autoimmunity in a population-based cohort from Sardinia. *J. Affect. Disord.* **2016**, *191*, 82–87. [[CrossRef](#)] [[PubMed](#)]

12. Delitala, A.P.; Delitala, G.; Sioni, P.; Fanciulli, G. Thyroid hormone analogs for the treatment of dyslipidemia: Past, present, and future. *Curr. Med. Res. Opin.* **2017**, *33*, 1985–1993. [[CrossRef](#)] [[PubMed](#)]
13. Delitala, A.P. Subclinical Hyperthyroidism and the Cardiovascular Disease. *Horm. Metab. Res.* **2017**, *49*, 723–731. [[CrossRef](#)] [[PubMed](#)]
14. Delitala, A.P.; Steri, M.; Fiorillo, E.; Marongiu, M.; Lakatta, E.G.; Schlessinger, D.; Cucca, F. Adipocytokine correlations with thyroid function and autoimmunity in euthyroid sardinians. *Cytokine* **2018**, *111*, 189–193. [[CrossRef](#)]
15. Chiamolera, M.I.; Wondisford, F.E. Minireview: Thyrotropin-releasing hormone and the thyroid hormone feedback mechanism. *Endocrinology* **2009**, *150*, 1091–1096. [[CrossRef](#)]
16. Cheng, S.Y.; Leonard, J.L.; Davis, P.J. Molecular aspects of thyroid hormone actions. *Endocr. Rev.* **2010**, *31*, 139–170. [[CrossRef](#)]
17. Maioli, M.; Contini, G.; Santaniello, S.; Bandiera, P.; Pigliaru, G.; Sanna, R.; Rinaldi, S.; Delitala, A.P.; Montella, A.; Bagella, L.; et al. Amniotic fluid stem cells morph into a cardiovascular lineage: Analysis of a chemically induced cardiac and vascular commitment. *Drug Des. Devel. Ther.* **2013**, *7*, 1063–1073.
18. Nicholls, J.J.; Brassill, M.J.; Williams, G.R.; Bassett, J.H. The skeletal consequences of thyrotoxicosis. *J. Endocrinol.* **2012**, *213*, 209–221. [[CrossRef](#)]
19. Rizzoli, R.; Poser, J.; Burgi, U. Nuclear thyroid hormone receptors in cultured bone cells. *Metabolism* **1986**, *35*, 71–74. [[CrossRef](#)]
20. Abu, E.O.; Bord, S.; Horner, A.; Chatterjee, V.K.; Compston, J.E. The expression of thyroid hormone receptors in human bone. *Bone* **1997**, *21*, 137–142. [[CrossRef](#)]
21. Mundy, G.R.; Shapiro, J.L.; Bandelin, J.G.; Canalis, E.M.; Raisz, L.G. Direct stimulation of bone resorption by thyroid hormones. *J. Clin. Invest.* **1976**, *58*, 529–534. [[CrossRef](#)] [[PubMed](#)]
22. Abe, E.; Marians, R.C.; Yu, W.; Wu, X.B.; Ando, T.; Li, Y.; Iqbal, J.; Eldeiry, L.; Rajendren, G.; Blair, H.C.; et al. TSH is a negative regulator of skeletal remodeling. *Cell* **2003**, *115*, 151–162. [[CrossRef](#)]
23. Hase, H.; Ando, T.; Eldeiry, L.; Brebene, A.; Peng, Y.; Liu, L.; Amano, H.; Davies, T.F.; Sun, L.; Zaidi, M.; et al. TNF $\alpha$  mediates the skeletal effects of thyroid-stimulating hormone. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 12849–12854. [[CrossRef](#)]
24. Bassett, J.H.; Williams, A.J.; Murphy, E.; Boyde, A.; Howell, P.G.; Swinhoe, R.; Archanco, M.; Flamant, F.; Samarut, J.; Costagliola, S.; et al. A lack of thyroid hormones rather than excess thyrotropin causes abnormal skeletal development in hypothyroidism. *Mol. Endocrinol.* **2008**, *22*, 501–512. [[CrossRef](#)] [[PubMed](#)]
25. Martini, G.; Gennari, L.; De Paola, V.; Pilli, T.; Salvadori, S.; Merlotti, D.; Valleggi, F.; Campagna, S.; Franci, B.; Avanzati, A.; et al. The effects of recombinant TSH on bone turnover markers and serum osteoprotegerin and RANKL levels. *Thyroid* **2008**, *18*, 455–460. [[CrossRef](#)] [[PubMed](#)]
26. Giusti, M.; Cecoli, F.; Ghiara, C.; Rubinacci, A.; Villa, I.; Cavallero, D.; Mazzuoli, L.; Mussap, M.; Lanzi, R.; Minuto, F. Recombinant human thyroid stimulating hormone does not acutely change serum osteoprotegerin and soluble receptor activator of nuclear factor-kappaB ligand in patients under evaluation for differentiated thyroid carcinoma. *Hormones* **2007**, *6*, 304–313. [[CrossRef](#)]
27. Hutchings, G.; Moncrieff, L.; Dompe, C.; Janowicz, K.; Sibiak, R.; Bryja, A.; Jankowski, M.; Mozdziak, P.; Bukowska, D.; Antosik, P.; et al. Bone Regeneration, Reconstruction and Use of Osteogenic Cells; from Basic Knowledge, Animal Models to Clinical Trials. *J. Clin. Med.* **2020**, *9*, 139. [[CrossRef](#)]
28. Bianco, P.; Gehron Robey, P. Marrow stromal stem cells. *J. Clin. Invest.* **2000**, *105*, 1663–1668. [[CrossRef](#)]
29. Arai, F.; Miyamoto, T.; Ohneda, O.; Inada, T.; Sudo, T.; Brasel, K.; Miyata, T.; Anderson, D.M.; Suda, T. Commitment and differentiation of osteoclast precursor cells by the sequential expression of c-Fms and receptor activator of nuclear factor kappaB (RANK) receptors. *J. Exp. Med.* **1999**, *190*, 1741–1754. [[CrossRef](#)]
30. Li, J.; Sarosi, I.; Yan, X.Q.; Morony, S.; Capparelli, C.; Tan, H.L.; McCabe, S.; Elliott, R.; Scully, S.; Van, G.; et al. RANK is the intrinsic hematopoietic cell surface receptor that controls osteoclastogenesis and regulation of bone mass and calcium metabolism. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 1566–1571. [[CrossRef](#)]
31. Parfitt, A.M.; Travers, R.; Rauch, F.; Glorieux, F.H. Structural and cellular changes during bone growth in healthy children. *Bone* **2000**, *27*, 487–494. [[CrossRef](#)]
32. Recker, R.; Lappe, J.; Davies, K.M.; Heaney, R. Bone remodeling increases substantially in the years after menopause and remains increased in older osteoporosis patients. *J. Bone Min. Res.* **2004**, *19*, 1628–1633. [[CrossRef](#)] [[PubMed](#)]

33. Waung, J.A.; Bassett, J.H.; Williams, G.R. Thyroid hormone metabolism in skeletal development and adult bone maintenance. *Trends Endocrinol. Metab.* **2012**, *23*, 155–162. [[CrossRef](#)] [[PubMed](#)]
34. Eller-Vainicher, C.; Falchetti, A.; Gennari, L.; Cairoli, E.; Bertoldo, F.; Vescini, F.; Scillitani, A.; Chiodini, I. Diagnosis of endocrine disease: Evaluation of bone fragility in endocrine disorders. *Eur. J. Endocrinol.* **2019**, *180*, R213–R232. [[CrossRef](#)]
35. Delitala, A.P.; Orru, M.; Filigheddu, F.; Pilia, M.G.; Delitala, G.; Ganau, A.; Saba, P.S.; Decandia, F.; Scuteri, A.; Marongiu, M.; et al. Serum free thyroxine levels are positively associated with arterial stiffness in the SardinIA study. *Clin. Endocrinol.* **2015**, *82*, 592–597. [[CrossRef](#)]
36. Harvey, C.B.; O’Shea, P.J.; Scott, A.J.; Robson, H.; Siebler, T.; Shalet, S.M.; Samarut, J.; Chassande, O.; Williams, G.R. Molecular mechanisms of thyroid hormone effects on bone growth and function. *Mol. Genet. Metab.* **2002**, *75*, 17–30. [[CrossRef](#)]
37. Krolner, B.; Jorgensen, J.V.; Nielsen, S.P. Spinal bone mineral content in myxoedema and thyrotoxicosis. Effects of thyroid hormone(s) and antithyroid treatment. *Clin. Endocrinol.* **1983**, *18*, 439–446. [[CrossRef](#)]
38. Langdahl, B.L.; Loft, A.G.; Eriksen, E.F.; Mosekilde, L.; Charles, P. Bone mass, bone turnover, calcium homeostasis, and body composition in surgically and radioiodine-treated former hyperthyroid patients. *Thyroid* **1996**, *6*, 169–175.
39. Vestergaard, P.; Mosekilde, L. Hyperthyroidism, bone mineral, and fracture risk—A meta-analysis. *Thyroid* **2003**, *13*, 585–593. [[CrossRef](#)]
40. Bauer, D.C.; Ettinger, B.; Nevitt, M.C.; Stone, K.L.; Study of Osteoporotic Fractures Research, G. Risk for fracture in women with low serum levels of thyroid-stimulating hormone. *Ann. Intern. Med.* **2001**, *134*, 561–568. [[CrossRef](#)]
41. Jodar, E.; Munoz-Torres, M.; Escobar-Jimenez, F.; Quesada-Charneco, M.; Lund del Castillo, J.D. Bone loss in hyperthyroid patients and in former hyperthyroid patients controlled on medical therapy: Influence of aetiology and menopause. *Clin. Endocrinol.* **1997**, *47*, 279–285. [[CrossRef](#)] [[PubMed](#)]
42. Sahin, S.B.; Ayaz, T.; Sumer, F.; Ilkkilic, K.; Algun, E. Bone mineral density in geriatric patients with toxic nodular goiter. *Aging Clin. Exp. Res.* **2015**, *27*, 221–226. [[CrossRef](#)]
43. Dhanwal, D.K.; Gupta, N. Bone mineral density trends in Indian patients with hyperthyroidism—Effect of antithyroid therapy. *J. Assoc. Physicians India* **2011**, *59*, 561–567. [[PubMed](#)]
44. Amato, G.; Mazziotti, G.; Sorvillo, F.; Piscopo, M.; Lalli, E.; Biondi, B.; Iorio, S.; Molinari, A.; Giustina, A.; Carella, C. High serum osteoprotegerin levels in patients with hyperthyroidism: Effect of medical treatment. *Bone* **2004**, *35*, 785–791. [[CrossRef](#)]
45. Ahmed, L.A.; Schirmer, H.; Berntsen, G.K.; Fonnebo, V.; Joakimsen, R.M. Self-reported diseases and the risk of non-vertebral fractures: The Tromso study. *Osteoporos Int.* **2006**, *17*, 46–53. [[CrossRef](#)]
46. Delitala, A.P.; Pilia, M.G.; Ferreli, L.; Loi, F.; Curreli, N.; Balaci, L.; Schlessinger, D.; Cucca, F. Prevalence of unknown thyroid disorders in a Sardinian cohort. *Eur. J. Endocrinol.* **2014**, *171*, 143–149. [[CrossRef](#)]
47. Greenspan, S.L.; Greenspan, F.S. The effect of thyroid hormone on skeletal integrity. *Ann. Intern. Med.* **1999**, *130*, 750–758. [[CrossRef](#)]
48. Gurlek, A.; Gedik, O. Effect of endogenous subclinical hyperthyroidism on bone metabolism and bone mineral density in premenopausal women. *Thyroid* **1999**, *9*, 539–543. [[CrossRef](#)] [[PubMed](#)]
49. Kumeda, Y.; Inaba, M.; Tahara, H.; Kurioka, Y.; Ishikawa, T.; Morii, H.; Nishizawa, Y. Persistent increase in bone turnover in Graves’ patients with subclinical hyperthyroidism. *J. Clin. Endocrinol. Metab.* **2000**, *85*, 4157–4161. [[PubMed](#)]
50. Kvetny, J. Subclinical hyperthyroidism in patients with nodular goiter represents a hypermetabolic state. *Exp. Clin. Endocrinol. Diabetes* **2005**, *113*, 122–126. [[CrossRef](#)]
51. Tauchmanova, L.; Nuzzo, V.; Del Puente, A.; Fonderico, F.; Esposito-Del Puente, A.; Padulla, S.; Rossi, A.; Bifulco, G.; Lupoli, G.; Lombardi, G. Reduced bone mass detected by bone quantitative ultrasonometry and DEXA in pre- and postmenopausal women with endogenous subclinical hyperthyroidism. *Maturitas* **2004**, *48*, 299–306. [[CrossRef](#)] [[PubMed](#)]
52. Morris, M.S. The association between serum thyroid-stimulating hormone in its reference range and bone status in postmenopausal American women. *Bone* **2007**, *40*, 1128–1134. [[CrossRef](#)] [[PubMed](#)]
53. Belaya, Z.E.; Melnichenko, G.A.; Rozhinskaya, L.Y.; Fadeev, V.V.; Alekseeva, T.M.; Dorofeeva, O.K.; Sasonova, N.I.; Kolesnikova, G.S. Subclinical hyperthyroidism of variable etiology and its influence on bone in postmenopausal women. *Hormones* **2007**, *6*, 62–70. [[PubMed](#)]

54. Rosario, P.W. Bone and heart abnormalities of subclinical hyperthyroidism in women below the age of 65 years. *Arq. Bras. Endocrinol. Metabol.* **2008**, *52*, 1448–1451. [[CrossRef](#)] [[PubMed](#)]
55. Lee, W.Y.; Oh, K.W.; Rhee, E.J.; Jung, C.H.; Kim, S.W.; Yun, E.J.; Tae, H.J.; Baek, K.H.; Kang, M.I.; Choi, M.G.; et al. Relationship between subclinical thyroid dysfunction and femoral neck bone mineral density in women. *Arch. Med. Res.* **2006**, *37*, 511–516. [[CrossRef](#)] [[PubMed](#)]
56. van Rijn, L.E.; Pop, V.J.; Williams, G.R. Low bone mineral density is related to high physiological levels of free thyroxine in peri-menopausal women. *Eur. J. Endocrinol.* **2014**, *170*, 461–468. [[CrossRef](#)] [[PubMed](#)]
57. Garin, M.C.; Arnold, A.M.; Lee, J.S.; Robbins, J.; Cappola, A.R. Subclinical thyroid dysfunction and hip fracture and bone mineral density in older adults: The cardiovascular health study. *J. Clin. Endocrinol. Metab.* **2014**, *99*, 2657–2664. [[CrossRef](#)]
58. Wirth, C.D.; Blum, M.R.; da Costa, B.R.; Baumgartner, C.; Collet, T.H.; Medici, M.; Peeters, R.P.; Aujesky, D.; Bauer, D.C.; Rodondi, N. Subclinical thyroid dysfunction and the risk for fractures: A systematic review and meta-analysis. *Ann. Intern. Med.* **2014**, *161*, 189–199. [[CrossRef](#)]
59. Blum, M.R.; Bauer, D.C.; Collet, T.H.; Fink, H.A.; Cappola, A.R.; da Costa, B.R.; Wirth, C.D.; Peeters, R.P.; Asvold, B.O.; den Elzen, W.P.; et al. Subclinical thyroid dysfunction and fracture risk: A meta-analysis. *JAMA* **2015**, *313*, 2055–2065. [[CrossRef](#)]
60. Rosario, P.W.; Carvalho, M.; Mourao, G.F.; Calsolari, M.R. Comparison of Antithyroglobulin Antibody Concentrations Before and After Ablation with <sup>131</sup>I as a Predictor of Structural Disease in Differentiated Thyroid Carcinoma Patients with Undetectable Basal Thyroglobulin and Negative Neck Ultrasonography. *Thyroid* **2016**, *26*, 525–531. [[CrossRef](#)]
61. Siru, R.; Alfonso, H.; Chubb, S.A.P.; Golledge, J.; Flicker, L.; Yeap, B.B. Subclinical thyroid dysfunction and circulating thyroid hormones are not associated with bone turnover markers or incident hip fracture in older men. *Clin. Endocrinol.* **2018**, *89*, 93–99. [[CrossRef](#)] [[PubMed](#)]
62. Segna, D.; Bauer, D.C.; Feller, M.; Schneider, C.; Fink, H.A.; Aubert, C.E.; Collet, T.H.; da Costa, B.R.; Fischer, K.; Peeters, R.P.; et al. Association between subclinical thyroid dysfunction and change in bone mineral density in prospective cohorts. *J. Intern. Med.* **2018**, *283*, 56–72. [[CrossRef](#)] [[PubMed](#)]
63. Buscemi, S.; Verga, S.; Cottone, S.; Andronico, G.; D’Orio, L.; Mannino, V.; Panzavecchia, D.; Vitale, F.; Cerasola, G. Favorable clinical heart and bone effects of anti-thyroid drug therapy in endogenous subclinical hyperthyroidism. *J. Endocrinol. Investig.* **2007**, *30*, 230–235. [[CrossRef](#)] [[PubMed](#)]
64. Kumar, S.; Nadeem, S.; Stan, M.N.; Coenen, M.; Bahn, R.S. A stimulatory TSH receptor antibody enhances adipogenesis via phosphoinositide 3-kinase activation in orbital preadipocytes from patients with Graves’ ophthalmopathy. *J. Mol. Endocrinol.* **2011**, *46*, 155–163. [[CrossRef](#)]
65. Lim, H.; Devesa, S.S.; Sosa, J.A.; Check, D.; Kitahara, C.M. Trends in Thyroid Cancer Incidence and Mortality in the United States, 1974–2013. *JAMA* **2017**, *317*, 1338–1348. [[CrossRef](#)]
66. Biondi, B.; Cooper, D.S. Benefits of thyrotropin suppression versus the risks of adverse effects in differentiated thyroid cancer. *Thyroid* **2010**, *20*, 135–146. [[CrossRef](#)]
67. Taelman, P.; Kaufman, J.M.; Janssens, X.; Vandecauter, H.; Vermeulen, A. Reduced forearm bone mineral content and biochemical evidence of increased bone turnover in women with euthyroid goitre treated with thyroid hormone. *Clin. Endocrinol.* **1990**, *33*, 107–117. [[CrossRef](#)]
68. Diamond, T.; Nery, L.; Hales, I. A therapeutic dilemma: Suppressing doses of thyroxine significantly reduce bone mineral measurements in both premenopausal and postmenopausal women with thyroid carcinoma. *J. Clin. Endocrinol. Metab.* **1991**, *72*, 1184–1188. [[CrossRef](#)]
69. Stall, G.M.; Harris, S.; Sokoll, L.J.; Dawson-Hughes, B. Accelerated bone loss in hypothyroid patients overtreated with L-thyroxine. *Ann. Intern. Med.* **1990**, *113*, 265–269. [[CrossRef](#)]
70. Ahmann, A.J.; Burman, K.D. The role of T lymphocytes in autoimmune thyroid disease. *Endocrinol. Metab. Clin. N. Am.* **1987**, *16*, 287–326. [[CrossRef](#)]
71. Gam, A.N.; Jensen, G.F.; Hasselstrom, K.; Olsen, M.; Nielsen, K.S. Effect of thyroxine therapy on bone metabolism in substituted hypothyroid patients with normal or suppressed levels of TSH. *J. Endocrinol. Investig.* **1991**, *14*, 451–455. [[CrossRef](#)]
72. Kim, E.H.; Jeon, Y.K.; Pak, K.; Kim, I.J.; Kim, S.J.; Shin, S.; Kim, B.H.; Kim, S.S.; Lee, B.J.; Lee, J.G.; et al. Effects of Thyrotropin Suppression on Bone Health in Menopausal Women with Total Thyroidectomy. *J. Bone Metab.* **2019**, *26*, 31–38. [[CrossRef](#)] [[PubMed](#)]

73. Faber, J.; Galloe, A.M. Changes in bone mass during prolonged subclinical hyperthyroidism due to L-thyroxine treatment: A meta-analysis. *Eur. J. Endocrinol.* **1994**, *130*, 350–356. [[CrossRef](#)] [[PubMed](#)]
74. Appetecchia, M. Effects on bone mineral density by treatment of benign nodular goiter with mildly suppressive doses of L-thyroxine in a cohort women study. *Horm. Res.* **2005**, *64*, 293–298. [[CrossRef](#)] [[PubMed](#)]
75. Reverter, J.L.; Holgado, S.; Alonso, N.; Salinas, I.; Granada, M.L.; Sanmarti, A. Lack of deleterious effect on bone mineral density of long-term thyroxine suppressive therapy for differentiated thyroid carcinoma. *Endocr. Relat. Cancer* **2005**, *12*, 973–981. [[CrossRef](#)] [[PubMed](#)]
76. Poomthavorn, P.; Mahachoklertwattana, P.; Ongphiphadhanakul, B.; Preeyasombat, C.; Rajatanavin, R. Exogenous subclinical hyperthyroidism during adolescence: Effect on peak bone mass. *J. Pediatr. Endocrinol. Metab.* **2005**, *18*, 463–469. [[CrossRef](#)]
77. Mendonca Monteiro de Barros, G.; Madeira, M.; Vieira Neto, L.; de Paula Paranhos Neto, F.; Carvalho Mendonca, L.M.; Correa Barbosa Lima, I.; Corbo, R.; Fleiuss Farias, M.L. Bone mineral density and bone microarchitecture after long-term suppressive levothyroxine treatment of differentiated thyroid carcinoma in young adult patients. *J. Bone Min. Metab.* **2016**, *34*, 417–421. [[CrossRef](#)]
78. Yoon, B.H.; Lee, Y.; Oh, H.J.; Kim, S.H.; Lee, Y.K. Influence of Thyroid-stimulating Hormone Suppression Therapy on Bone Mineral Density in Patients with Differentiated Thyroid Cancer: A Meta-analysis. *J. Bone Metab.* **2019**, *26*, 51–60. [[CrossRef](#)]
79. Biondi, B.; Cappola, A.R.; Cooper, D.S. Subclinical Hypothyroidism: A Review. *JAMA* **2019**, *322*, 153–160. [[CrossRef](#)]
80. Delitala, A.P.; Filigheddu, F.; Orru, M.; AlGhatrif, M.; Steri, M.; Pilia, M.G.; Scuteri, A.; Lobina, M.; Piras, M.G.; Delitala, G.; et al. No evidence of association between subclinical thyroid disorders and common carotid intima medial thickness or atherosclerotic plaque. *Nutr. Metab. Cardiovasc. Dis.* **2015**, *25*, 1104–1110. [[CrossRef](#)]
81. Delitala, A.P.; Fanciulli, G.; Maioli, M.; Delitala, G. Subclinical hypothyroidism, lipid metabolism and cardiovascular disease. *Eur. J. Intern. Med.* **2017**, *38*, 17–24. [[CrossRef](#)] [[PubMed](#)]
82. Delitala, A.P.; Fanciulli, G.; Pes, G.M.; Maioli, M.; Delitala, G. Thyroid Hormones, Metabolic Syndrome and Its Components. *Endocr. Metab. Immune Disord. Drug Targets* **2017**, *17*, 56–62. [[CrossRef](#)] [[PubMed](#)]
83. Delitala, A.P.; Scuteri, A.; Fiorillo, E.; Lakatta, E.G.; Schlessinger, D.; Cucca, F. Role of Adipokines in the Association between Thyroid Hormone and Components of the Metabolic Syndrome. *J. Clin. Med.* **2019**, *8*, 764. [[CrossRef](#)] [[PubMed](#)]
84. Polovina, S.P.; Miljic, D.; Zivojinovic, S.; Milic, N.; Micic, D.; Brkic, V.P. The impact of thyroid autoimmunity (TPOAb) on bone density and fracture risk in postmenopausal women. *Hormones* **2017**, *16*, 54–61. [[PubMed](#)]
85. Faber, J.; Jensen, I.W.; Petersen, L.; Nygaard, B.; Hegedus, L.; Siersbaek-Nielsen, K. Normalization of serum thyrotrophin by means of radioiodine treatment in subclinical hyperthyroidism: Effect on bone loss in postmenopausal women. *Clin. Endocrinol.* **1998**, *48*, 285–290. [[CrossRef](#)] [[PubMed](#)]
86. Rosario, P.W. Radioiodine therapy in elderly patients with subclinical hyperthyroidism due to non-voluminous nodular goiter and its effect on bone metabolism. *Arq. Bras. Endocrinol. Metabol.* **2013**, *57*, 144–147. [[CrossRef](#)]
87. Arata, N.; Momotani, N.; Maruyama, H.; Saruta, T.; Tsukatani, K.; Kubo, A.; Ikemoto, K.; Ito, K. Bone mineral density after surgical treatment for Graves' disease. *Thyroid* **1997**, *7*, 547–554. [[CrossRef](#)]
88. Yonem, O.; Dokmetas, H.S.; Aslan, S.M.; Erselcan, T. Is antithyroid treatment really relevant for young patients with subclinical hyperthyroidism? *Endocr. J.* **2002**, *49*, 307–314. [[CrossRef](#)]
89. Delitala, A.P.; Scuteri, A.; Maioli, M.; Mangatia, P.; Vilardi, L.; Erre, G.L. Subclinical hypothyroidism and cardiovascular risk factors. *Minerva Med.* **2019**, *110*, 530–545. [[CrossRef](#)]
90. Lee, K.; Lim, S.; Park, H.; Woo, H.Y.; Chang, Y.; Sung, E.; Jung, H.S.; Yun, K.E.; Kim, C.W.; Ryu, S.; et al. Subclinical thyroid dysfunction, bone mineral density, and osteoporosis in a middle-aged Korean population. *Osteoporos. Int.* **2019**, *31*, 547–555. [[CrossRef](#)]
91. Ding, B.; Zhang, Y.; Li, Q.; Hu, Y.; Tao, X.J.; Liu, B.L.; Ma, J.H.; Li, D.M. Low Thyroid Stimulating Hormone Levels Are Associated with Low Bone Mineral Density in Femoral Neck in Elderly Women. *Arch. Med. Res.* **2016**, *47*, 310–314. [[CrossRef](#)]

92. Rosario, P.W.; Carvalho, M.; Calsolari, M.R. Symptoms of thyrotoxicosis, bone metabolism and occult atrial fibrillation in older women with mild endogenous subclinical hyperthyroidism. *Clin. Endocrinol.* **2016**, *85*, 132–136. [[CrossRef](#)]
93. Saler, T.; Ahbab, S.; Saglam, Z.A.; Keskek, S.O.; Kurnaz, S. Endogenous subclinical hyperthyroidism may not lead to bone loss in premenopausal women. *Hippokratia* **2014**, *18*, 240–244. [[PubMed](#)]
94. Ahn, K.H.; Lee, S.H.; Park, H.T.; Kim, T.; Hur, J.Y.; Kim, Y.T.; Kim, S.H. Effect of adiponectin and sex steroid hormones on bone mineral density and bone formation markers in postmenopausal women with subclinical hyperthyroidism. *J. Obstet. Gynaecol. Res.* **2010**, *36*, 370–376. [[CrossRef](#)] [[PubMed](#)]
95. Ugur-Altun, B.; Altun, A.; Arikan, E.; Guldiken, S.; Tugrul, A. Relationships existing between the serum cytokine levels and bone mineral density in women in the premenopausal period affected by Graves' disease with subclinical hyperthyroidism. *Endocr. Res.* **2003**, *29*, 389–398. [[CrossRef](#)] [[PubMed](#)]
96. Foldes, J.; Tarjan, G.; Szathmari, M.; Varga, F.; Krasznai, I.; Horvath, C. Bone mineral density in patients with endogenous subclinical hyperthyroidism: Is this thyroid status a risk factor for osteoporosis? *Clin. Endocrinol.* **1993**, *39*, 521–527. [[CrossRef](#)] [[PubMed](#)]
97. Moon, J.H.; Jung, K.Y.; Kim, K.M.; Choi, S.H.; Lim, S.; Park, Y.J.; Park, D.J.; Jang, H.C. The effect of thyroid stimulating hormone suppressive therapy on bone geometry in the hip area of patients with differentiated thyroid carcinoma. *Bone* **2016**, *83*, 104–110. [[CrossRef](#)]
98. Eftekhari, M.; Asadollahi, A.; Beiki, D.; Izadyar, S.; Gholamrezanezhad, A.; Assadi, M.; Fard-Esfahani, A.; Fallahi, B.; Takavar, A.; Saghari, M. The long term effect of levothyroxine on bone mineral density in patients with well differentiated thyroid carcinoma after treatment. *Hell. J. Nucl. Med.* **2008**, *11*, 160–163.
99. Nuzzo, V.; Lupoli, G.; Esposito Del Puente, A.; Rampone, E.; Carpinelli, A.; Del Puente, A.E.; Oriente, P. Bone mineral density in premenopausal women receiving levothyroxine suppressive therapy. *Gynecol. Endocrinol.* **1998**, *12*, 333–337. [[CrossRef](#)]
100. Lecomte, P.; Lecureuil, N.; Osorio-Salazar, C.; Lecureuil, M.; Valat, C. Effects of suppressive doses of levothyroxine treatment on sex-hormone-binding globulin and bone metabolism. *Thyroid* **1995**, *5*, 19–23. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).