

Genetic monitoring of brackish-water populations: the Mediterranean toothcarp *Aphanius fasciatus* (Cyprinodontidae) as a model

Ferruccio Maltagliati*

Dipartimento di Zoologia e Antropologia Biologica, Università di Sassari, Corso Margherita di Savoia 15, 07100 Sassari, Italy

ABSTRACT: The measurement of genetic variability and assessment of population genetic losses are important components of environmental management programs. Twenty-three natural populations of the Mediterranean brackish-water toothcarp *Aphanius fasciatus* were investigated using different statistical approaches based on genetic data at 13 polymorphic allozyme loci. In general, no differences between values of within-population genetic variability estimates occurred. The Wilcoxon sign-rank test for heterozygosity excess due to a recent bottleneck was conducted on the array of populations. In addition, a qualitative descriptor of allele frequency distribution was used to infer bottlenecks. Only populations from the Orbetello lagoon and La Salina at Elba Island revealed significant heterozygosity excess under both the infinite allele model (IAM) and stepwise mutation model (SMM). A recent dystrophic crisis may account for the genetic loss detected in the population of *A. fasciatus* from the Orbetello lagoon, whereas exceptionally high predation pressure and/or the increased mortality rate created by local hydrological conditions could be responsible for the bottleneck in the population at La Salina. Tests for bottlenecks have proved effective tools for genetic monitoring of *A. fasciatus* populations.

KEY WORDS: *Aphanius fasciatus* · Allozymes · Heterozygosity · Bottleneck · Genetic monitoring · Conservation

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

The loss of genetic variation within natural populations may occur through bottlenecks, namely severe reductions of population size over a relatively short period. Bottlenecks may determine reductions of within-population genetic diversity owing to the loss of alleles through genetic drift or random fluctuations in allele frequencies (Spellerberg 1996, Storfer 1996, Meffe et al. 1997). However, in some cases severe reductions of population size have not resulted in a substantial loss of genetic variability (Pimm et al. 1989, Queney et al. 2000). Furthermore, both theoretical and empirical evi-

dence suggest that sometimes bottlenecks can cause a short-term increase of genetic variation by converting epistatic variation into additive genetic variation (Goodnight 1988, Bryant & Meffert 1990), but at present no evidence exists about the long-term benefits of bottlenecks.

Aphanius fasciatus Nardo, 1827 is a Mediterranean endemic cyprinodont fish, occurring in coastal brackish-water habitats where it is usually found in large populations. Its distribution is discontinuous, owing to the natural fragmentation of its habitats. Thus, the overall genetic diversity of the species is almost completely determined by the among-population rather than within-population genetic variability (Maltagliati 1998a,b, 1999). Meffe & Vrijenhoek (1988) suggested that species with natural patterns of variation that result in low local population genetic variability but high among-populations variability are worth preserv-

*Present address: Dipartimento di Scienze dell'Uomo e dell'Ambiente, Università di Pisa, Via A. Volta 6, 56126 Pisa, Italy. E-mail: maltagli@discat.unipi.it

ing. In addition, they stressed that a risk to which species genetically structured in this way are subjected is the inbreeding depression caused by a reduction of the effective population size. Within this context, declines in genetic variation in *A. fasciatus* should tend to be local, circumscribed to single disjunct habitats, considering that the species has been demonstrated to be strongly genetically structured with reduced gene flow among populations (Maltagliati 1998a, 1999). Some populations of *A. fasciatus* have declined dramatically, in a few cases even to extinction, due to problems such as brackish-water habitat degradation, pollution of continental and coastal waters, destruction and reduction of salt-works, or introduction of exotic fishes (Bianco 1995). Since knowledge of historical variation in population size is not available for *A. fasciatus* populations, surveys of genetic variation can be employed to determine whether currently large populations have experienced a bottleneck in the past.

The aim of the present study was to assess the possible occurrence of genetic loss in *Aphanius fasciatus* populations using recently introduced methods in order to make an attempt to define the causative factors which have determined genetic loss. Two main approaches were used: (1) differences among heterozygosity values of 23 populations were verified; and (2) the possibility that low levels of within-population genetic variability are due to genetic loss resulting from recent bottlenecks was assessed using the methods developed by Cornuet & Luikart (1996) and Luikart et al. (1998a). These methods rely on the determination of multilocus genotypes at a single point in time, based on the expectation that a bottlenecked population will: (1) temporarily demonstrate an excess of heterozygosity over that expected under the mutation-drift equilibrium (Cornuet & Luikart 1996); and (2) cause a characteristic mode-shift distortion in the distribution of allele frequencies at selectively neutral loci (Luikart et al. 1998a).

MATERIALS AND METHODS

Data on genetic variability of *Aphanius fasciatus* at 13 polymorphic allozyme loci were extracted from a previous study on 11 populations from central Mediterranean brackish-water habitats (Maltagliati 1999). Furthermore, 12 populations were newly analysed using the same genetic markers. A total of 23 populations were analysed (Fig. 1). New samples were collected in Sardinian, Corsican and Tuscan coastal brackish-water habitats in the period between June and October 2000 (Fig. 1). In all brackish-water basins where sampling was carried out, the large population sizes allowed the collection of at least 60 individuals per locality. Details

of sample collections and electrophoretic procedures adopted are outlined in previous papers (Maltagliati 1998b, 1999). The mean number of individuals analysed per locus (\pm SE) ranged from 47.8 ± 3.3 to 66.8 ± 3.9 individuals (Table 1). Loci analysed were *FBP** (fructose biphosphatase, E.C. 3.1.3.11), *GDH-2** (glucose dehydrogenase, E.C. 1.1.1.47), *GPD-1** and *GPD-2** (glycerole-3-phosphate dehydrogenase, E.C. 1.1.1.8), *G6PDH-1** and *G6PDH-2** (glucose-6-phosphate dehydrogenase, E.C. 1.1.1.49), *GPI-1** and *GPI-2** (glucose-6-phosphate isomerase, E.C. 5.3.1.9), *LDH-3** (L-lactate dehydrogenase, E.C. 1.1.1.27), *ME-2** and *ME-3** (malic enzyme E.C. 1.1.1.40), *MPI** (mannose-6-phosphate isomerase, E.C. 5.3.1.8) and *PGM** (phosphoglucomutase, E.C. 5.4.2.2).

The heterozygosity based on Hardy-Weinberg expectations (H_e) was used to estimate within-sample genetic variability. Levels of genetic variability can be compared using this parameter provided genotypic proportions are in Hardy-Weinberg equilibrium, otherwise the observed heterozygosity (H_o = proportion of individuals sampled that are actually heterozygous) should be used. Differences in levels of expected heterozygosity were tested for significance between all possible pairs of samples by paired *t*-tests of arcsine square-root transformed values of single locus H_e (Archie 1985). Two-tailed *t*-tests were used because there was no expected direction of differences in H_e values. Multiple tests were adjusted with the sequential Bonferroni correction with an initial $\alpha = 0.05$ to correct for Type I error (Hochberg 1988).

Tests for recent bottlenecks were performed using the program BOTTLENECK (Piry et al. 1999). A




Fig 1. Localities where *Aphanius fasciatus* was sampled. (●) Samples analysed in a previous study (Maltagliati 1999); (○) newly gathered samples. Locality names and numbers are reported in Table 1, Column 1

2-tailed Wilcoxon sign-rank test for heterozygosity excess (Luikart et al. 1998b) was applied to detect recent bottlenecks. This test relies on the assumption of allele neutrality and mutation-drift equilibrium; it is based on the principle that near the mutation-drift equilibrium the expected heterozygosity (H_{eq}) equals the measured Hardy-Weinberg equilibrium heterozygosity (H_e) in non-bottlenecked populations. Instead, if a population has been subjected to a recent bottleneck, the mutation-drift equilibrium is temporarily disrupted and H_e will be significantly greater than H_{eq} calculated from the number of alleles sampled (Nei et al. 1975, Cornuet & Luikart 1996, Luikart & Cornuet 1998). The Wilcoxon sign-rank test allows the detection of the heterozygosity excess due to the faster loss of low frequency alleles and thus determines whether a population has been recently bottlenecked. In fact, the loss of low-frequency alleles does not account for an equal loss in heterozygosity (Nei et al. 1975, Luikart & Cornuet 1998). However, in non-bottlenecked populations about 50% of the loci are expected to have a slight heterozygote excess and 50% slight heterozygote deficiency, due to genetic drift and/or sampling error. The Wilcoxon sign-rank test for heterozygosity excess allows the reliable detection of small bottlenecks occurring in natural pop-

ulations when at least 5 polymorphic loci and 30 individuals are assayed (Luikart & Cornuet 1998). The calculation of H_{eq} depends on the model of mutation considered to analyse the loci (Cornuet & Luikart 1996). In the present study, H_{eq} values were calculated for the stepwise mutation model (SMM, Kimura & Ohta 1978) and the infinite allele model (IAM, Kimura & Crow 1964), which are considered the 2 extreme models of mutation (Chakraborty & Jin 1992). Classically, allozyme data are assumed to fit the IAM (Chakraborty et al. 1980), but most loci probably evolve according to a model intermediate between SMM and IAM (Di Rienzo et al. 1994, Luikart & Cornuet 1998).

In addition, a qualitative descriptor of allele frequency distribution was used to infer bottlenecks. This test relies on the fact that population bottlenecks cause a characteristic mode-shift distortion in the distribution of allele frequencies at selectively neutral loci. Bottlenecks cause alleles at low-frequency class (<0.1) to temporarily become less abundant than alleles in 1 or more intermediate allele frequency classes (Luikart et al. 1998a). The application of this qualitative method to the present data is appropriate because sample size largely exceeds the minimum requested of 30 specimens (Luikart et al. 1998a).

Table 1. Wilcoxon sign-rank tests for heterozygosity excess (Luikart et al. 1998b) in 23 populations of *Aphanius fasciatus*. L: number of polymorphic loci; N: mean number of individuals sampled per locus; H_e : Hardy-Weinberg expected heterozygosity; IAM: infinite alleles model; SMM: stepwise mutation model; LH_{exc} : number of loci with heterozygosity excess; p: probability of no significant heterozygosity excess. Significant values of probability are in bold

Population	L	N (\pm SE)	H_e (\pm SE)	Wilcoxon sign-rank test					
				IAM			SMM		
				obs LH_{exc}	exp LH_{exc}	p	obs LH_{exc}	exp LH_{exc}	p
1 Biguglia pond	9	57.3 \pm 0.9	0.178 \pm 0.050	5	3.61	0.180	4	4.14	0.410
2 Porto Vecchio salt-works	9	58.6 \pm 0.9	0.186 \pm 0.052	6	3.55	0.125	4	4.05	0.248
3 Santa Giulia pond	9	56.2 \pm 1.5	0.212 \pm 0.062	6	3.61	0.064	5	4.09	0.102
4 Balistra pond	9	55.0 \pm 1.9	0.159 \pm 0.055	3	3.63	0.455	3	4.11	0.455
5 Raza 'e Juncu pond	11	50.5 \pm 2.1	0.180 \pm 0.058	5	4.44	0.319	4	5.02	0.449
6 Porto Taverna pond	10	54.3 \pm 2.9	0.181 \pm 0.051	6	4.13	0.278	4	4.69	0.688
7 Cala Gilgolu pond	9	47.8 \pm 3.3	0.200 \pm 0.054	6	3.71	0.082	6	4.24	0.248
8 Stagno Longu	11	55.4 \pm 1.8	0.202 \pm 0.041	7	4.40	0.103	6	5.02	0.416
9 Muravera pond	7	56.1 \pm 1.5	0.148 \pm 0.051	4	2.81	0.188	3	3.16	0.344
10 Quartu pond	7	58.2 \pm 0.9	0.150 \pm 0.049	5	2.89	0.148	4	3.32	0.289
11 Mistras pond	9	51.2 \pm 2.5	0.116 \pm 0.042	3	3.61	0.633	3	4.13	0.898
12 Casaraccio pond	5	60.9 \pm 0.8	0.077 \pm 0.034	2	1.96	0.406	2	2.26	0.891
13 Pilo pond	7	50.2 \pm 3.1	0.117 \pm 0.043	3	2.84	0.344	3	3.20	0.656
14 La Salina/Elba Island	6	59.4 \pm 1.4	0.157 \pm 0.061	4	2.36	0.039	4	2.72	0.039
15 Perelli canal	6	62.4 \pm 0.9	0.135 \pm 0.051	4	2.36	0.078	3	2.72	0.219
16 Diaccia marsh	6	59.2 \pm 3.4	0.107 \pm 0.049	3	2.35	0.422	2	2.71	0.500
17 Orbetello lagoon	5	57.5 \pm 2.0	0.137 \pm 0.057	4	1.96	0.031	3	2.28	0.047
18 Sabaudia lagoon	8	64.8 \pm 0.5	0.174 \pm 0.059	5	3.30	0.098	5	3.74	0.273
19 Marsala lagoon	11	66.8 \pm 3.9	0.135 \pm 0.043	4	4.66	0.793	2	5.42	0.997
20 Tunis southern lake	9	58.2 \pm 2.5	0.186 \pm 0.054	5	3.80	0.248	4	4.36	0.590
21 Khniss lagoon	9	61.2 \pm 1.9	0.158 \pm 0.096	4	3.76	0.500	3	4.36	0.820
22 Lesina lagoon	6	55.7 \pm 1.3	0.043 \pm 0.027	1	2.39	0.922	1	2.72	0.992
23 Comacchio lagoon	4	63.4 \pm 1.2	0.023 \pm 0.013	0	1.69	1.000	0	1.95	1.000

RESULTS

The number of polymorphic loci (i.e. polyallelic) was variable among the populations analysed ranging from 4 to 11 (Table 1). Probability values obtained by pairwise *t*-tests on arcsine square-root transformed values of expected heterozygosity gave 26 significant cases out of 253; but they reduced to only 1 after the sequential Bonferroni adjustment (comparison Stagno Longu–Comacchio lagoon, $p < 0.001$). The mean number of individuals assayed per locus (ranging from 47.8 to 66.8) abundantly exceeded the minimum number of 20 to 30 required for the reliability of the Wilcoxon sign-rank test (Luikart & Cornuet 1998). Genotypic

proportions of all loci in all populations were in Hardy-Weinberg equilibrium (data not shown), as required for the reliability of the test, because loci that are not in equilibrium could bias the results of the test (Luikart & Cornuet 1998). Two out of 23 populations showed evidence of recent bottlenecks: Orbetello lagoon and La Salina populations (Table 1). Significant heterozygote excess was detected in these populations under both the IAM and SMM (Table 1). In addition, shifted distributions of allele frequencies were revealed in those populations (Fig. 2). Furthermore, the population from Cala Gilgolu exhibited significant distortion of allele frequency distribution (Fig. 2), but it was not significant by Wilcoxon sign-rank test (Table 1).




Fig 2. Distribution of allele frequencies in each population of *Aphanius fasciatus* analysed. This graphical method shows that a population has been recently bottlenecked if fewer alleles are found in the low-frequency class (0 to 0.1) than in 1 or more intermediate frequency classes (Luikart et al. 1998a). Mode-shift distortions were observed in the populations followed by an asterisk

DISCUSSION

The estimates of within-population genetic variability, in general, were not significantly different, with only 1 exception between 2 geographically distant populations (Stagno Longu vs Comacchio lagoon), nor were correlations with habitat surface area found (data not shown). This implies that within-population genetic diversity may be strictly related to variable local conditions.

Another result that stands out from this work is that 2 of the populations analysed (Orbetello lagoon and La Salina) appear to have been subjected to a recent bottleneck. Reductions of size in brackish-water populations following natural and/or anthropogenic catastrophic events are common phenomena (Lardicci et al. 1997, 2001), which sometimes can reach self-organised criticality. Since the 1960s, periodical dystrophic crises in the Orbetello lagoon have been periodically documented (Lardicci et al. 1997, 2001, Cartei et al. 1998). These crises were due to increasing urban water discharge and aquaculture waste waters, which caused large mortality events with fluctuations of fish population sizes. In particular, the crisis that occurred in the summer of 1993 was of exceptional intensity, with abundant macrophyte blooms causing extensive mortality of *Aphanius fasciatus* and other fish due to the reduction of water oxygen concentration to 0 in most areas of the basin (Lardicci et al. 1997, 2001, Cartei et al. 1998, C. Lardicci pers. comm.). As a consequence the Italian Ministry of the Environment declared the Orbetello lagoon an 'area at high environmental risk'. Results of the present study are consistent with the hypothesis that the extent of the mortality event documented in the Orbetello population of *A. fasciatus* produced the effects of a bottleneck, despite the wide surface area of the biotope and the large size of the extant population of *A. fasciatus*. It is likely that in the years following that crisis the population has been rapidly reconstituted by reproduction of individuals that survived in refugia, such as small freshwater tributaries, or zones of the lagoon more proximate to the sea. The other population that appeared bottlenecked was that at Elba Island, which lives in a small-size biotope strongly influenced by tidal changes. At low tide, large zones of the biotope are emersed and groups of individuals of *A. fasciatus* remain confined within small shallow tide pools. These micro-environments may provide extreme environmental conditions in both the cold and warm seasons, contributing to a high mortality rate in the population at La Salina on Elba Island. Moreover, individuals of *A. fasciatus* trapped in the tide pools are more vulnerable to predation by birds and other natural predators. Thus, high levels of physiological stress and predation pressure related to unusual environmental conditions may account for the

genetic loss detected in this population. Results also suggest that the Cala Gilgolu population needs further monitoring because it exhibits distortion in allele frequencies, a method which is considered less powerful than the Wilcoxon sign-rank test (Piry et al. 1999).

From a conservation perspective, it has been suggested that heterozygosity estimates be used in making decisions about the management of populations and species (Vrijenhoek et al. 1985, Leberg 1992). However, the conclusion that can be drawn from the present study is that the extent of heterozygosity values per se do not appear to be effective indicators for genetic monitoring, at least when the events determining genetic losses are relatively recent. Tests for population bottlenecks can represent useful procedures for large-scale assessment of the effects of natural and/or anthropogenic stress on natural populations. These tests had recent effective applications to assess the effects of a viral disease on a wild rabbit population (Queney et al. 2000) and on supportive breeding in brown trout (Hansen et al. 2000) as well as monitoring natterjack toad population declines (Beebe & Rowe 2001). Results of the present study showed that these methods are also effective in monitoring the effect of natural and anthropogenic stress on brackish-water populations of *Aphanius fasciatus*. Because only 1 sample per locality is required and the number of loci analysed need not be large, these methods are attractive candidates for large-scale application.

Acknowledgements. This manuscript has benefited from the comments of G. Cognetti and J. A. Commito. I also wish to thank A. Pinna and G. Vargiu for their valuable assistance during the laboratory work, and G. Calvisi, A. Castelli, M. Carcupino, B. Cristo, M. Curini Galletti and A. Floris for their help during sampling campaigns. The research was financially supported by the 1999 inter-university grant by the Italian Ministry of University and Scientific and Technological Research on 'Environmental stress and biodiversity in brackish waters' and by the Interreg II project.

LITERATURE CITED

- Archie JW (1985) Statistical analysis of heterozygosity data: independent sample comparisons. *Evolution* 39:623–637
- Beebe T, Rowe G (2001) Application of genetic bottleneck testing to the investigation of amphibian declines: a case study with natterjack toads. *Conserv Biol* 15:266–270
- Bianco PG (1995) Mediterranean endemic freshwater fishes of Italy. *Biol Conserv* 72:159–170
- Bryant EH, Meffert LM (1990) Multivariate phenotypic differentiation among bottleneck lines of the housefly. *Evolution* 44:660–668
- Cartei P, Innamorati M, Melillo C (1998) Omeostasi trofica modulata dal mare ed ipertrofia autoctona lagunare. *Biol Mar Medit* 5:41–46 (in Italian)
- Chakraborty R, Jin L (1992) Heterozygote deficiency, population substructure and their implication in DNA fingerprinting. *Hum Genet* 88:267–272

- Chakraborty R, Fuerst PA, Nei M (1980) Statistical studies on protein polymorphism in natural populations. III. Distribution of allele frequencies and the number of alleles per locus. *Genetics* 94:1039–1063
- Cornuet JM, Luikart G (1996) Description and evaluation of two tests for detecting recent bottlenecks. *Genetics* 144: 2001–2014
- Di Rienzo A, Peterson AC, Garza JC, Valdes AM, Slatkin M, Freimer NB (1994) Mutational processes of simple sequence repeat loci in human populations. *Proc Natl Acad Sci USA* 91:3166–3170
- Goodnight CJ (1988) Epistasis and the effect of founder events on the additive genetic variance. *Evolution* 42: 441–454
- Hansen MM, Nielsen EE, Ruzzante DE, Bouza C, Mensberg KLD (2000) Genetic monitoring of supportive breeding in brown trout (*Salmo trutta* L.), using microsatellite DNA markers. *Can J Fish Aquat Sci* 57:2130–2139
- Hochberg Y (1988) A sharper Bonferroni procedure for multiple tests of significance. *Biometrika* 75:800–802
- Kimura M, Crow JF (1964) The number of alleles that can be maintained in a finite population. *Genetics* 49:725–738
- Kimura M, Ohta T (1978) Stepwise mutation model and distribution of allelic frequencies in a finite population. *Proc Natl Acad Sci USA* 75:2868–2872
- Lardicci C, Rossi F, Castelli A (1997) Analysis of macrozoobenthic community structure after severe dystrophic crises in a Mediterranean coastal lagoon. *Mar Pollut Bull* 34:536–547
- Lardicci C, Como S, Corti S, Rossi F (2001) Recovery of the macrozoobenthic community after severe dystrophic crises in a Mediterranean coastal lagoon (Orbetello, Italy). *Mar Pollut Bull* 42:202–214
- Leberg PL (1992) Effect of population bottlenecks on genetic diversity as measured by allozyme electrophoresis. *Evolution* 46:477–494
- Luikart G, Cornuet JM (1998) Empirical evaluation of a test for identifying recently bottlenecked populations from allele frequency data. *Conserv Biol* 12:228–237
- Luikart G, Allendorf FW, Cornuet JM, Sherwin WB (1998a) Distortion of allele frequency distributions provided a test for recent population bottlenecks. *J Hered* 89:238–247
- Luikart G, Sherwin WB, Steele BM, Allendorf FW (1998b) Usefulness of molecular markers for detecting population bottlenecks via monitoring genetic change. *Mol Ecol* 7: 963–974
- Maltagliati F (1998a) Does the Mediterranean killifish *Aphanius fasciatus* (Teleostei: Cyprinodontidae) fit the one-dimensional stepping-stone model of gene flow? *Environ Biol Fishes* 53:385–392
- Maltagliati F (1998b) A preliminary investigation of allozyme genetic variation and population geographical structure in *Aphanius fasciatus* from Italian brackish-water habitats. *J Fish Biol* 52:1130–1140
- Maltagliati F (1999) Genetic divergence in natural populations of the Mediterranean brackish-water killifish *Aphanius fasciatus*. *Mar Ecol Prog Ser* 179:155–162
- Maruyama T (1970) Rate of decrease of genetic variability in subdivided populations. *Biometrika* 57:299–312
- Meffe GK, Vrijenhoek RC (1988) Conservation genetics in the management of desert fishes. *Conserv Biol* 2:157–169
- Meffe GK, Carroll CR and contributors (1997) Principles of conservation biology. Sinauer Associates, Sunderland, MA
- Nei M, Maruyama T, Chakraborty R (1975) The bottleneck effect and genetic variability in populations. *Evolution* 29: 1–10
- Pimm SL, Gittleman JL, McCracken GF, Gilpin M (1989) Plausible alternatives to bottlenecks to explain reduced genetic diversity. *Trends Ecol Evol* 4:176–178
- Piry SG, Luikart G, Cornuet JM (1999) BOTTLENECK: a computer program for detecting recent reductions in the effective population size using allele frequency data. *J Hered* 90:502–503
- Queney G, Ferrand N, Marchandeu S, Azevedo M, Mougél F, Branco M, Monnerot M (2000) Absence of a genetic bottleneck in wild rabbit (*Oryctolagus cuniculus*) population exposed to a severe viral epizootic. *Mol Ecol* 9: 1253–1264
- Spellerberg IF (1996) Conservation biology. Longman Group, Harlow
- Storfer A (1996) Quantitative genetics: a promising approach for the assessment of genetic variation in endangered species. *Trends Ecol Evol* 11:343–348
- Vrijenhoek RC, Douglas ME, Meffe GK (1985) Conservation genetics of endangered fish populations in Arizona. *Science* 229:400–402

Editorial responsibility: Simon Archer (Contributing Editor), Guildford, United Kingdom

Submitted: June 6, 2001; Accepted: February 5, 2002
Proofs received from author(s): May 10, 2002