

XXXV

Società Chimica
Italiana



Dipartimento di Chimica e
Farmacia
Università di Sassari
Istituto di Chimica
Biomolecolare del CNR di
Sassari

Comitato Scientifico:

Prof. Paolo Scrimin
Prof. Raffaele Riccio
Prof. Roberto Ballini
Prof. Valeria Conte
Prof. Marco D'Ischia
Prof. Gianluca Farinola
Prof. Francesco Sannicolò

Conferenze Plenarie:

Dott. Catia Bastioli, Matrica/Novamont
Dott. Walter Cabri, Indena
Prof. Cinzia Chiappe, Uni Pisa
Prof. Antonella Dalla Cort, Uni Roma1
Prof. Bartolo Gabriele, Uni Calabria
Prof. Cesare Gennari, Uni Milano
Prof. Andrea Mazzanti, Uni Bologna
Dott. Alessandro Mordini, CNR Firenze
Prof. Carmen Nàjera, Uni Alicante
Dott. Vincenzo Palermo, ISOF CNR Bologna
Dott. Stefano Protti, Uni Pavia
Dott. Haymo Ross, EurJOCWiley
Prof. Claudio Trombini, Uni Bologna
Dott. Antonio Zanotti-Gerosa, JohnsonMatthey

Comitato Organizzatore:

Prof. Ugo Azzena
Dott. Massimo Carraro
Dott.ssa Giovanna Delogu
Dott.ssa Lidia De Luca
Dott. Davide Fabbri
Dott. Mauro Marchetti
Dott.ssa Luisa Pisano
Dott. Andrea Porcheddu
Dott.ssa Gloria Rassu
Dott. Pietro Spanu
Dott. Pietro Allegrini

Convegno della Divisione di Chimica Organica

Sassari, 9-13 Settembre 2013
Campus Universitario Via Vienna




Synthesis and Study of Polyhydroxylated Phenol Derivatives with Potential Cosmetic and Phytoiatric Applications

¹Maria Antonietta Dettori, ¹Davide Fabbri, ²Xenia Fois, ¹Roberto Dallochio,
¹Alessandro Dessi, ³Pier Andrea Serra, ²Roberto Pantaleoni, ³Gaia Rocchitta e
¹Giovanna Delogu

¹CNR-Istituto di Chimica Biomolecolare, UOS Sassari, Traversa La Crucca 3, 07100 Sassari. ²CNR-Istituto per lo Studio degli Ecosistemi, Traversa La Crucca 3, 07100 Sassari. ³UNISS-Dipartimento di Medicina Clinica e Sperimentale, V.le S. Pietro 43/b, 07100 Sassari
 a.dettori@icb.cnr.it

Tyrosinase (polyphenol oxidase, E.C. 1.14.18.1) and laccase (phenol oxidase, E.C. 1.10.3.2) are multifunctional copper-containing enzymes, that are keys in melanin biosynthesis, melanisation in animals and browning in plants. Tyrosinase inhibitors can therefore be clinically useful for treatment of some dermatological disorders associated with melanin hyperpigmentation, these inhibitors are also known to be useful in cosmetics as whitening agents (1). The involvement of laccase in cuticle sclerotization or tanning is essential to insect survival (2). In the past few decades, a number of polyphenols tyrosinase inhibitors and laccase inhibitors from both natural and synthetic sources, including polyhydroxylated flavonoids, stilbenes and terpenoids have intensively investigated (3). To our knowledge, however, only few biphenyl inhibitors have been reported to date (4).

Our study is aimed to prepare new monomer and dimer phenol derivatives as potential inhibitors of melanin production starting from natural hydroxylated aromatic units. In the figure are reported some of the new synthesized derivatives (**1**, **2** and **3**) representative of gingerdiones, chalcones, and C-prenylated biphenols classes, respectively.



Docking studies of the new compounds with crystal structure of tyrosinase and laccase, were carried on, respectively. Most of the compounds were prepared according to the guidelines of Sustainable Chemistry and they were tested as inhibitors in tyrosinase and laccase assays for potential cosmetic and phytoiatric applications.

- (1) Chang, T. S. *Int. J. Mol. Sci.* **2009**, *10*, 2440-2475.
- (2) Dittmer, N. T.; Kanost, M. R. *Insect Biochem. Mol. Biol.* **2010**, *40*, 179-188.
- (3) Khan, M. T. H.; Khan, S. B.; Ather, A. *Bioorg. Med. Chem.* **2006**, *14*, 938-943.
- (4) Kai, B. *et al. Bioorg. Med. Chem.* **2010**, *18*, 6708-6714.

This work has been supported by Sardinia Autonomous Region - L.R. 7 August 2007, n. 7