

Correspondence

Cat Got Your Tongue?

TO THE EDITOR—A 69-year-old woman with a history of seizures, which were well controlled with sodium valproate, and depression presented with acute onset of a painful swollen tongue that resulted in dysphagia, dysphonia, and drooling of saliva. She had not begun treatment with any new medications and was not using an angiotensin-converting enzyme inhibitor. There was no history of recent dental problems, and her only other medication was dothiepin.

On examination, her temperature was 37.6°C. There was gross swelling of the floor of the mouth and tongue, with no erythema. There was no exudate from the submandibular gland duct. The epiglottis and larynx were normal when visualized by direct laryngoscopy. Her neck was tender to palpation in the submental and submandibular regions, but there was no lymphadenopathy. There was no rash or stridor, and chest auscultation revealed normal vesicular breath sounds without any wheeze. The patient was treated with nebulized adrenaline, intravenous dexamethasone, and meropenem.

The C-reactive protein level was elevated to 142 mg/L (normal, <6 mg/L), and there was a neutrophilia with a marked left shift. CT of the neck revealed some patchy hypodensity in the inferior half of the tongue, which was consistent with edema/inflammatory change. Artifact from amalgam fillings obscured the superior half of the tongue. The salivary glands had a normal appearance, and there was no retropharyngeal or prevertebral collection. The nasopharynx, oropharynx, and laryngopharynx were patent, with no airway compromise.

Two sets of blood culture samples taken at admission to the hospital grew *Pasteu-*

rella multocida. Subsequent history revealed that the patient had a cat in her home, but she could not recall a recent bite or scratch. Antibiotic treatment was changed to intravenous ampicillin and then oral amoxicillin, and the patient made a good recovery.

P. multocida is a small, gram-negative coccobacillus that colonizes the nasopharynx and gastrointestinal tracts of many animals [1]. Most human *P. multocida* infections are caused by dog and cat bites. These bites may result in various clinical syndromes, including cellulitis, subcutaneous abscesses, osteomyelitis, septic arthritis, pneumonia, meningitis, endocarditis, intra-abdominal infection, and septicemia [1]. With regard to upper respiratory tract infection, *P. multocida* has been associated with sinusitis, otitis media, mastoiditis, tonsillitis, peritonsillar abscess, and epiglottitis [1–3]. This is, to our knowledge, the first reported case of *P. multocida* bacteremia manifesting with glossitis.

The earliest written example of the phrase “cat got your tongue” in the *Oxford English Dictionary* is from 1911; the expression is used when addressing someone who is refusing to speak [4]. Several other theories about the origins of this phrase exist: (1) in the Middle East, liars, as punishment, had their tongues removed and fed to the king’s cats; (2) during the Middle Ages, it was thought that, if you saw a witch, her cat would steal or control your tongue such that you could not report the sighting; and (3) the fear of being whipped by the cat o’ nine tails would paralyze the victim into silence [5].

Acknowledgments

Potential conflicts of interest. All authors: no conflicts.

Peter Kelley and David Gordon

Department of Microbiology and Infectious Diseases, Flinders Medical Centre, Bedford Park, South Australia, Australia

References

- Weber DJ, Wolfson JS, Swartz MN, Hooper DC. *Pasteurella multocida* infections: report of 34 cases and review of the literature. *Medicine* 1984;63:133–54.
- Ramdeen GD, Smith RJ, Smith EA, Baddour LM. *Pasteurella multocida* tonsillitis: case report and review. *Clin Infect Dis* 1995;20:1055–77.
- Wine N, Lim Y, Fierer J. *Pasteurella multocida* epiglottitis. *Arch Otolaryngol Head Neck Surg* 1997;123:759–61.
- Oxford English Dictionary Online. Cat got your tongue. Available at <http://dictionary.oed.com/>. Accessed 22 January 2007.
- Ask Yahoo. Where did the phrase “cat got your tongue” come from? Available at <http://ask.yahoo.com/20061102.html>. Accessed 23 July 2007.

Reprints or correspondence: Dr. David Gordon, Dept. of Microbiology and Infectious Disease, Flinders Medical Centre, Bedford Park, South Australia (d.gordon@flinders.edu.au).

Clinical Infectious Diseases 2008;46:146

© 2008 by the Infectious Diseases Society of America. All rights reserved. 1058-4838/2008/4601-0026\$15.00
DOI: 10.1086/523723

Drug Interactions between Warfarin and Efavirenz or Lopinavir-Ritonavir in Clinical Treatment

TO THE EDITOR—Warfarin undergoes liver metabolism by CYP2C9 [1], which is an isoenzyme of cytochrome P450 whose activity could be affected by antiretrovirals. The effect of coadministration in HIV-positive patients has been reported only for nevirapine, which has been shown to decrease warfarin concentration, probably via CYP2C9 induction [2]; no clinical data are yet available for other antiretrovirals. We describe 2 cases concerning coadministration of warfarin with efavirenz or lopinavir-ritonavir.

Case 1 occurred in a 34-year-old black

woman who was receiving stable antiretroviral therapy with didanosine, lamivudine, and efavirenz and who developed an extensive deep vein thrombosis of the right leg with involvement of inferior vena cava and right iliac veins. The patient underwent placement of a vena cava filter and anticoagulant treatment. After 12 days, the patient was discharged from the hospital with a prescription for warfarin (mean daily dose, 5 mg); she had an international normalized ratio (INR) within the optimal range (2–3) and a platelet count of 432,000 platelets/ μ L. The patient was readmitted to the hospital 22 days after discharge because of macrohematuria associated with an INR of 7, a platelet count of 58,000 platelets/ μ L, and normal renal function, without clinical or other pharmacological factors known to increase warfarin activity. The patient's most recent INR measurement, obtained 3 days before, had shown an INR of 2.7, and the patient denied overdosing. After admission, warfarin treatment was transiently discontinued, vitamin K supplementation was initiated, bleeding rapidly ceased, and the patient's INR normalized. The patient's 12-h efavirenz concentration in plasma was 3125 ng/mL. The patient was finally discharged with a normalized platelet count and a mean daily dose of warfarin of 1.25 mg (one-fourth of the initial dose).

Case 2 occurred in a 66-year-old white man who underwent mitral valve replacement due to *Streptococcus viridans* endocarditis after stopping a previously effective HAART regimen that had included zidovudine, lamivudine, and lopinavir-ritonavir (400 mg/100 mg administered twice per day). The patient was discharged from the hospital with a warfarin daily dose ranging between 3.75 mg and 5 mg to maintain an INR within the optimal range (2–3.5). Two months later, the previous HAART regimen was reintroduced. After several days, a progressive increase in warfarin dosing up to 10 mg daily was needed to maintain an INR within the optimal range. Lopinavir plasma trough con-

centration was 4580 ng/mL. After 2 years, when the patient's treatment regimen was simplified to zidovudine-lamivudine-abacavir plus tenofovir, the warfarin dosage was decreased to 3.75–5 mg daily.

These are, to our knowledge, the first clinical reports on coadministration of warfarin with efavirenz or lopinavir-ritonavir. In both cases, the need for warfarin dose adjustments was probably attributable to the opposite effects of these 2 drugs on CYP2C9.

Efavirenz was reported to be a moderate inhibitor in vitro of such isoenzymes, potentially increasing the levels and/or effects of CYP2C9 substrates [3]. For instance, efavirenz was shown to increase phenytoin concentrations, because this anticonvulsant is known to be primarily metabolized by CYP 2C9 and 2C19 [4]. In our case, in which no explanations other than drug interaction were found for the increase in warfarin activity, the magnitude of interaction was impressive, leading to warfarin overdosing. Although concomitant transient thrombocytopenia (probably attributable to previous use of heparin) had probably facilitated bleeding, a marked increase in INR is, per se, considered to indicate a high risk of hemorrhage [5]. Therefore, clinicians should be aware of such possible occurrences in their patients.

On the other hand, ritonavir is known to be an inducer of CYP2C9 [6]. The effect of lopinavir-ritonavir (400 mg/100 mg twice daily) on the metabolism of S-warfarin (administered in a 10-mg single dose) was studied in a group of 7 healthy volunteers, resulting in a 22% decrease in the warfarin area under the curve [7]. In our patient, lopinavir-ritonavir treatment was confirmed to decrease warfarin activity, but the magnitude of the interaction was higher than expected, requiring a doubling of the warfarin dose. This case highlights that data regarding drug interactions obtained from single-dose studies involving healthy volunteers could not exactly predict the extent of such interactions at the steady-state concentration in patients.

In conclusion, the clinical cases reported here showed clinically significant drug interactions when efavirenz or lopinavir-ritonavir were concomitantly administered with warfarin, suggesting the need for careful clinical monitoring and warfarin dose adjustment.

Acknowledgments

Potential conflicts of interest. All authors: no conflicts.

Stefano Bonora,¹ Massimiliano Lanzafame,² Antonio D'Avolio,¹ Laura Trentini,¹ Emanuela Lattuada,² Ercole Concia,² and Giovanni Di Perri¹

¹Department of Infectious Diseases, University of Torino, and ²Unit of Infectious Diseases, "G.B. Rossi" Hospital, University of Verona, Verona, Italy

References

1. Ufer M. Comparative pharmacokinetics of vitamin K antagonists: warfarin, phenprocoumon and acenocoumarol. *Clin Pharmacokinet* 2005; 44:1227–46.
2. Dionisio D, Mininni S, Bartolozzi D, Esperti F, Vivarelli A, Leoncini F. Need for increased dose of warfarin in HIV patients taking nevirapine. *AIDS* 2001; 15:277–8.
3. Smith PF, DiCenzo R, Morse GD. Clinical pharmacokinetics of non-nucleoside reverse transcriptase inhibitors. *Clin Pharmacokinet* 2001; 40:893–905.
4. Robertson SM, Penzak SR, Lane J, Pau AK, Mican JM. A potentially significant interaction between efavirenz and phenytoin: a case report and review of the literature. *Clin Infect Dis* 2005; 41:e15–8.
5. Dentali F, Ageno W, Crowther M. Treatment of coumarin-associated coagulopathy: a systematic review and proposed treatment algorithms. *J Thromb Haemost* 2006; 4:1853–63.
6. Barry M, Gibbons S, Back D, Mulcahy F. Protease inhibitors in patients with HIV disease: clinically important pharmacokinetic considerations. *Clin Pharmacokinet* 1997; 32: 194–209.
7. Yeh R, Gaver V, Park J, et al. Lopinavir/ritonavir induces CYP2C9 and 2C19 activity as measured by warfarin and omeprazole biomarkers in healthy human volunteers [abstract 4.1]. In: Program and abstracts of the 5th International Workshop on Clinical Pharmacology of HIV Therapy (Rome). The Netherlands: Virology Education, 2004; 10.

Reprints or correspondence: Dr. Stefano Bonora, Clinica delle Malattie Infettive, Ospedale Amedeo di Savoia, Corso Svizzera 164, 10149 Torino, Italy (stefano.bonora@unito.it).

Clinical Infectious Diseases 2008; 46:146–7

© 2008 by the Infectious Diseases Society of America. All rights reserved. 1058-4838/2008/4601-0027\$15.00
DOI: 10.1086/524086

***Mycobacterium avium* Subspecies *paratuberculosis* Bacteremia in Type 1 Diabetes Mellitus: An Infectious Trigger?**

TO THE EDITOR—*Mycobacterium avium* subspecies *paratuberculosis* (MAP) is the established cause of paratuberculosis in ruminants (i.e., Johne disease). The bacterium is shed in the milk of infected cows and survives pasteurization. Recently, an association between MAP and Crohn disease has been suggested, wherein MAP has been found to persist in a cell wall-deficient form, escaping clearance by the host immune system [1, 2].

Type 1 (insulin-dependent) diabetes mellitus (T1DM) reflects the interactions of polygenic traits with ill-defined environmental factors, and it is unknown what initiates and maintains autoimmunity to self-antigens expressed in the pancreatic islets of Langerhans [3]. Consumption of cow's milk early in life has been a recognized risk factor in the development of this disease, and environmental microorganisms are thought to trigger autoimmune responses in genetically susceptible individuals [4]. MAP bacilli have recently been hypothesized to trigger molecular mimicry and killing of pancreatic islet cells by the immune system.

We attempted to test the association of MAP with T1DM in an area of endemicity, such as Sardinia, by testing patients with T1DM for the presence of MAP bacilli in peripheral blood. A total of 96 participants, composed of 46 patients with T1DM and 50 healthy control subjects, were tested for the presence of MAP-specific IS900 signature (figure 1) using total DNA extracted from PBMCs. Informed consent from patients, as well as other necessary clearances, were procured before blood samples were obtained. PCR was performed to detect MAP DNA, as described elsewhere [2]. IS900 fragment identity was confirmed by sequencing (GenBank accession group 1517012) and by BLAST analysis against sequences in the




Figure 1. Representative PCR results showing amplification (or otherwise) of 298-bp genomic fragment corresponding to IS900 in diabetic patients (lanes 1–8). Lanes 9 and 10, Control subjects without type 1 diabetes mellitus. Lanes 11 and 12, *Mycobacterium avium* subspecies *paratuberculosis*-positive and *M. avium* subspecies *paratuberculosis*-negative control subjects, respectively. M, molecular marker (100-bp ladder).

National Center for Biotechnology Information database [5].

A total of 29 (63%) of 46 blood samples from diabetic patients were found to be positive for MAP, whereas only 8 (16%) of the 50 samples from healthy control subjects were positive for MAP. Although a majority of diabetic patients with positive PCR results had a family history of diabetes or other genetic and/or autoimmune disorders, 14 diabetic individuals with positive PCR results did not have any history of diabetes or other autoimmune diseases in their families. Although the differences in the outcome of PCR positivity were statistically validated and found to be significant ($\chi^2 = 10.07$; $P \leq .01$), it is our understanding that MAP-positive control subjects might possibly be representing either the genetic resistance or the asymptomatic forms of variable duration that generally precede the clinical presentation of T1DM.

Genetic evidence suggests that there are specific states of immune dysfunction that promote both T1DM and mycobacterial infection [6, 7]. Deficiency of vitamin D has been implicated as a risk factor for T1DM. Interestingly, vitamin D is also implicated in limiting mycobacterial infection by upregulating expression of an antimicrobial peptide [8].

The island of Sardinia has a high incidence of Crohn disease and other autoimmune diseases, such as T1DM, with a very high prevalence of MAP in Sardinian patients with Crohn disease [2, 3]. Because MAP is present in almost one-half of the sheep herds tested in Sardinia, it is prob-

ably endemically contaminating water, milk, and animal feed [9].

Finding evidence of MAP bacteremia in patients with T1DM is a novel observation that might provide an important foundation in establishing an infectious etiology for T1DM. These results also might possibly have implications for countries that have the greatest livestock populations and high incidence of MAP concurrent with the highest numbers of patients with T1DM. Although it is tempting to suggest that MAP could be a potential cause of autoimmune responses associated with T1DM in Sardinia, we believe that a large-scale study involving patients from different genetic and geographic backgrounds might further validate our findings.

Acknowledgments

Financial support. The Italian Ministero Italiano Università E Ricerca (2005064503_003) and the University of Sassari FAR 2006.

Potential conflicts of interest. All authors: no conflicts.

Leonardo A. Sechi,¹ Daniela Paccagnini,¹ Sara Salza,¹ Adolfo Pacifico,¹ Niyaz Ahmed,² and Stefania Zanetti¹

¹Dipartimento di Scienze Biomediche, Università di Sassari, Sassari, Italy; and ²Pathogen Evolution Group, Centre for DNA Fingerprinting and Diagnostics, Hyderabad, India

References

1. Naser SA, Ghobrial G, Romero C, Valentine JF. Culture of *Mycobacterium avium* subspecies *paratuberculosis* from the blood of patients with Crohn's disease. *Lancet* 2004; 364:1039–44.
2. Sechi LA, Scanu AM, Molicotti P, et al. Detection and isolation of *Mycobacterium avium* subspecies *paratuberculosis* from intestinal mucosal

- biopsies of patients with and without Crohn's disease in Sardinia. *Am J Gastroenterol* **2005**; 100:1529–36.
3. Daneman D. Type 1 diabetes. *Lancet* **2006**; 367: 847–58.
 4. Dow. CT. Paratuberculosis and type I diabetes: is this the trigger? *Med Hypotheses* **2006**; 67: 782–5.
 5. National Center for Biotechnology Information database. <http://www.ncbi.nlm.nih.gov/>. Accessed 20 July 2007.
 6. Kissler S, Stern P, Takahashi K, Hunter K, Peterson LB, Wicker LS. In vivo RNA interference demonstrates a role for Nrp1 in modifying susceptibility to type 1 diabetes. *Nat Genet* **2006**; 38:479–83.
 7. Sechi LA, Gazouli M, Sieswerda LE, et al. Relationship between Crohn's disease, infection with *Mycobacterium avium* subspecies paratuberculosis and SLC11A1 gene polymorphisms in Sardinian patients. *World J Gastroenterol* **2006**; 12:7161–4.
 8. Zasloff M. Fighting infections with vitamin D. *Nat Med* **2006**; 12:388–90.
 9. Pickup RW, Rhodes G, Bull TJ, et al. *Mycobacterium avium* subsp. *paratuberculosis* in lake catchments, in river water abstracted for domestic use, and in effluent from domestic sewage treatment works: diverse opportunities for environmental cycling and human exposure. *Appl Environ Microbiol* **2006**; 72:4067–77.

Reprints or correspondence: Dr. Leonardo A. Sechi, Dipartimento di Scienze Biomediche, Università di Sassari, Viale San Pietro 43b, 07100 Sassari, Italy (sechila@uniss.it).

Clinical Infectious Diseases **2008**; 46:148–9
 © 2008 by the Infectious Diseases Society of America. All rights reserved. 1058-4838/2008/4601-0028\$15.00
 DOI: 10.1086/524084

Untangling the Immunological Implications of Nadir on CD4⁺ Cell Recovery during Suppressive Highly Active Antiretroviral Therapy

TO THE EDITOR—Moore et al. [1] recently described a significant difference in CD4⁺ cell recovery following HIV RNA suppression among patients receiving HAART according to nadir CD4⁺ cell count, with individuals whose nadir CD4⁺ cell counts were <200 cells/ μ L not achieving a protective CD4⁺ cell count.

A lack of CD4⁺ cell rescue despite complete HAART-driven viremia suppression represents an everyday dilemma for HIV/AIDS clinicians [2]. Invariably, CD4⁺ cell count nadir proved to be a major and reliable determinant of suboptimal CD4⁺

cell recovery and clinical outcome [1–8]. However, the mere quantification of CD4⁺ cell count nadir fails to qualitatively estimate the ultimate immunological mechanism(s) hindering CD4⁺ cell rescue. Only by detailing the immunological holes in nadir-driven T cell homeostasis will it be possible to devise the most efficacious treatment approach.

By limiting immune recovery assessment to total CD4⁺ cell count in a large patient cohort, Moore et al. [1], although gaining statistical power, miss an insight into the immunological role of CD4⁺ cell count nadir. On the contrary, we feel that stringently focused studies are needed that address the time course of specific immune pathways by nadir CD4⁺ cell count strata. It would be interesting to know whether the authors have found distinct dynamics in immunophenotype and other immunologic parameters according to CD4⁺ cell count ranges.

Given their major regulatory role in T cell homeostasis, common γ -chain cytokines, including IL-2, IL-7, IL-15, and IL-21 [9], might be ideal markers of the direction of immune recovery. Thus, although Moore et al. [1] recommend HAART initiation at a CD4⁺ cell count >350 cells/ μ L for the most robust immune recovery, we advocate a broad immunologic investigation of regulatory cytokine networks to obtain an indication of pretherapy immune damage and CD4⁺ cell rescue potential, as well as possible clinical relapses.

As an example of the exploitation of γ -chain cytokine kinetics, we would like to share our experience with the IL-7 and IL-7R system [10] in a cohort of 18 antiretroviral-naive, HIV-positive patients with advanced infection (nadir CD4⁺ cell count, <150 cells/ μ L) whom we observed prospectively for the first 12 months of HAART. These patients displayed different virological and immunological responses to HAART: 12 patients had concordant responses (HIV RNA level, \leq 50 copies/mL; CD4⁺ cell count, \geq 200 cells/ μ L), and 6 patients had discordant responses (HIV

RNA level, \leq 50 copies/mL; CD4⁺ cell count, \leq 200 cells/ μ L). Despite having a higher median baseline plasma IL-7 level than that for patients with concordant responses, patients with discordant responses had a tendency toward reduced CD4⁺ cell count and lower IL-7R α availability, which, by indicating free IL-7 mainly resulting from receptor down-modulation rather than compensatory production, also rules out a functional potential on T cell homeostasis. In fact, during HAART, despite a progressive reduction in IL-7, only patients with concordant responses displayed a substantial increase in CD4⁺ cell and IL-7R α expression. These findings, by clearly pointing to opposite IL-7 and IL-7R dynamics, also indicate that a lack of IL-7 and IL-7R regulatory control over CD4⁺ cell homeostasis could be the basis of discordant responses, which cautions against IL-7–based treatment.

In conclusion, it is time to investigate the early integration of classic quantitative determinations of CD4⁺ cell count nadir proposed by Moore et al. [1] with additional qualitative measures of CD4⁺ cell count nadir–associated immune imbalances. The effort should be to highlight the immunological role of CD4⁺ cell count nadir, not only as a quantitative reflection of T cell depletion, but mainly as evidence of more-complex T cell qualitative impairment. The intriguing hypothesis that a lack of IL-7– and IL-7R–mediated regulatory function is behind discordant responses in antiretroviral-naive patients with advanced infection clearly indicates the complexity of the immunopathogenetic mechanisms of CD4⁺ cell immune recovery. The possibility of monitoring major T cell homeostasis regulators, such as γ -chain cytokines, by possibly disclosing upstream breakdown offers the appealing prospect of targeted immune interventions.

Acknowledgments

Financial support. Istituto Superiore di Sanità, Italy “National research program on AIDS” (50D.06; 30D.34; 30D.39), “Ermenegildo Zegna” Foundation, Angelantoni Scientifica Spa, and

Fondo Interno Ricerca Scientifica e Tecnologica (FIRST), University of Milan.

Potential conflicts of interest. All authors: no conflicts.

**Laurenzia Ferraris,¹ Giusi Maria Bellistri,²
Valentina Pegorer,² Camilla Tincati,²
Luca Meroni,¹ Massimo Galli,¹
Antonella d'Arminio Monforte,² Andrea Gori,³
and Giulia Marchetti²**

¹Department of Clinical Sciences, "Luigi Sacco" Hospital, and ²Department of Medicine, Surgery and Dentistry, Clinic of Infectious Diseases, "San Paolo" Hospital, University of Milan, Milan, and ³Division of Infectious Diseases, "San Gerardo" Hospital, Monza, Italy

References

1. Moore RD, Keruly JC. CD4⁺ cell count 6 years after commencement of highly active antiretroviral therapy in persons with sustained virologic suppression. *Clin Infect Dis* **2007**; *44*: 441–6.
2. Grabar S, Le Moing V, Goujard C, et al. Clinical outcome of patients with HIV-1 infection according to immunologic and virologic response after 6 months of highly active antiretroviral therapy. *Ann Intern Med* **2000**; *133*: 401–10.
3. Egger M, May M, Chene G, et al. Prognosis of HIV-1-infected patients starting highly active antiretroviral therapy: a collaborative analysis of prospective studies. *Lancet* **2002**; *360*:119–29.
4. Florence E, Lundgren J, Dreezen C, et al. Factors associated with a reduced CD4 lymphocyte count response to HAART despite full viral suppression in the EuroSIDA study. *HIV Med* **2003**; *4*:255–62.
5. Kaufmann GR, Furrer H, Ledergerber B, et al. Characteristics, determinants, and clinical relevance of CD4 T cell recovery to <500 cells/microL in HIV type 1-infected individuals receiving potent antiretroviral therapy. *Clin Infect Dis* **2005**; *41*:361–72.
6. Anthony KB, Yoder C, Metcalf JA, et al. Incomplete CD4 T cell recovery in HIV-1 infection after 12 months of highly active antiretroviral therapy is associated with ongoing increased CD4 T cell activation and turnover. *J Acquir Immune Defic Syndr* **2003**; *33*: 125–33.
7. Benveniste O, Flahault A, Rollot F, et al. Mechanisms involved in the low-level regeneration of CD4⁺ cells in HIV-1-infected patients receiving highly active antiretroviral therapy who have prolonged undetectable plasma viral loads. *J Infect Dis* **2005**; *191*:1670–9.
8. Marchetti G, Gori A, Casabianca A, et al. Comparative analysis of T-cell turnover and homeostatic parameters in HIV-infected patients with discordant immune-virological responses to HAART. *AIDS* **2006**; *20*:1727–36.
9. Alves NL, Arosa FA, van Lier RA. Common gamma chain cytokines: dissidence in the details. *Immunol Lett* **2007**; *108*:113–20.

10. Fry T, Mackall C. Interleukin-7: from bench to clinic. *Blood* **2002**; *99*:3892–904.

Reprints or correspondence: Dr. Giulia Marchetti, Dept. of Medicine, Surgery and Dentistry, Clinic of Infectious Diseases, "San Paolo" Hospital, University of Milan, via A. di Rudini, 8- 20142 Milan, Italy (giulia.marchetti@unimi.it).

Clinical Infectious Diseases **2008**; *46*:149–50

© 2008 by the Infectious Diseases Society of America. All rights reserved. 1058-4838/2008/4601-0029\$15.00
DOI: 10.1086/524087

Cholera or Choleric?

TO THE EDITOR—The cover of the 1 September 2007 issue of *Clinical Infectious Diseases* shows a 14th-century painting based on an 11th-century Arabic book that was described as an illustration of a treatment of cholera. However, there is no evidence that the disease that we now call cholera, caused by *Vibrio cholerae*, existed in Europe or Arabia at those times. The first evidence of the presence of this disease spreading westward outside the Indian subcontinent arose in the early 19th century [1].

Prior to the emergence of what we now know as cholera, the term "cholera" was sometimes applied to what were then considered to be various disorders traceable to disturbances of bile, including a "choleric" or angry disposition [2]. It appears to be likely that the cover illustration was intended to show treatment of that disorder, because the patient appears to be agitated, with breasts exposed, and seems to be biting a towel. There is no sign of the diarrhea, vomiting, dehydration, or collapse that are associated with what we now know as cholera.

Acknowledgments

Potential conflicts of interest. D.R.N.: no conflicts.

David R. Nalin

References

1. Pollitzer R. Cholera. Geneva: World Health Organization Monograph, **1959**.
2. Barua D, Greenough WB III, eds. Cholera. New York: Plenum, **1992**.

Reprints or correspondence: Dr. David R. Nalin, 100 Lucky Hill Rd., West Chester, PA 19382 (davidnalin@aol.com).

Clinical Infectious Diseases **2008**; *46*:150

© 2008 by the Infectious Diseases Society of America. All rights reserved. 1058-4838/2008/4601-0030\$15.00
DOI: 10.1086/524088

Interstrain Antigenic Variability of Mumps Viruses

TO THE EDITOR—We would like to congratulate Peltola et al. [1] on their excellent review. As the authors emphasize, protection from clinical disease is not perfect, even after 2 doses of mumps component vaccine; both primary and secondary vaccine failure were discussed as potential reasons. Another possibility for lack of protection after mumps vaccination (and wild-type virus infection) is the existence of heterologous mumps viruses that may escape immune protection because of interstrain antigenic variability [2, 3]. Here, we report 3 such laboratory-confirmed cases of symptomatic mumps virus infection in patients with documented pre-existing immunity.

The first case occurred in a 12-year-old girl who had received 2 mumps vaccinations (with Jeryl-Lynn vaccine strain in both July 1990 and April 1997) and presented with a lymphadenitis colli. At that time, antimumps IgG antibodies were determined in a serum specimen (IgG antibody concentration, 1400 IU/mL; IgM antibody negative). Two weeks later, bilateral parotid swelling appeared; a second ELISA performed in the same laboratory revealed a 17-fold increase in antimumps IgG antibody concentration to 24,000 IU/mL (IgM antibody negative).

The second case occurred in a 9-year-old boy who had received 2 documented mumps vaccinations (with Rubini vaccine strain in July 1995 and with Jeryl-Lynn vaccine strain in March 2000) and presented with acute bilateral parotid swelling; his antimumps IgG antibody concentration was 2900 IU/mL (IgM antibody negative). Four weeks later, convalescent-phase serologic examination revealed a

5.5-fold increase in antimumps IgG antibody concentration to 16,000 IU/mL.

The third case occurred in a 30-year-old male pediatric resident with no history of mumps vaccination who had mumps disease during childhood. He had an antimumps IgG antibody concentration of 33 IU/mL in a serum sample obtained for screening for newly employed health care workers at our institution. One year later, the patient developed acute bilateral parotid swelling. Examination of a serum sample obtained at that time revealed an 8.5-fold increase in IgG antibody concentration to 277 IU/mL; this sample was tested together with the previous sample and was antimumps IgM antibody negative. Neutralizing assays confirmed reinfection, despite pre-existing neutralizing antimumps serum antibodies. Ten days later, the patient's 26-year-old female partner (with no history of mumps disease or vaccination) developed a primary mumps infection (IgM antibody positive). These cases illustrate that symptomatic mumps reinfection may occur in the presence of pre-existing specific mumps virus serum antibodies.

Acknowledgments

We thank A. Tischer for providing the neutralizing assays.

Potential conflicts of interest. U.H. and J.B.: no conflicts.

Ulrich Heininger and Jan Bonhoeffer

University Children's Hospital Basel,
Basel, Switzerland

References

1. Peltola H, Kulkarni PS, Kapre SV, Paunio M, Jadhav SS, Dhere RM. Mumps outbreaks in Canada and the United States: time for new thinking on mumps vaccines. *Clin Infect Dis* 2007;45:459–66.
2. Nojd J, Teele T, Samuelsson A, Orvell C. Mumps virus neutralizing antibodies do not protect against reinfection with a heterologous mumps virus genotype. *Vaccine* 2001;19:1727–31.
3. Rubin S, Mauldin J, Chumakov K, Vanderzanden J, Iskow R, Carbone K. Serological and phylogenetic evidence of monotypic immune responses to different mumps virus strains. *Vaccine* 2006;24:2662–8.

Reprints or correspondence: Dr. Ulrich Heininger, Infectious Disease and Vaccines, University Children's Hospital Basel (UKBB), Postbox, 4005 Basel, Switzerland (Ulrich.Heininger@unibas.ch).

Clinical Infectious Diseases 2008;46:150–1

© 2008 by the Infectious Diseases Society of America. All rights reserved. 1058-4838/2008/4601-0031\$15.00
DOI: 10.1086/524089

Prevention of Traveler's Diarrhea: A Call to Reconvene

TO THE EDITOR—More than 40 years have passed since the initial reports of traveler's diarrhea (TD), and much has been learned about the epidemiology of this illness, which is most often attributed to colonization of the intestinal track with pathogenic bacteria of a broad variety [1, 2]. Although our understanding has improved and effective treatment has been made available, individuals are still traveling from developed countries to lesser developed countries, and ~1 of 3 will experience an acute illness; some of these individuals will be incapacitated, but most will improve within a few days [3].

In addition, >50 years ago, the initial descriptions of postinfectious functional bowel disorders were reported [4, 5], and in the past decade, a proliferation of studies summarized in 2 recent systematic reviews have concluded that ~1 of 10 people who develop TD will acquire postinfectious irritable bowel syndrome (PI-IBS), despite normal preexisting bowel habits [6, 7]. Although PI-IBS is perhaps not as debilitating as less frequent sequelae of reactive arthritis [8], Guillain Barré syndrome [9], or inflammatory bowel disease [10], the attributable burden of PI-IBS is measurable. A quick calculation reveals that, among the 100 million travelers each year from industrialized countries who are at intermediate to high risk of diarrhea, ~30 million will develop TD, and 3 million new cases of PI-IBS will arise [11]. PI-IBS has been reported to persist in 57% of patients 6 years after its onset in one study, and postinfectious functional bowel disorders (predominantly PI-IBS) persisted in 76% of patients 5 years after its

onset in another study [12, 13]. These illnesses not only decrease the quality of life of those afflicted, but the economic impact is considerable. In a review of 18 economic studies conducted in the United States and the United Kingdom, the direct cost per patient-year was estimated to be US\$348–\$8750 (in 2002), the annual number of lost days of work was 9–22 days, and the indirect cost per patient-year was estimated to be \$355–\$3344 [14]. Unquestionably, this represents significant social and economic impact.

In addition to travel-related PI-IBS, the fraction of cases of IBS secondary to domestically acquired foodborne illness needs to be considered. IBS is the most common chronic medical illness, seen in up to 15% of the US population and accounting for nearly one-third of all costs in gastroenterology as a result of investigation, management, and work loss among those seeking care [15–17]. Although the attributable fraction of IBS cases caused by domestic foodborne illness is unknown, it is likely to be large, given the frequency of these illnesses [18]. Thus, the traditional thought of foodborne illness as a self-limited condition also needs serious reconsideration and should inform future food security policy decisions.

Finally, shortly after the first reports of TD 40 years ago, reports of studies evaluating agents for use in prophylaxis followed. A number of effective regimens have been described; however, widespread use of prophylaxis, particularly antimicrobial agents, has been discouraged—a position formalized in a 1985 National Institutes of Health consensus meeting that has been unchallenged [19]. We feel that PI-IBS is a development that challenges this consensus.

The balance of chemoprophylaxis risk, compared with the consequences of TD and sequelae, needs reassessment. Although certain questions remain regarding the natural history of PI-IBS, effective treatment, and consequences of chemoprophylaxis, it is time that the questions regarding what can be done to prevent TD

and who could benefit from preventive interventions are reconsidered.

Acknowledgments

Potential conflicts of interest. M.P. and M.S.R.: no conflicts.

Mark Pimentel^{1,2} and Mark S. Riddle³

¹Cedars-Sinai Medical Center, Burns and Allen Research Institute, and ²University of California—Los Angeles, Geffen School of Medicine, Los Angeles, California; and ³Enteric Diseases Department, Naval Medical Research Center, Silver Spring, Maryland

References

1. Hyllner S, Heinlaid H. Diarrhea in tourists and hygiene—observations during a field study of diarrheal symptoms in Scandinavian tourists in the Canary Islands during the years 1960–63 [in Swedish]. *Nordisk Hygienisk Tidsskrift* **1964**;45:1–13.
2. Kean BH. The diarrhea of American travelers abroad. *Minn Med* **1959**;42:1749–51.
3. Traveler's diarrhea: a 50-year perspective. Proceedings of a conference organized by the International Society of Travel Medicine. June 12–14, 2004. Jackson Hole, Wyoming, USA. *Clin Infect Dis* **2005**;41(Suppl 8):S521–86.
4. Chaudhary NA, Truelove SC. The irritable colon syndrome: a study of the clinical features, predisposing causes, and prognosis in 130 cases. *Q J Med* **1962**;31:307–22.
5. Stewart GT. Post-dysenteric colitis. *Br Med J* **1950**;1:405–9.
6. Thabane M, Kottachchi DT, Marshall JK. Systematic review and meta-analysis: the incidence and prognosis of post-infectious irritable bowel syndrome. *Aliment Pharmacol Ther* **2007**;26:535–44.
7. Halvorson HA, Schlett CD, Riddle MS. Postinfectious irritable bowel syndrome—a meta-analysis. *Am J Gastroenterol* **2006**;101:1894–9; quiz 942.
8. Yates JA, Stetz LC. Reiter's syndrome (reactive arthritis) and travelers' diarrhea. *J Travel Med* **2006**;13:54–6.
9. Butzler JP. *Campylobacter*, from obscurity to celebrity. *Clin Microbiol Infect* **2004**;10:868–76.
10. Garcia Rodriguez LA, Ruigomez A, Panes J. Acute gastroenteritis is followed by an increased risk of inflammatory bowel disease. *Gastroenterology* **2006**;130:1588–94.
11. Steffen R, DuPont H, Wilder-Smith A. *Manual of travel medicine and health*. 3rd ed. Hamilton, Ontario: B.C. Decker, **2007**.
12. Neal KR, Barker L, Spiller RC. Prognosis in post-infective irritable bowel syndrome: a six year follow up study. *Gut* **2002**;51:410–3.
13. Tornblom H, Holmval P, Svenungsson B, Lindberg G. Gastrointestinal symptoms after infectious diarrhea: a five-year follow-up in a Swedish cohort of adults. *Clin Gastroenterol Hepatol* **2007**;5:461–4.
14. Maxion-Bergemann S, Thielecke F, Abel F, Bergemann R. Costs of irritable bowel syndrome in the UK and US. *Pharmacoeconomics* **2006**;24:21–37.
15. Cremonini F, Talley NJ. Irritable bowel syndrome: epidemiology, natural history, health care seeking and emerging risk factors. *Gastroenterol Clin North Am* **2005**;34:189–204.
16. Quigley EM, Bytzer P, Jones R, Mearin F. Irritable bowel syndrome: the burden and unmet needs in Europe. *Dig Liver Dis* **2006**;38:717–23.
17. American Gastroenterological Association. *The burden of gastrointestinal diseases*. Bethesda, MD: American Gastroenterological Association, **2001**. Available at: <http://www.gastro.org/user-assets/Documents/burden-report.pdf>. Accessed 20 January 2007.
18. Mead PS, Slutsker L, Dietz V, et al. Food-related illness and death in the United States. *Emerg Infect Dis* **1999**;5:607–25.
19. Travelers' diarrhea. NIH Consensus Development Conference. *JAMA* **1985**;253:2700–4.

Reprints or correspondence: Dr. Mark S. Riddle, Enteric Disease Dept., Naval Medical Research Center, Silver Spring, MD 20817 (markriddlemd@hotmail.com).

Clinical Infectious Diseases **2008**;46:151–2

© 2008 by the Infectious Diseases Society of America. All rights reserved. 1058-4838/2008/4601-0032\$15.00
DOI: 10.1086/524090