


STRUCTURAL MODEL OF A Ni(II) COMPLEX WITH A 30-AMINOACID PEPTIDE THROUGH AN NMR STUDY

M.A. Zoroddu, M. Peana, S. Medici

Department of Chemistry, University of Sassari, Via Vienna 2, 07100, Sassari, ITALY
zoroddu@uniss.it

Multidimensional NMR spectroscopy is a useful tool for the calculation of structures or structural models of metal-peptide complexes in solution.¹ We applied bidimensional NMR techniques to study the interactions of Ni(II) ions with a 30-aminoacid peptide, a fragment of the C-terminal tail of Cap 43 protein. This protein is strictly connected to nickel exposure in cells, since it seems to be specifically expressed as a response to the presence of this metal in the cellular medium^{2,3} and it is also related to cancer development; an abnormal level of Cap43 protein has been detected in a number of tumour tissues.^{2,4} The striking feature of Cap 43 is a three-repeated decapeptide sequence at its C-terminus; each 10-aminoacid fragment (TRSRSHSTSEG) bearing a histidinic residue, which has been indicated as an anchoring site for metal binding in numerous cases. We previously reported that each fragment is able to coordinate a Ni(II) ion in a very effective way.^{5,6}

Structure calculations for the peptide-metal complex were performed for a single monohistidinic fragment on the basis of the ROE cross-correlations observed in the 2D ¹H-¹H ROESY spectra at pH = 10. The metal complex involves an imidazolic nitrogen of histidine residue, three amidic nitrogens of the backbone and an oxygen atom from a deprotonated serine residue which takes part to the formation of a square pyramidal structure.



The structural model calculated (figure below) allowed us a better understanding of the features of nickel coordination with the C-terminal region of Cap43 protein.

References

1. E. Gaggelli, H. Kozłowski, D. Valensin, G. Valensin, *Mol. BioSyst.* 1 (2005) 79-84. M. A. Zoroddu, M. Peana, S. Medici, *Dalton Trans.*, (2007) 379-384
2. D. Zhou, Salnikow K, M. Costa, *Cancer Res*, 58 (1998) 2182-2189
3. K. Salnikow, T. Kluz, M. Costa, *Toxicol. Appl. Pharmacol.* 160 (1999) 127-132
4. K. Salnikow, D. Zhou, T. Kluz, C. Wang, M. Costa, in: A. Sarkar (Ed.), *Metal and Genetics*, Kluwer Academic, Plenum Publishers, New York (1999) 131-144
5. M.A. Zoroddu, T. Kowalik-Jankowska, H. Kozłowski, K. Salnikow, M. Costa, *J. Inorg. Biochem.* 85 (2001) 47-54.
6. M.A. Zoroddu, M. Peana, T. Kowalik-Jankowska, H. Kozłowski, M. Costa, *J. Inorg Biochem.* 98 (2004) 931-939

STRUCTURAL MODEL OF A Ni(II) COMPLEX WITH A 30-AMINOACID PEPTIDE THROUGH AN NMR STUDY

M.A. Zoroddu, M. Peana, S. Medici

Department of Chemistry, University of Sassari, Via Vienna 2, 07100, Sassari, ITALY

zoroddu@uniss.it

Multidimensional NMR spectroscopy is a useful tool for the calculation of structures or structural models of metal-peptide complexes in solution.¹ We applied bidimensional NMR techniques to study the interactions of Ni(II) ions with a 30-aminoacid peptide, a fragment of the C-terminal tail of Cap 43 protein. This protein is strictly connected to nickel exposure in cells, since it seems to be specifically expressed as a response to the presence of this metal in the cellular medium^{2,3} and it is also related to cancer development; an abnormal level of Cap43 protein has been detected in a number of tumour tissues.^{2,4} The striking feature of Cap 43 is a three-repeated decapeptide sequence at its C-terminus; each 10-aminoacid fragment (TRSRSHSTSEG) bearing a histidinic residue, which has been indicated as an anchoring site for metal binding in numerous cases. We previously reported that each fragment is able to coordinate a Ni(II) ion in a very effective way.^{5,6} Structure calculations for the peptide-metal complex were performed for a single monohistidinic fragment on the basis of the ROE crosscorrelations observed in the 2D ¹H-¹H ROESY spectra at pH = 10. The metal complex involves an imidazolic nitrogen of histidine residue, three amidic nitrogens of the backbone and an oxygen atom from a deprotonated serine residue which takes part to the formation of a square pyramidal structure. The structural model calculated (figure below) allowed us a better understanding of the features of nickel coordination with the C-terminal region of Cap43 protein.

References

1. E. Gaggelli, H. Kozlowski, D. Valensin, G. Valensin, *Mol. BioSyst.* 1 (2005) 79- 84. M. A. Zoroddu, M. Peana, S. Medici, *Dalton Trans.*, (2007) 379-384
2. D. Zhou, Salnikow K, M. Costa, *Cancer Res*, 58 (1998) 2182-2189
3. K. Salnikow, T. Kluz, M. Costa, *Toxicol. Appl. Pharmacol.* 160 (1999) 127-132
4. K. Salnikow, D. Zhou, T. Kluz, C. Wang, M. Costa, in: A. Sarkar (Ed.), *Metal and Genetics*, Kluwer Academic, Plenum Publishers, New York (1999) 131–144
5. M.A. Zoroddu, T. Kowalik-Jankowska, H. Kozlowski, K. Salnikow, M. Costa, *J. Inorg. Biochem.* 85 (2001) 47-54.
6. M.A. Zoroddu, M. Peana, T. Kowalik-Jankowska, H. Kozlowski, M. Costa, *J. Inorg Biochem.* 98 (2004) 931-939