

XXXV

Società Chimica
Italiana



Dipartimento di Chimica e
Farmacia
Università di Sassari
Istituto di Chimica
Biomolecolare del CNR di
Sassari

Comitato Scientifico:

Prof. Paolo Scrimin
Prof. Raffaele Riccio
Prof. Roberto Ballini
Prof. Valeria Conte
Prof. Marco D'Ischia
Prof. Gianluca Farinola
Prof. Francesco Sannicolò

Conferenze Plenarie:



Dott. Catia Bastioli, Matrica/Novamont
Dott. Walter Cabri, Indena
Prof. Cinzia Chiappe, Uni Pisa
Prof. Antonella Dalla Cort, Uni Roma1
Prof. Bartolo Gabriele, Uni Calabria
Prof. Cesare Gennari, Uni Milano
Prof. Andrea Mazzanti, Uni Bologna
Dott. Alessandro Mordini, CNR Firenze
Prof. Carmen Nàjera, Uni Alicante
Dott. Vincenzo Palermo, ISOF CNR Bologna
Dott. Stefano Protti, Uni Pavia
Dott. Haymo Ross, EurJOCWiley
Prof. Claudio Trombini, Uni Bologna
Dott. Antonio Zanotti-Gerosa, JohnsonMatthey

Comitato Organizzatore:

Prof. Ugo Azzena
Dott. Massimo Carraro
Dott.ssa Giovanna Delogu
Dott.ssa Lidia De Luca
Dott. Davide Fabbri
Dott. Mauro Marchetti
Dott.ssa Luisa Pisano
Dott. Andrea Porcheddu
Dott.ssa Gloria Rassu
Dott. Pietro Spanu
Dott. Pietro Allegrini

Convegno della Divisione di Chimica Organica

Sassari, 9-13 Settembre 2013
Campus Universitario Via Vienna




Green and Convenient Arene-Catalyzed Reductive Lithiation of Heterocyclic Compounds

Ashenafi Damtew Mamuye, Ugo Azzena, Luisa Pisano, Massimo Carraro, Mario Pittalis

Dipartimento di Chimica e Farmacia, Università di Sassari, via Vienna 2, I-07100-Sassari, Italy

mamuye@uniss.it

In the last decades reductive lithiation established itself as a particularly powerful procedure for the generation of functionalized organometals. In addition, the enormous role of this reaction in the reductive ring opening and functionalization of heterocycles has been reported (Scheme 1) [1].



Scheme 1. Reductive lithiation of different classes of heterocyclic compounds; X = NR, O, S; n = 1,2.

Almost always, the efficiency of this reaction strongly depends on the presence of a polycyclic aromatic hydrocarbon (PAH), employed as an electron shuttle, either in catalytic (CA) or stoichiometric (SA) amounts. Indeed, formation of their soluble and intensively coloured radical anions is considered to speed up the electron-transfer process, thus avoiding formation of by-products [2]. According to its ability to dissolve PAH's radical anions, tetrahydrofuran (THF) emerged as the solvent of choice for these reactions [3]. However, several factors (water solubility, relatively low boiling point, expensive purification, drying and recycling) makes THF a poorly eco-efficient solvent. Herein we report on the employment of 2-methyltetrahydrofuran (2-MeTHF) and cyclopentyl methyl ether (CPME) as alternative green solvents for the generation of PAH's radical anions and their employment in the generation of organolithium reagents.

Accordingly, critical experiments were performed for the generation of lithium radical anions of several PAHs and a diene in both solvents, under both CA and SA protocols. Almost all of the above mentioned hydrocarbons rapidly generate radical anions with different fascinating visible colors at 0 °C. Based on this, influence of different electron shuttles and their efficiency in the reductive lithiation of *N*-phenylaziridine, phthalan will be disclosed. Moreover, we will report an interesting example of chemoselectivity concerning to the reductive lithiation of 4-chlorobenzyl methyl ether in a green solvent.

References:

- [1] Yus, M.; Foubelo, F. *Adv. Heterocycl. Chem.*, **2006**, 91, 135-158, and references therein.
- [2] Azzena, U., Pilo, L., Piras, E. *Tetrahedron*, **2000**, 56, 3775-3780, and references therein.
- [3] Cohen, T., Kreethadumrongdat, T., Liu, X.; Kulkarni, V. *J. Am. Chem. Soc.*, **2001**, 123, 3478-3483.