Replacing organic with mineral N fertilization does not reduce nitrate leaching in double crop forage systems under Mediterranean conditions

Questa è la versione Post print del seguente articolo:

Original

Replacing organic with mineral N fertilization does not reduce nitrate leaching in double crop forage systems under Mediterranean conditions / Demurtas, Ce; Seddaiu, Giovanna; Ledda, Luigi; Cappai, C; Doro, L; Carletti, A; Roggero, Pier Paolo. - In: AGRICULTURE, ECOSYSTEMS & ENVIRONMENT. - ISSN 0167-8809. - 219:(2016), pp. 83-92. [10.1016/j.agee.2015.12.010]

Availability: This version is available at: 11388/59477 since: 2022-05-23T23:52:01Z

Publisher:

Published DOI:10.1016/j.agee.2015.12.010

Terms of use:

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

Publisher copyright

note finali coverpage

(Article begins on next page)

Elsevier Editorial System(tm) for Agriculture, Ecosystems and Environment Manuscript Draft

Manuscript Number: AGEE13437R3

Title: Replacing organic with mineral N fertilization does not reduce nitrate leaching in double crop forage systems under Mediterranean conditions

Article Type: Research Paper

Keywords: Disc lysimeter; Silage maize; Italian ryegrass; Cattle slurry; Manure; Nitrate Vulnerable Zone.

Corresponding Author: Dr. Giovanna Seddaiu, Ph.D.

Corresponding Author's Institution: University of Sassari

First Author: Clara E Demurtas

Order of Authors: Clara E Demurtas; Giovanna Seddaiu, Ph.D.; Luigi Ledda; Chiara Cappai; Luca Doro; Alberto Carletti; Pier Paolo Roggero

Manuscript Region of Origin: ITALY

Abstract: This research evaluated the impact of four nitrogen (N) fertilizer management systems on nitrate losses in an irrigated forage system under Mediterranean conditions within a Nitrate Vulnerable Zone (NVZ). The experiment was conducted from June 2009 to May 2012 in an intensive dairy cattle farm that produces silage maize and Italian ryegrass in a double cropping system. A monthly monitoring of the nitrate concentrations in the soil solution was carried out using 10 cm diameter disc lysimeters. The N fertilization systems had a target N application of 316 and 130 kg ha-1 for maize and Italian ryegrass, respectively. Four systems were compared: cattle manure (MA); cattle slurry (SL); cattle slurry + mineral N fertilizer (SM); mineral N fertilization (MI). A clear seasonal dynamics of nitrate concentration was observed in the three years and was similar among treatments, with maximum occurring in autumnwinter. On average, at soil depths between 50 and 90 cm, nitrate concentrations in the soil solution were intermediate for the MI treatment (146 \pm 10.4 mg L-1), in between those of SL or SM (202 \pm 11.3 and 164 \pm 9.4 mg L-1 respectively) and MA (78 \pm 4.9 mg L-1). Despite the high average concentrations, only in some sampling dates the nitrate concentration was significantly higher than 50 mg L-1. The estimated annual N leaching losses below 90 cm soil depth ranged from 42 (MA) to 110 (SL) kg N ha-1. These findings highlighted that, under Mediterranean conditions and with

the N input rates, nitrate leaching in autumn-winter cannot be easily controlled through N fertilizer management because it is mainly associated to the natural water surplus and the low N uptake from the winter crop. The cattle manure has proved to be the most conservative in terms of N leaching, while replacing organic with mineral sources of N did not reduce nitrate leaching.

Highlights

- The impact of mineral and organic fertilizers on nitrate leaching was assessed
- Disc lysimeters were used for the nitrate content monitoring in the soil solution
- Cattle manure proved to be the most conservative in terms of nitrate content
- The replacement of organic with mineral fertilizer did not reduce nitrate leaching
- Autumn-winter soil water nitrates were almost independent from fertilizer systems

1	Replacing organic wit	h mineral N fertilization	does not reduce nitrat	te leaching in double crop
<u> </u>	Replacing of guille with	in initional i v iortinization	does not reduce minu	to reacting in aduote crop

- 2 forage systems under Mediterranean conditions
- 3
- 4 Clara Ella Demurtas^a, Giovanna Seddaiu^{a,b*}, Luigi Ledda^{a,b}, Chiara Cappai^a, Luca Doro^a,
- 5 Alberto Carletti^a, Pier Paolo Roggero^{a,b}
- 6 ^a Nucleo di Ricerca sulla Desertificazione NRD, University of Sassari, Viale Italia, 39, 07100,
- 7 Sassari, Italy
- ^b Dipartimento di Agraria, University of Sassari, Viale Italia, 39, 07100, Sassari, Italy
- 9 *E-mail address:* <u>cldemurtas@uniss.it</u> (C.E. Demurtas); <u>gseddaiu@uniss.it</u> (G. Seddaiu);
- 10 <u>lledda@uniss.it</u> (L. Ledda); <u>ccappai@uniss.it</u> (C. Cappai); <u>ldoro@uniss.it</u> (L. Doro);
- 11 <u>acarletti@uniss.it</u> (A. Carletti); <u>pproggero@uniss.it</u> (P.P. Roggero).
- 12
- 13 *Corresponding author: Tel.: +39 079 229392; fax: +39 079 229222. E-mail: gseddaiu@uniss.it.
- 14 Dipartimento di Agraria, University of Sassari, Via E. De Nicola, 07100, Sassari (Italy), (G.
- 15 Seddaiu).
- 16
- 17
- 18
- 19

20 Abstract

This research evaluated the impact of four nitrogen (N) fertilizer management systems on nitrate 21 losses in an irrigated forage system under Mediterranean conditions within a Nitrate Vulnerable 22 23 Zone (NVZ). The experiment was conducted from June 2009 to May 2012 in an intensive dairy cattle farm that produces silage maize and Italian ryegrass in a double cropping system. A 24 monthly monitoring of the nitrate concentrations in the soil solution was carried out using 10 cm 25 diameter disc lysimeters. The N fertilization systems had a target N application of 316 and 130 26 kg ha⁻¹ for maize and Italian ryegrass, respectively. Four systems were compared: cattle manure 27 28 (MA); cattle slurry (SL); cattle slurry + mineral N fertilizer (SM); mineral N fertilization (MI). A 29 clear seasonal dynamics of nitrate concentration was observed in the three years and was similar among treatments, with maximum occurring in autumn-winter. On average, at soil depths 30 between 50 and 90 cm, nitrate concentrations in the soil solution were intermediate for the MI 31 treatment (146 \pm 10.4 mg L⁻¹), in between those of SL or SM (202 \pm 11.3 and 164 \pm 9.4 mg L⁻¹ 32 respectively) and MA (78 \pm 4.9 mg L⁻¹). Despite the high average concentrations, only in some 33 sampling dates the nitrate concentration was significantly higher than 50 mg L⁻¹. The estimated 34 annual N leaching losses below 90 cm soil depth ranged from 42 (MA) to 110 (SL) kg N ha⁻¹. 35 36 These findings highlighted that, under Mediterranean conditions and with the N input rates, 37 nitrate leaching in autumn-winter cannot be easily controlled through N fertilizer management because it is mainly associated to the natural water surplus and the low N uptake from the winter 38 crop. The cattle manure has proved to be the most conservative in terms of N leaching, while 39 40 replacing organic with mineral sources of N did not reduce nitrate leaching. 41

42 Key words: disc lysimeter, silage maize, Italian ryegrass, cattle slurry, manure, Nitrate
43 Vulnerable Zone.

45 Introduction

Intensive, grain-fed livestock production systems are considered a major source of nitrate (NO_3) 46 leaching and pollution. The mitigation of NO_3^- leaching from agricultural soils has become of 47 global outstanding relevance (Hooda et al., 2000). The European Nitrate Directive (ND, 48 1991/676/EEC) intends to reduce diffuse NO_3^- pollution of water bodies by designating Nitrate 49 Vulnerable Zones (NVZs) that need to be managed in a way that prevent NO₃⁻ buildup. The ND 50 established a 50 mg L^{-1} threshold for NO₃⁻ concentration in ground and surface water (European 51 Union, 1998). In NVZs, the use of organic fertilizers is restricted to 170 kg N ha⁻¹ with a 52 53 restricted time window for its application (e.g. between November and February in the 54 Mediterranean area). The only restriction for the mineral fertilizer supply is that of not exceeding crop requirements. This is because mineral fertilization is assumed to have higher efficiency than 55 organic fertilizers (Hernández et al., 2013). Moreover, the ND does not make any distinction in 56 terms of amount or timing between different types of organic fertilizers, assuming that the 57 amount of leached N is not influenced by the organic N source and management. 58 59 The implementation of ND prescriptions has also often resulted in increased costs for crop fertilization and particularly for disposing the excess N from the farm effluents out of the NVZ 60 (de Roest et al., 2007). 61 The impact of different N fertilization sources on NO_3^- leaching is not clearly established since 62 NO₃⁻ dynamics are controlled mainly by a complex set of interrelationships between rainfall 63 patterns (Fang et al., 2013), irrigation (Jamali et al., 2015) and fertilization management 64 practices, as well as by soil type and cropping systems (Jabloun et al., 2015; Barros et al., 2012). 65 In many NVZs, the increased demand for high-quality milk has led to an evolution of the dairy 66

67 cattle housing from the bedded-pack to the cubicle housing system. This resulted in a higher

68 production of slurry than farmyard manure (Phillips, 2010), even though the bedded pack system

69 was recognized to be compatible with high quality milk and better animal welfare (Atzori et al.,

70 2009 a).

71 The development of site-specific sustainable fertilization management options for mitigating

 NO_3^{-} pollution is therefore a key priority for policy decision makers and land managers

73 (Zavattaro et al., 2012).

74 The NO₃⁻ pollution is an issue in Mediterranean semi-arid countries (Daudén et al., 2004). In the

75 Mediterranean regions of the northern hemisphere, 50-90% of total NO_3^{-1} losses by leaching were

found to occur particularly between October and February (De Paz et al., 2009; Trindade et al.,

1997; Goss et al., 1988). The application of N fertilizers in autumn reinforces the natural

accumulation of NO_3^- in the soil at the start of the rainy period (Trindade et al., 2009), thus

79 promoting significant N leaching (Trindade et al., 1997). The dynamic of these processes relates

80 to both the N source and application scheme.

Experiments in Mediterranean conditions showed that the NO₃⁻ leaching potential from the 81 application of slurries could be lower than those from N mineral fertilizer application (Trindade 82 et al., 2009; Daudén and Quílez, 2004). However, it was also recognized that NO₃⁻ leaching can 83 be a serious problem when organic effluents are used (Giola et al., 2012; Daudén et al., 2004; 84 85 Chambers et al., 2000; Trindade et al., 1997). Hence it is still an open question whether the replacement of organic with mineral N fertilizers can effectively mitigate NO₃⁻ leaching in NVZ 86 and if the type of organic effluent can make a difference in improving the environmental 87 88 sustainability of intensive dairy farming under Mediterranean conditions. This issue is

89 particularly relevant for intensive livestock farming areas under Mediterranean conditions, like

90 the district of Arborea (Sardinia, Italy). This farming district is characterized by very high

91 stocking rate and high productivity of its dairy cattle system, one of the highest in Europe

92 (Manca, 2009), with a stocking rate of some 5 livestock unit ha^{-1} . In these conditions, the annual

animal effluent production is somehow double than the maximum applicable to farmland

94 according to the ND (Nguyen et al, 2014). This study showed that to comply with the ND,

95 Arborea farmers must buy additional mineral N fertilizer to meet the total N crop requirements

96 and must also rent additional land or pay to dispose the cattle effluents outside the NVZ, thus

- 97 increasing the production costs. This situation is clearly unsustainable and has given rise to
- 98 controversy about the effectiveness of the ND measures to mitigate NO_3^{-} pollution while
- 99 maintaining the profitability of the dairy farming.
- 100 The objective of this study was to compare a range of N fertilization options for dairy farming
- 101 systems in NVZ under Mediterranean conditions, to support the development of sustainable
- 102 fertilization strategies by exploiting the recycling of the farm N resources.
- 103 In this paper, we tested two hypotheses: i) in the Mediterranean intensive forage systems, the
- seasonal dynamics of the NO_3 concentrations in the soil solution is very wide and can be
- 105 effectively controlled by the fertilization management systems; ii) the mineral fertilizers are less
- 106 impacting on NO_3^{-1} losses than organic fertilizers in NVZ.
- 107 We measured the dynamics of the NO_3^- concentration in the soil solution and assessed the
- 108 potential N leaching as influenced by four N fertilization management systems for an irrigated
- 109 double forage cropping system in sandy soil under Mediterranean conditions.
- 110

111 1. Material and methods

112 **1.1. Site and crop management**

113 The field experiment was conducted from June 2009 until May 2012 in a private farm located in the NVZ of Arborea, Italy (39°47' N, 8°33' E, 3 m asl). In the district of Arborea, some 28,000 114 115 Bovine Livestock Units are reared by 160 farms on a 5,000 ha irrigated plain. This plain was drained in the 1930's from wetlands and salt marshes. The groundwater table dynamics is 116 117 somehow regulated by the local Water User Association dewatering pump system, which controls the drainage of the whole area. The groundwater table depth was monitored during the 118 119 three year of experiment through 12 piezometers installed in the experimental field. 120 In this area, about two thirds of the livestock effluent volume is represented by slurry and one third by manure. Considering the N content of the two effluents, the N load from manure and 121 122 slurry in this NVZ is approximately 38 % and 62 %, respectively.

123 The climate is Mediterranean and the mean annual temperature and precipitation are 16.7 °C and 568 mm, respectively (1959-2012). Some 73 % of the annual rainfall occurs between October 124 125 and March and the average annual aridity index (rainfall/reference evapotranspiration) is 0.49 126 (semi-arid area). The soil was classified as Psammentic Palexeralfs (USDA, 2006) and the soil 127 slope was less than 0.5 %. The soil properties are reported in Table 1. The field capacity of soil 128 profile was determined according to Romano and Santini (2002). The water balance dynamic 129 from May 2009 to June is shown in Figure 1. The common forage cropping system is based on 130 the double-crop rotation of silage maize (Zea mays L.) grown from June to September and Italian 131 ryegrass (Lolium multiflorum Lam.), grown from October to May. In this cropping system, 132 organic fertilizers (slurry and manure) are applied regularly prior to sowing of each crop and 133 combined with mineral fertilization. During the Italian ryegrass cycle, mineral fertilizers are 134 often applied at the end of the winter to compensate for insufficient soil mineral N availability, 135 while maize is fertilized in summer either at seeding or at the 4-6 leaves stage. We compared four fertilization systems at the same target N rate (316 and 130 kg ha⁻¹ for maize 136 137 and ryegrass respectively), set on the basis of the N fertilization prescriptions for NVZs, and of 138 the business as usual local fertilization practices, which should correspond to the expected N uptakes of silage maize and Italian ryegrass. The experimental design was a randomized 139 140 complete block design with four replicates and a plot size of $12 \text{ m} \times 60 \text{ m}$. The treatments were: 141 i) manure (MA): cattle farmyard manure applied before the sowing of each crop with a 142 conventional spreader and followed by rotary tillage. About 70 % of the total amount was spread 143 to maize at the end of May and 30 % to ryegrass in October; 144 ii) slurry (SL): cattle slurry applied before sowing with a conventional spreader and immediately 145 incorporated into the soil through rotary tillage using the same proportions as the MA treatment 146 for maize and ryegrass;

147 iii) slurry+mineral (SM): following the NVZs prescriptions, slurry was applied at a

148 corresponding target rate of 100 and 70 kg ha⁻¹ of N for maize and ryegrass respectively and

- mineral fertilizer (ENTEC 26[®]) at a rate of 216 and 60 kg ha⁻¹ of N applied before sowing for
 maize and at the end of ryegrass tillering respectively;
- iv) mineral (MI): mineral fertilizer (ENTEC 26[®]) applied before sowing for maize and at the end
 of ryegrass tillering (mid February beginning of March).

153 The amount of total N rate actually supplied with organic fertilizers (Table 2) was calculated ex 154 post on the basis of the volume of fertilizer distributed and its actual nutrient concentration. The 155 fertilizer was sampled from the field spreader at each distribution event, for a total of six 156 sampling dates. Four and eight samples per sampling date were collected respectively for manure 157 and slurry and stored frozen until the determination of total N content. The cattle slurry was 158 stored in concrete storage pits at the farm and was homogenized prior to surface application 159 using a vacuum-tank spreader. The cattle manure was stored for five months before spreading 160 and it was sampled just before application.

161 A mixture of four varieties and hybrids of Italian ryegrass (cultivars Meritra, Ivan, Littorio and 162 Mowester) was sown around mid-October each year. Seedbed was prepared by harrowing and 163 milling. Auxiliary sprinkler irrigation was provided to ryegrass when necessary to minimize crop water stress (a total of about 80 mm per cycle). Ryegrass hay was harvested in mid-May at early 164 165 earing stage. The Calcio hybrid (FAO 700) of maize was sown during the first ten days of June 166 and it was harvested between the 15th and the 20th of September at the dough stage (target of 33 167 % DM for the whole plant). The maize seedbed was prepared using a rotavator, ripper and harrow. The sprinkler irrigation for maize started at seeding in June until one week before 168 harvest. The total water volumes were on average 4600 m³ ha⁻¹. Weeds were controlled with the 169 pre-emergence herbicide Lumax[®]. 170

171

1.2. Soil solution sampling and analysis

Two tension disc lysimeters (10 cm diameter, PRENART[®]) were installed in each plot below the
Ap soil horizons, at soil depths between 50 and 90 cm (average depth 68 cm). A total of 32 disc
lysimeters were installed. Special care was taken in repacking the soil from different layers in the

175 original position. The disc lysimeters were connected to above ground with plastic pipes at the border of each plot. Near the soil surface a 20 cm rubber collar was put around the plastic pipe to 176 177 prevent water from seeping from the surface down through the pipe. The soil solution was 178 sampled through the disc lysimeters at monthly intervals by applying a suction of -70 kPa for 179 about 45 minutes using an electric pump. Before collecting the sample to be analyzed, pipes were cleaned out of the water left from the previous sampling in the connecting pipe and disc 180 181 lysimeter. Collected samples were stored under cooled conditions (4 °C) and filtered in the lab 182 through a 0.2 μ m membrane filter. The NO₃⁻ concentration was determined by ion 183 chromatography (anion column Alltech model allsep anion 7 µm, 100 mm).

184 **1.3. Water balance**

185 The water balance of the maize-Italian ryegrass cropping system was estimated using the EPIC 186 model (Williams, 1995). The model was set using local soil and weather data, information about 187 the actual management, including irrigation volumes and amount of fertilizers distributed, and crop related data such as plants density and crop growing period. EPIC simulated actual 188 189 evapotranspiration from the Penman Monteith equation considering the actual crop management. 190 Daily weather data were collected from a weather station located about 5 km away from the experimental field (39°45' N; 8°34' E. 13 m asl, OR, Italy). Soil characteristics were obtained 191 192 analyzing soil samples collected in the experimental field and water holding capacity was 193 measured in the field through systematic gravimetric and tensiometric measurements between saturation and field capacity. The actual physical and chemical soil characteristic (including the 194 195 hydraulic properties) were used as inputs in the EPIC model. Because the groundwater table 196 depth is controlled by the local Water Use Association (WUA), part of the subroutine that 197 simulates the groundwater table dynamics in EPIC was modified considering a deeper 198 groundwater table in spring and summer and a shallower groundwater table in autumn and 199 winter. The modification in EPIC allowed to overwrite the initial maximum and minimum depth 200 for the groundwater table fluctuation and to use different maximum and minimum depths in

201 different periods of the year. The model was calibrated for the crops yield and ground water table dynamics. For each crop a total of 12 yield observations (3 years \times 4 replicates) were available. 202 203 Six observations were used for the calibration and the remaining six observations were used for 204 the validation. For the groundwater table depth, a total of 24 observations were available from 205 November 2009 to May 2012. The first 12 observations were used for the calibration and the last 12 observations were used for the validation. The calibration and validation were evaluated using 206 relative root mean square error (RRMSE; Bellocchi et al., 2002), modeling efficiency (EF; 207 208 Loague and Green, 1991), slope and intercept of the regression line and coefficient of determination (R²). After the calibration and validation, the EPIC model was used to estimate the 209 210 percolation below the average lysimeters depth. The calculation of the total percolation, including groundwater table fluctuations, was performed for those days in which a water surplus 211 (Precipitation + Irrigation > ET) occurred. 212

213

1.4. Nitrate leaching and N balance

Assuming that the soil water percolation between two soil water sampling dates had the average

215 NO_3^- concentration of the sampled soil water, the amount of NO_3^- leached (kg N ha⁻¹) was

216 calculated using the average NO_3^{-} concentration between two subsequent sampling dates times

and the volume of percolation between the two dates as estimated with EPIC.

218 The cumulative N leaching was estimated on the basis of average NO_3^- concentration and the

soil water percolation (as simulated by the EPIC model) for each sampling date.

Crop yield, aboveground biomass at harvest and N removal were measured every year but in this
paper we report only the data on crop N removal. The entire plots were harvested using farm
machineries and the removed products (fresh aboveground maize or ryegrass hay) were weighed
immediately using electronic weighing cells positioned in a flat place under the tractor cart. The

biomass dry matter content at harvest was assessed by sampling 1 kg of ryegrass hay or fresh

chopped maize that was immediately cooled in plastic bags and hence dried in a forced-air oven

at 65 °C for 72 h. The crop N removal was estimated by multiplying the crop yield and N

concentration determined on harvested biomass samples by the Kjeldahl method (Mipaaf, 2006).

228 N surplus was defined as the difference between N fertilizer input at plot scale and crop N

removal (Grignani and Zavattaro, 2000). While, combining the data on crop N removal and

230 leached N we also estimated by difference the residual soil N pools available for N

- 231 immobilization, volatilization or denitrification.
- **1.5. Statistical analysis**

The NO₃⁻ concentration and yearly cumulative N (removal, leaching and residual) were analyzed 233 234 with the PROC MIXED procedure in SAS (SAS institute, 1999), suitable for analyzing mixed 235 effects and repeated measures with non-constant variance and any covariance structure models. 236 The independence assumption on the error terms required for the ANOVA of a factorial model (Montgomery, 1997) was likely not met. Therefore, the appropriate assumption on the error 237 238 terms for this experiment was the non normal distribution with heterogeneous (non-constant) 239 variance by sampling date and a given covariance structure. The appropriate covariance structure 240 for this particular experiment was determined using -2 Log Likelihood (-2RLL), Akaike's 241 Information Criterion (AIC), Corrected Akaike's Information Criterion (AICC) and Schwarz's Bayesian Criterion (BIC), which are essentially log likelihood values penalized for the number 242 of parameters estimated (Littell et al., 1996). Also, according to the fit statistic -2RLL, AIC 243 244 AICC and BIC, the best covariance structure was determined to be the Compound Symmetry 245 covariance type. The statistical test results given in the following section are based on the transformation of original NO₃⁻ concentration data as reported Shen et al. (2006). 246 247 A one-tailed t test was performed on data from each sampling date to determine whether treatments significantly exceed the EU threshold of 50 mg L^{-1} of NO₃⁻ concentration using the 248 249 pooled error variances when appropriate. 250 All data analyses were performed using Microsoft Excel® and SAS Statistical Package (SAS

251 institute, 1999).

252

253 **2. Results**

- 254 **2.1.** Nitrate concentration in the soil solution
- 255 The average concentration of NO_3^- in the investigated soil depths showed a high variability and
- 256 it fluctuated from 0.8 ± 0.3 (mean \pm SE, June 2010) to 586 ± 122 (November 2011) mg L⁻¹ in
- 257 MA, from 6 ± 3 (May 2010) to 593 \pm 104 (January 2012) mg L⁻¹ in SL, from 13 \pm 3 (June 2009)
- 258 to 460 ± 54 (December 2011) mg L⁻¹ in SM and from 2.6 ± 0.4 (June 2009) to 390 ± 25
- 259 (December 2010) mg L^{-1} in MI (Figs. 2-4).
- 260 The dynamics of NO_3^- concentration (mg L⁻¹) in the soil solution showed a strong seasonal

261 pattern for all treatments, characterized by two maximum values, one in the winter, during the

- ryegrass crop phase, and one in the summer, during the maize crop phase (Figs. 2-4). Minimum
- values were typically observed at ryegrass and maize harvest, in late spring and September. In all

treatments, the NO_3^- concentrations decreased to reach a minimum in April and May, with the

exception of spring 2012 (Fig. 4), when the observed NO_3^- concentrations remained high until

the end of April in all treatments. Plots receiving the MI treatment showed high NO_3^{-1}

- 267 concentration in all autumns, even though N fertilizer was not applied until February.
- In the three years, the mean NO_3^- concentration significantly exceeded the 50 mg L⁻¹ on 5, 14,
- 269 15 and 8 dates out of the total 33 sampling dates in MA, SL, SM and MI respectively (Figs. 2-4).
- 270 Of these dates, 5/5, 8/14, 7/15 and 7/8 respectively in MA, SL, SM and MI occurred in the

271 November-February period and the remaining occurred in the spring 2012.

272 Of the 33 sampling dates, only 6 showed a significant difference between treatments. In

- 273 particular, 4 dates in the third year (Jan-Mar-Apr-May) had higher NO₃⁻ concentration in SL
- 274 with respect to MI and MA, while SM was not significantly different from all the other
- treatments. In December 2010 the NO_3^- concentration was lower in MA treatment than in SL
- and MI treatments, despite the mineral fertilizer was not applied yet.

The lowest average NO_3^- concentration was observed in MA. In 2009 and 2010 MA showed an average annual NO_3^- concentration of about $46 \pm 16 \text{ mg L}^{-1}$, but in 2012 the NO_3^- dynamics of MA was similar to that observed in the other treatments (Fig. 4).

280 **2.2. EPIC model calibration and validation**

281 Model performance for crop yields and groundwater table dynamics were satisfactory for all the data considered and for both the calibration and validation phases (Table 3). For silage maize the 282 283 RRMSE values indicate small deviations of the simulated values from the observed ones and the 284 EF > 0 indicates that the model is a good predictor of the observed values. For Italian ryegrass 285 the deviations of the model results from the observed values were acceptable in terms of RRMSE 286 and good for EF values. The model was also able to reasonably simulate the variation during time of the groundwater table depth. The deviation of the simulated data from the observed data 287 288 was about 26 % and 20 % during the calibration and validation process respectively and the EF 289 value was of 0.37 for the calibration and 0.54 for the validation. Moreover, the slope and 290 intercept of the regression line were close to the optimal values.

291 **2.3.** Soil water percolation

292 The estimated average annual percolation simulated by EPIC was 252 mm, 292 mm and 268 mm during June-May in 2009-2010, 2010-2011 and 2011-2012, respectively. In all years, percolation 293 294 events occurred almost exclusively from October to February with an influence of cumulative 295 monthly precipitation (y) on estimated cumulative monthly percolation (x), as evidenced by the significant linear regression between these two variables (y = 0.327x + 4.891; R²=0.66**). The 296 297 average soil water percolation in October-January was about 130 mm and it represented 1/2 of the total annual percolation (60 % between Oct-Feb). The rainfall in autumn 2010 (392 mm in 298 299 Nov-Dec) was considerably higher than the long term average recorded for the study area (158 300 mm) and this was mirrored by the calculated drainage volumes (Fig. 3). From April to September no relevant drainage was estimated, the percolation was on average about 60 mm as 301 302 the total water supply (730 mm = 194 mm rain and 536 mm irrigation) rarely exceeded crop

transpiration demands (670 mm). The groundwater table depth ranged between 0.25m (in

304 January-February) and 1.80 m in July-August. Groundwater table depth reached the highest

305 values from June to September, while the water table was the shallowest during the autumn from

- 306 November to February in coincidence with rainfall events.
- 307

2.4. Nitrate leaching and N balance

308 The average N concentration of slurry was 0.26 % but ranged between 0.09 and 0.47, while that

of manure was 0.71 % (0.45 - 0.83). The average ammonia N concentration was 0.12 % in slurry

and 0.17 % in farmyard manure. The nitric nitrogen was 0.002 % in slurry and 0.01 % in

311 manure.

312 The N input of organic fertilizers varied widely (Tab. 2) according to the variable N

313 concentration of the animal effluent.

The C:N ratio and the organic N concentration in the animal effluents were respectively $8.01 \pm$

315 0.85 (mean \pm SE) - 0.14 \pm 0.01 % for slurry and 13.02 \pm 1.17 - 0.51 \pm 0.04 % for manure.

316 The estimated N leaching and residual N were significantly influenced by the fertilization

treatment (P < 0.05) and no significant year \times treatment interaction was observed (data not

shown). The estimated residual N was variable among years, particularly for SL. The exception

319 was the MI treatment, in which residual N variability was much lower than all other treatments,

320 likely due to a lower and more constant rate of N inputs relative to that observed for organic

fertilizers (r = 0.47, ns). For this reason, N inputs on SL and SM explained as much as 97 % of

322 the variability of the residual N (P < 0.01), while there was no linear correlation between N input

and N leaching considering all treatments (r = 0.42) or between N surplus and N leaching (r = 0.42)

324 0.30, ns). The estimated annual N leaching losses below 90 cm soil depth were reported in Table

4. The relative proportion of the N surplus lost with leaching was high (62 %) just for MI while

it represented less than about 17 % of the surplus when organic fertilizers were used.

327 On average, N removal roughly represented between 1/3 (SL) and 1/2 (SM) of the N input in all

328 treatments with organic fertilizers, while it reached 2/3 of the total N input in MI (Tab. 4). N

329 leaching in MA was 32 % lower than the average of the other fertilization systems.

330 The dynamics of the cumulated N leaching showed the same pattern in the three years of

experiment (Fig.5). On average, 78 % of the total N leaching was in autumn (44 %) and winter

332 (34 %). Almost no soil water percolation and hence no leaching was recorded during the maize

333 crop phase, despite the relatively high NO₃⁻ concentration in soil solution, but we estimated a

high percolation in April.

335 The total cumulated annual water drainage simulated by EPIC was reported in previous

paragraph. If the NO_3^{-1} losses were expressed as a function of the cumulative drainage (data not

shown), the loss of NO_3^- corresponding to a drainage concentration of 50 mg L⁻¹ of NO_3^- was

28.4, 33.1 and 25.9 kg ha⁻¹ in year 1, 2 and 3 respectively. The estimated NO₃⁻ losses reached

these thresholds between 125 mm and 170 mm of drainage in MI and MA respectively in the

first year; between 60 mm and 125 mm in SL and MA in the second year and about 35 mm for

all treatments in the third year.

342 **3. Discussion**

The NO₃⁻ concentration dynamics in the soil solution was very variable during the year and was 343 344 influenced by weather and crop N uptake. This was consistent with our first hypothesis. About 345 68 % of the average total annual NO_3^- leaching occurred between October and February, which corresponds to the period in which 57 % of the annual percolation occurred, as a consequence of 346 347 the unbalance between rainfall and evapotranspiration, as usual under Mediterranean conditions (Carneiro et al., 2012; Arregui and Quemada, 2006). In this period, N uptake of Italian ryegrass 348 349 is very low as the crop is in the establishment-tillering phase (Carneiro et al., 2012; De Paz et al., 350 2009).

A wide variation in N concentrations in soil water collected using the same methodology under
 similar agro-environmental conditions of this study was also observed by Carneiro et al.(2012)

and Trindade et al. (1997). The wide variation in the soil water NO_3^{-1} concentration at a given 353 sampling time was attributed to the heterogeneous distribution of fertilizers into the soil, to the 354 355 soil composition variability and the corresponding expected uneven infiltration pattern (Cuttle et al., 1992). However, the wide variability of soil water NO₃⁻ concentration may also be related to 356 357 microbiological processes (Liang et al., 2010). Moreover, the high N content variability of 358 manure and slurry determined *a posteriori* in our study, led to application of a higher N supply 359 than the amount planned, and, hence, higher than crop requirements. This unintentional over-360 application led to a high N surplus, as similarly reported also by Trindade et al. (2009), generating a source of difficulty for the accurate comparison of NO_3^- concentration data in the 361 362 soil solution among the four N fertilization systems. The N input from SL and SM treatments was significantly higher than that from MI treatment, while the nitrate leaching rates for SL, SM 363 364 and MI were similar. This was interpreted as possibly due to the ammonium Nimmobilization through microbial decomposition of organic matter in the slurry after application, that plays a 365 366 significant role in the organic N retention in soil (Sørensen, 2004). This immobilised N is 367 stabilised and slowly released a few months after application (Sørensen, 2004). We can then 368 assume that much more N was immobilized after slurry or manure incorporation than after 369 mineral fertilizer application. Moreover, the significantly lower N leaching achieved from MA 370 treatment could be associated to the higher C:N ration of farmyard manure than slurry. However, 371 neither the N input nor the N surplus showed a significant linear correlation with N leaching considering all treatments, similarly to what reported by Morari et al. (2012) and Zavattaro et al. 372 373 (2012). In contrast to our results, Silgram et al. (2001) reported that the leaching losses are 374 linearly related to N inputs, over-simplifying a complex N loss function which depends on the 375 interactions between over-winter rainfall, soil type, cropping, and the rate/timing of 376 fertilizer/manure applications. Since in our study the NO_3^- leaching was estimated considering 377 only losses associated to percolation in the days when water surplus occurred (i.e. 378 Rain+Irrigation-ET), it is likely that the total N losses were underestimated. However, the rate of estimated N leaching was not affected neither by the N input or N surplus and the percolation was not very high, on average 271 mm per year. The shallow groundwater in autumn-winter could have contributed with washing away to the N leaching rate, in particular under the winter crop as shown by the drop of NO_3^- concentration at the end of the winter, while the contribution of the deep groundwater table to NO_3^- leaching and maize water requirements was assumed to be negligible.

The average monthly NO_3^- concentration was lower in April and May, except for the third year, when we observed low winter rainfall (and percolation) and higher NO_3^- concentrations from December to February and in July, as found for other irrigated Mediterranean cropping systems (Ibrikci et al., 2015). In particular, the high NO_3^- concentrations were recorded after fertilization and precipitation/irrigation events, in agreement with what found by Perego et al. (2012) in summer on maize.

391 In the present study, the observed high soil water NO_3^- concentration in the autumn was 392 independent of the fertilization system, hence the NO₃⁻ concentrations in the soil solution were 393 not effectively controlled by the fertilization management. The potential impact of the presence 394 of the nitrification inhibitor under MI was assumed to be negligible below a soil depth of 40 cm 395 as reported Yu et al. (2007). In this period, the MI treatment had not been fertilized yet, but showed high NO₃⁻ concentration. This could be interpreted hypothesizing that the maximum 396 NO₃⁻ concentrations observed in autumn were the outcome of the accumulation of NO₃⁻ in the 397 soil after the maize harvest, in a period in which soil temperature and water content were not 398 399 limiting for microbial activity (Liang et al., 2010), crop uptake was zero and the maize crop 400 residues were incorporated into the soil before ryegrass seeding. Although the residual amount of soil NO₃⁻ after maize harvest was not measured in this experiment, we estimated it from the 401 NO₃⁻ concentration between two subsequent sampling dates (September and November) across 402 403 maize harvest dates and the soil water percolation volume estimated by EPIC. The outcomes of this estimate showed that on average the estimated NO_3^- at maize harvest ranged from 10 kg N 404

ha⁻¹ to 30 kg N ha⁻¹ and this is in accordance to the results obtained from the application of the linear relationship proposed by Andrasky et al. (2000) for 1.3-1.5 m soil depth. Several studies reported that silage maize often shows high residual mineral soil N amounts ranging from 48 kg N ha⁻¹ under low N input up to 278 kg N ha⁻¹ under high fertilized treatments (Morari et al., 2012; Kayser et al., 2011) leading to high over-winter NO₃⁻ leaching (Trindade et al., 2009; Le Gall et al., 1997; Simon and Le Corre, 1988).

411 The estimated N leaching loss is within the range reported by Carneiro et al. (2012), while sometimes we found that the mean annual leaching was lower than that reported by Perego et al. 412 (2012) in Northern Italy, where the annual drainage (about 350 mm year⁻¹) was higher than that 413 414 observed in our experiment (about 271 mm year⁻¹). The magnitude of the losses was mainly determined by the NO₃⁻ concentration in the soil solution in November and December. 415 The N mineralized between October and March may also have represented an important NO₃⁻ 416 417 source. Vertes and Decau (1992) estimated that 90 % of N leaching losses were originated from 418 N mineralization. Therefore, these authors recommended for preventing high soil NO₃⁻ 419 concentration in the autumn to apply conservative fertilization and cropping practices for the 420 summer crop to avoid high residual-NO₃⁻ values, in order to favour the early establishment of 421 the autumn–winter crops or to minimize the amount of N supplied at the winter crop, particularly 422 when sown late (Carneiro et al., 2012). The same authors suggested that the mineral N should be 423 applied only through top-dressing applications. This may explain why in similar agroenvironmental conditions, it was observed that N applied at tillering was recovered more 424 425 efficiently than that applied at emergence (López-Bellido et al., 2005; Kirda et al., 2001). In agreement with this study, Trindade et al. (2009) reported that the treatments where mineral-N 426 fertilizer was applied had higher soil NO_3^{-1} -leaching potential after the maize crop. Moreover, 427 Smith and Chambers (1993) suggested that wherever cool and wet winters prevail, N leaching 428 can be reduced using organic fertilizers with a low mineral to organic N ratio, as farmyard 429

manure. On the contrary, Shepherd and Newell-Price (2013) reported that annual applications of

430

431 farmyard manure increased NO_3^- leaching by 39 % more than the inorganic fertilizer, under a 7-

432 yr rotation (wheat-potatoes, wheat-winter barley, sugar beet- wheat). This could be related to the

433 site-specific dynamics of the soil organic matter: in this study, the soil organic matter

434 accumulated over the years from long-term organic fertilization may have acted as a buffer of

435 organic N available for mineralization (Indraratne et al., 2009; Kaur et al., 2008; Gutser et al.,

436 2005; Schröder et al., 2005; Hao et al., 2003).

437 The findings of this study suggest that under Mediterranean conditions N fertilization in autumn 438 could be omitted to mitigate the leaching of NO_3^- in the autumn-winter period without affecting 439 crop yields. López-Bellido et al. (2012) showed that the postponing of the N fertilizer application 440 to tillering-stem elongation phases for winter crops is usually more efficient in terms of plant N uptake. Besides, this is confirmed by the findings of Trindade et al. (2009) who reported no 441 442 significant effects on Italian ryegrass annual dry matter yields of the N application in October compared to no N fertilization using similar total N inputs to our study. Moreover, despite the 443 ND prescriptions are forbidding the supply of organic or mineral fertilization from November to 444 445 February, this practice did not mitigate the NO_3^{-} leaching.

Studies conducted in the Netherlands under maize and grassland crops reported that about 25 % 446 447 of the total N input applied as cattle slurry was lost by leaching and it was independent of its rate 448 application (Schröder and Dilz, 1987). So that the N losses depends on the source as well. An 449 investigation by Worrall et al. (2009) showed the lack of objective success in the majority of the 450 32 NVZs studied in the United Kingdom. This has called into doubt the use of input management 451 as a means of limiting NO_3^{-} pollution, especially when the areas designated are small. The lack of objective success for NVZ designation suggests that NO₃⁻ pollution control strategies based 452 453 only on input management need to be rethought.

454 The residual N input that is not uptaken by the crop or leached, can be subjected to various

455 processes such as volatilization (Atzori et al, 2009), immobilization (Divito et al., 2011; Kaur et

456 al., 2008) or denitrification (Morari et al., 2012). However the assessment of the relative

457 contribution of these processes to the fate of the residual N was not the object of this study and is458 worthy of further studies.

459 The lowest average NO_3^{-} concentration was observed in MA and this was associated to the

460 higher C:N ratio and proportion of organic N in farmyard manure than in slurry. Similar findings

461 were reported by several scholars under a range of agro-environmental conditions (Sänger et al.,

462 2010; Bertora et al., 2008; Moral et al., 2005). Ever more frequently the amount of manure

463 produced in this farming livestock system decreases when cattle management shifts from a bed-

464 pack system to cubicle housing as occurred in this study area. In fact the production of manure

465 instead of slurry is related to structural changes of the livestock housing system, which in turn is

466 indirectly related to farming targets of milk quality and animal welfare.

467 However, the bed-pack management and particularly the straw availability are key components

468 for achieving good milk quality (Wenz et al., 2007) and animal welfare standards (ADAS, 1994;

469 Singh et al., 1994) that can be compatible with the production of manure, which proved to be

470 much more sustainable in terms of environmental impacts of intensive dairy farming under

471 Mediterranean conditions.

472

473 **4.** Conclusions

The study supported the hypothesis that the NO_3^- dynamics in the soil water in a maize-ryegrass double cropping forage system under Mediterranean conditions are variable, independently of the fertilization system. This implies that the NO_3^- losses can be only partially controlled by the fertilization system, while the complex interactions between the winter rainfall, the crop sequence and the soil organic N buffer are far more important.

479 Among the fertilization systems under comparison, the cattle manure proved to be the most

480 conservative, while the replacement of organic with mineral N fertilization systems was not

481 effective in mitigating NO₃⁻ leaching, despite the lower yearly residual N estimated.

The maximum soil water NO_3^{-1} concentrations observed in the autumn-winter were almost 482 independent from the type of fertilizer and actual N input rate. This was explained by the 483 484 impossibility of controlling effectively the leaching associated to the natural water surplus in winter and the observed soil water NO₃⁻ concentration dynamics in autumn-winter. It seems that 485 the autumn-winter NO_3^{-} leaching in the maize-ryegrass double cropping is unavoidable in highly 486 fertilized systems. The fertilization with mineral N instead of organic N does not reduce N 487 leaching in winter. The average annual NO₃⁻ losses and soil water NO₃⁻ concentrations were 488 489 significantly lower using 100% farmyard manure as N fertilizer. This may have relevant 490 implications on the livestock effluent management system at the farm scale (e.g. from cubicle 491 housing to straw bedded-pack), that could be addressed to obtain fertilizers with higher C:N ratio in order to minimize the release of NO_3^- during the leaching period. However the organic 492 493 fertilizers must be carefully managed to prevent over- or under-application and to account for the cumulative environmental effects of application as well as storage. 494

495

496 Acknowledgements

This study was carried out within the Agroscenari project (2008-2013) funded by the Italian 497 498 Ministry of Agricultural, Food and Forestry Policies and the PRIN-ZVN project (2008–10). We 499 are grateful to Azienda Sardo farm for hosting our experimental activities and the Arborea 500 Farmers' Cooperative for their fruitful collaboration during this study. A special thanks is 501 provided to Dr. Armen Kemanian, Penn State University for his valuable comments on the early 502 stage of the manuscript. This research formed a part of a PhD program on "Crop productivity" carried out at the Dipartimento di Agraria of the, University of Sassari, Italy. Furthermore, we 503 504 thank Dr. Giulia Urracci and Dr. Laura Mula for their collaboration in the design of the 505 experiment and for the support in the field activities. Authors are also grateful to Alberto Stanislao Atzori for his support on the calculations of the N loads from livestock effluents at case 506 507 study scale.

509 **References**

- 510 ADAS, 1994. The design of dairy cow housing. CIGR Section II: Working Group Cattle
- 511 Housing, ADAS, Winchester, United Kingdom, 1994.
- 512 Andraski, T.W., Bundy, L.G., Brye, K.R., 2000. Crop management and corn nitrogen rate effects
- on nitrate leaching. J. Environ. Qual. 29, 1095–1103.
- 514 Arregui, L.M., Quemada, M., 2006. Drainage and nitrate leaching in a crop rotation under
- 515 different N-fertiliser strategies: application of capacitance probes. Plant Soil 288, 57–69.
- 516 Atzori, A.S., Carta, P., Pulina, G., 2009 a. Le pratiche che incidono sulla diffusione delle zoppie.
- 517 Supplemento a L'Informatore Agrario 21, 33-36.
- 518 Atzori, A.S., Boe, R., Carta, P., Fenu, A., Spanu, G., Francesconi, A.H.D., Cannas, A., 2009 b.
- 519 Estimation of nitrogen volatilization in the bedded-pack of dairy cow barns. Ital.J.Anim.Sci. 8
- 520 (Suppl. 2), 253-255.
- 521 Barros, R., Isidoro, D., Aragüés, R., 2012. Irrigation management, nitrogen fertilization and
- 522 nitrogen losses in the return flows of La Violada irrigation district (Spain). Agric. Ecosyst.
- 523 Environ. 155, 161–171.
- Bellocchi, G., Acutis, M., Fila, G., 2002. An Indicator of Solar Radiation Model Performance
 based on a Fuzzy Expert System. Agron. J. 94(6), 1222–1233.
- 526 Bertora, C., Zavattaro, L., Sacco, D., Monaco, S., Grignani, C., 2008. Soil organic matter
- 527 dynamics and losses in manured maize-based forage systems. Eur. J. Agron. 30, 177–186.
- 528 Carneiro, J.P., Coutinho, J., Trindade, H., 2012. Nitrate leaching from a maize × oats double-
- 529 cropping forage system fertilized with organic residues under Mediterranean conditions. Agric.
- 530 Ecosyst. Environ. 160, 29–39.
- 531 Chambers, B.J., Smith, K.A., Pain, B.F., 2000. Strategies to encourage better use of nitrogen in
- animal manures. Soil Use Manage. 16, 157–161.
- 533 Cuttle, S.P., Hallard, M., Daniel, G., Scurlock, R.V., 1992. Nitrate leaching from sheep-grazed
- 534 grass/clover and fertilized grass pastures. J. Agric. Sci., Cambridge 119, 335–343.

- 535 Daudén, A., Quílez, D., 2004. Pig slurry versus mineral fertilization on corn yield and nitrate
- 536 leaching in a Mediterranean irrigated environment. Eur. J. Agron. 21, 7–19.
- 537 Daudén, A., Quílez, D., Vera, M.V., 2004. Pig slurry application and irrigation effects on nitrate
- leaching in Mediterranean soil lysimeters. J. Environ. Qual. 33, 2290–2295.
- 539 De Paz, J.M., Albert, C., Delgado, J.A., 2009. NLEAP-GIS modelling in a Mediterranean region
- of Spain, in: Grignani, C., Acutis, M., Zavattaro, L., Bechini, L., Bertota, C., Gallina, P., Sacco,
- 541 D. (Eds.), Proceedings of the 16th Nitrogen Workshop Connecting Different Scales of Nitrogen
- 542 Use in Agriculture. Turin, 515–516.
- 543 de Roest, K., Montanari, C., Corradini, E. 2007. Il costo della Direttiva Nitrati . BIT SpA- Cassa
- 544 Padana. Leno (Brescia). Emilia Romagna: Centro Ricerche Produzioni Animali.
- 545 European Commission, 1991. Commission decision of 12 December 1991 concerning the
- 546 rotection of waters against pollution caused by nitrates from agricultural sources, 91/676/EC, in:
- 547 Official Journal, L 375, 31/12/1991, 1-8.
- 548 Divito, G.A., Sainz Rozas, H.R., Echeverría, H.E., Studdert, G.A., Wyngaard, N., 2011. Long
- term nitrogen fertilization: Soil property changes in an Argentinean Pampas soil under no tillage.
- 550 Soil & Tillage Research 114, 117–126.
- 551 European Union, 1998. Council Directive 98/83/CE of 3 November 1998 imposed to the surface
- waters devoted to the production of water for human consumption. Off. J. L 330, 32–54
- 553 (5/12/1998).
- 554 Fang, Q.X., Ma, L., Yu, Q., Hu, C.S., Li, X.X., Malone, R.W., 2013. Quantifying climate and
- 555 management effects on regional crop yield and nitrogen leaching in the North China Plain. J.
- 556 Environ. Qual. 42, 1466–1479.
- 557 Giola, P., Basso, B., Pruneddu, G., Giunta, F., Jones, J.W., 2012. Impact of manure and slurry
- applications on soil nitrate in a maize-triticale rotation: Field study and long term simulation
- analysis. Europ. J. Agronomy 38, 43–53.

- 560 Goss, M.J., Colbourn, P., Harris, G.L., Howse, K.R., 1988. Leaching of nitrogen under autumn-
- sown crops and the effects of tillage, in: Jenkinson, D.S., Smith, K.A. (Eds.), Nitrogen
- 562 Efficiency in Agricultural Soils. Elsevier Applied Science, Barking, Essex, UK, 269–282.
- 563 Grignani, C., Zavattaro, L., 2000. A survey on actual agricultural practices and their effects on
- the mineral nitrogen concentration of the soil solution. Eur. J. Agron. 12, 251–268.
- 565 Gutser, R., Ebertseder, T., Weber, A., Schraml, M., Schmidhalter, U., 2005. Short-term and
- 566 residual availability of nitrogen after long-term application of organic fertilizers on arable land. J
- 567 Plant Nutr Soil Sci 168:439–446. doi:10.1002/jpln.200520510.
- Hao, X., Chang, C., Travis, G.R., Zhang, F., 2003. Soil carbon and nitrogen response to 25 cattle
- 569 manure applications. J. Plant. Nutr. Soil Sci. 166, 239-245.
- 570 Hernández, D., Polo, A., Plaza, C., 2013. Long-term effects of pig slurry on barley yield and N
- 571 use efficiency under semiarid Mediterranean conditions. Eur. J. Agron. 44, 78-86.
- 572 Hooda, P.S., Edwards, A.C., Anderson, H.A., Miller, A., 2000. A review of water quality
- 573 concerns in livestock farming areas. Sci. Total Environ. 250, 143-167.
- 574 Ibrikci, H., Cetin, M., Karnez, E., Flügel, W.A., Tilkici, B., Bulbul, Y., Ryan, J., 2015.
- 575 Irrigation-induced nitrate losses assessed in a Mediterranean irrigation district. Agric. Water
- 576 Manage. 148, 223–231.
- 577 Indraratne, S.P., Hao, X., Chang, C., Godlinski, F., 2009. Rate of soil recovery following
- termination of long-term cattle manure applications. Geoderma 150, 415–423.
- Jabloun, M., Schelde, K., Tao, M.F., Olesena, J.E., 2015. Effect of temperature and precipitation
- 580 on nitrate leaching fromorganic cereal cropping systems in Denmark. Europ. J. Agronomy 62,
- 581 55-64.
- Jamali, H., Quayle, W.C., Baldock, J., 2015. Reducing nitrous oxide emissions and nitrogen
- 583 leaching losses from irrigated arable cropping in Australia through optimized irrigation
- scheduling. Agricultural and Forest Meteorology 208, 32–39.

- Le Gall, A., Legarto, J., Pflimlin, A., 1997. Place du mais et de la prairie dans les sistemes
- 586 fourragers laitiers. III-Incidence sur l'environment. Fourrages 150, 147–169.
- 587 Liang, X.Q., Li, H., He, M.M., Qian, Y.C., Liu, J., Nie, Z.Y., Ye, Y.S., Chen, Y.X., 2010.
- 588 Influence of N fertilization rates, rainfall, and temperature on nitrate leaching from a rainfed
- 589 winter wheat field in Taihu watershed. Phys. Chem. e Earth 36, 395-400.
- 590 Littell, R.C., Milliken, G.A., Stroup, W.W., Wolfinger, R.D., 1996. SAS System for Mixed
- 591 Models. Cary, NC: SAS Institute Inc.
- 592 Loague, K., Green, R.E., 1991. Statistical and graphical methods for evaluating solute transport
- 593 models: Overview and application. J. Contam. Hydrol. 7, 51–73.
- 594 López-Bellido, L., Munoz-Romero, V., Benítez-Vega, J., Fernández-García, P., Redondo, R.,
- 595 López-Bellido, R.J., 2012: Wheat response to nitrogen splitting applied to a Vertisol in different
- tillage systems and cropping rotations under typical Mediterranean climatic conditions. European
- 597 Journal of Agronomy 43, 24–32.
- 598 López-Bellido, L., López-Bellido, R.J., Redondo, R., 2005. Nitrogen efficiency in wheat under
- rainfed Mediterranean conditions as affected by spilt nitrogen application. Field Crop. Res. 94,
- 600 86–97.
- 601 Kayser, M., Benke, M., Isselstein, J., 2011. Little fertilizer response but high N loss risk of maize
- 602 growing on a productive organic-sandy soil. Agron Sustain Dev 31,709–718.
- 603 doi:10.1007/s13593-011-0046-9
- Kaur, T., Brar, B.S., Dhillon, N.S., 2008. Soil organic matter dynamics as affected by long-term
- 605 use of organic and inorganic fertilizers under maize-wheat cropping system. Nutr Cycl
- 606 Agroecosyst 81, 59–69.
- Kirda, C., Derici, M.R., Schepers, J.S., 2001. Yield response and N-fertiliser recovery of rainfed
 wheat growing in the Mediterranean region. Field Crop. Res. 71, 113–122.
- 609 Manca, A., 2009. L'applicazione della Direttiva Nitrati: ricadute e problematiche nella
- 610 zootecnia. Cagliari: Agenzia regionale per lo sviluppo in agricoltura.

- 611 Ministry of Agricultural, Food and Forestry Policies (Mipaaf), 2006. Osservatorio Nazionale
- 612 Pedologico e per la Qualità del Suolo Agricolo e Forestale Metodi di Analisi per i Fertilizzanti,
- 613 in: Trinchera, A., Leita L., Sequi P. (coordinators) Consiglio per la Ricerca e la sperimentazione
- 614 in Agricoltura, Istituto Sperimentale per la Nutrizione delle Piante Ed., Roma, Italy, 63-134.
- Montgomery, D.C., 1997. Design and Analysis of Experiments, th edition. New York, NY:
- 616 Wiley.
- 617 Moral, R., Moreno-Caselles, J., Perez-Murcia, M.D., Perez-Espinosa, A., Rufete, B., Paredes, C.,
- 618 2005. Characterisation of the organic matter pool in manures. Bioresource Technol. 96,
- **619** 153-158.
- 620 Morari, F., Lugato, E., Polese, R., Berti, A., Giardini, L., 2012. Nitrate concentrations in
- 621 groundwater under contrasting agricultural management practices in the low plains of Italy.
- 622 Agric. Ecosyst. Environ. 147, 47–56.
- 623 Nguyen, T.P.L., Seddaiu, G., Roggero, P.P., 2014. Hybrid knowledge for understanding complex
- 624 agri-environmental issues: nitrate pollution in Italy. Int. J. Agric. Sustain. 12, 164-182. doi:
- **625** 10.1080/14735903.2013.825995.
- 626 Perego, A., Basile, A., Bonfante, A., De Mascellis, R., Terribile, F., 2012. Nitrate leaching under
- 627 maize cropping systems in Po Valley (Italy). Agric. Ecosyst. Environ. 147, 57–65.
- 628 Phillips, C.J.C., 2010. Principles of Cattle Production. Landlinks. Press, Collingwood, Victoria.
- 629 Yu, Q.G., Chen, Y.X., Ye, X.Z., Tian, G.M., Zhang, Z.J, 2007. Influence of the DMPP (3,4-
- 630 dimethyl pyrazole phosphate) on nitrogen transformation and leaching in multi-layer soil
- 631 columns. Chemosphere 69, 825–831
- 632 Romano, N., Santini, A., 2002. The soil solution phase. Water retention and storage: Field water
- 633 capacity, in: Dane, J. H., Topp, C.G. (Eds.), Methods of Soil Analysis. Part 4 Physical Methods.
- 634 E-Publishing Inc., Madison, Wisconsin, USA, 723-725.

- 635 Sänger, A., Geisseler, D., Ludwig, B., 2010. Effects of rainfall pattern on carbon and nitrogen
- dynamics in soil amended with biogas slurry and composted cattle manure. J. Plant Nutr. Soil
- 637 Sci. 173, 692–698.
- 638 SAS Institute, 1999. SAS/STAT User's Guide, vol. 8. SAS Inst, Cary, NC.
- 639 Schröder, J.J., Jansen, A.G., Hilhorst, G.J., 2005. Long-term nitrogen supply from cattle slurry.
- 640 Soil Use Manag 21,196–204. doi:10.1111/j.1475-2743.2005.tb00125.x
- 641 Schröder, J.J., Dilz, K., 1987. Cattle slurry and farmyard manure as fertilizers for forage maize,
- 642 in: Van der Meer, H.G., Unwin, R.J., Van Dijk, T.A., Ennik, G.C. (Eds.), Animal Manure on
- 643 Grassland and Fodder Crops. Fertilizer or Waste. Developments in Plant Soil Scie. Martinus
- 644 Nijhoff Publishers, Dordrecht, 134–156.
- Shen, H., Brown, L.D., Zhi, H., 2006. Efficient estimation of log-normal means with application
 to pharmacokinetic data. Statist. Med. 25, 3023–3038.
- 647 Shepherd, M., Newell-Price, P., 2013. Manure management practices applied to a seven-course
- rotation on a sandy soil: effects on nitrate leaching. Soil Use Manage. 29, 210–219.
- 649 Silgram, M., Waring, R., Anthony, S., Webb, J., 2001. Intercomparasion of national & IPCC
- 650 methods for estimating N loss from agricultural land. Nutr. Cycl. Agroecosyst. 60, 189–195.
- 651 Simon, J.C., Le Corre, L., 1988. Lessivage d'azote en monoculture de mais, ensol granitique du
- 652 Finistere. Nitrogen leaching in maize monocolture, on granitic soil in Finistere. Fourrage, 114,
- **653** 193-207.
- 654 Singh, S.S., Ward, W.R., Hughes, J.W., Lautenbach, K., Murray, R.D., 1994. Behavior of Dairy-
- 655 Cows in a Straw Yard in Relation to Lameness. Veterinary Record 135, 251±3.
- 656 Smith, K.A., Chambers, B.J., 1993. Utilising the nitrogen content of organic manures on farms ±
- problems and practical solutions. Soil Use Manage. 9, 105-112.
- 658 Sørensen, P., 2004. Immobilisation, remineralisation and residual effects in subsequent crops of
- dairy cattle slurry nitrogen compared to mineral fertiliser nitrogen. Plant Soil 267, 285–296.

- 660 Trindade, H., Coutinho, J., Jarvis, S., Moreira, N., 2009. Effects of different rates and timing of
- application of nitrogen as slurry and mineral fertilizer on yield of herbage and nitrate-leaching
 potential of a maize/Italian ryegrass cropping system in north-west Portugal. Grass Forage Sci.
 663 64, 2–11.
- Trindade, H., Coutinho, J., Van Beusichem, M.L., Scholefield, D., Moreira, N., 1997. Nitrate
- leaching from sandy loam soil under a double-cropping forage system estimate from suction-
- probe measurements. Plant soil 195, 247-256.
- 667 US Department of Agriculture (USDA), 2006. Natural resources conservation service. Keys to
- soil taxonomy, 10th ed. Washington, DC, USA.
- 669 Vertes, F., Decau, M.L., 1992. Suivis d'azote minéral dans les sols: risque de lessivage de nitrate
- 670 selon le couvert végétal. Fourrages 129, 11–28.
- 671 Wenz, J.R., Jensen, S.M., Lombard, J.E., Wagner, B.A., Dinsmore, R.P., 2007. Herd
- 672 Management Practices and Their Association with Bulk Tank Somatic Cell Count on United
- 673 States Dairy Operations. J. Dairy Sci. 90, 3652–3659. doi:10.3168/jds.2006-592.
- 674 Williams, J.R., 1995. The EPIC model. In: Singh VP (ed) Computer models of watershed
- hydrology. Water Resources Publications, Highlands Ranch, 909–1000.
- 676 Worrall, F., Spencer, E., Burt, T.P., 2009. The effectiveness of nitrate vulnerable zones for
- 677 limiting surface water nitrate concentrations. J. Hydrol 370, 21–28.
- 678 Zavattaro, L., Monaco, S., Sacco, D., Grignani, C., 2012. Options to reduce N loss from maize in
- 679 intensive cropping systems in Northern Italy. Agr. Ecosyst. Environ. 147, 24–35.

Figure 1. Water balance dynamic from May 2009 to June 2012 under Maize-Italian Ryegrasssystem.

683

Figure 2. Dynamics of the NO_3^- concentration (mg L⁻¹, transformed by Shen et al., 2006) in the 684 soil solution and soil water percolation as simulated by EPIC model during the silage maize -685 Italian ryegrass rotation (2009–2010) for the different treatments (MA = Cattle Manure; SL =686 Cattle Slurry; SM = Cattle Slurry+ Mineral; MI = Mineral). Vertical black arrows indicate the 687 688 dates of fertilizers distribution. The horizontal black line indicates the legal threshold of 50 mg L^{-1} for drinkable water and the rectangles indicate the duration of cropping cycle. The asterisks 689 690 indicate the sampling dates when the NO₃⁻ concentration significantly exceeded 50 mg L^{-1} . 691 Figure 3. Dynamics of the NO₃⁻ concentration (mg L^{-1} , transformed by Shen et al., 2006) in the 692 693 soil solution and soil water percolation as simulated by EPIC model during the silage maize -694 Italian ryegrass rotation (2010–2011) for the different treatments (MA = Cattle Manure; SL =

695 Cattle Slurry; SM = Cattle Slurry+ Mineral; MI = Mineral). Vertical black arrows indicate the 696 dates of fertilizers distribution. The horizontal black line indicates the legal threshold of 50 mg 697 L^{-1} for drinkable water and the rectangles indicate the duration of cropping cycle. The asterisks 698 indicate the sampling dates when the NO₃⁻ concentration significantly exceeded 50 mg L^{-1} . 699

Figure 4. Dynamics of the NO_3^- concentration (mg L⁻¹, transformed by Shen et al., 2006) in the soil solution and soil water percolation as simulated by EPIC model during the silage maize – Italian ryegrass rotation (2011–2012) for the different treatments (MA = Cattle Manure; SL = Cattle Slurry; SM = Cattle Slurry+ Mineral; MI = Mineral). Vertical black arrows indicate the dates of fertilizers distribution. The horizontal black dotted line indicates the legal threshold of 50 mg L⁻¹ for drinkable water and the rectangles indicate the duration of cropping cycle. The

- asterisks indicate the sampling dates when the NO_3 concentration significantly exceeded 50 mg
- **707** L^{-1} .
- Figure 5. Distribution of cumulated N leaching (kg N ha⁻¹) from June 2009 to May 2012 for the
- 709 different treatments.

Table 1. Physical and chemical characteristics of the soil at the beginning of the experiment

711	(2009).
-----	---------

	Soil depth (cm)				
Trait	0-45	46-70	71-100	>100	
Clay, g kg ⁻¹	25 ±0.16	20 ± 0.15	28 ± 0.48	66 ± 1.04	
Sand, g kg ⁻¹	953 ± 0.20	940 ± 1.44	954 ± 0.47	912 ± 1.01	
Silt, g kg ⁻¹	21 ± 0.14	19 ± 0.13	22 ± 0.22	22 ± 0.31	
Bulk density, g cm ⁻³	1.55 ± 0.02	1.35 ± 0.11	1.22 ± 0.17	1.25 ± 0.18	
Water holding capacity, %Vol 0 kPa	41.6	49.2	52.5	52.3	
Field capacity, %Vol 33kPa	6.6	7.8	8.8	9.7	
Field capacity, %Vol 23kPa	17.7	20.5	22.3	23.2	
Wilting point, %Vol 1500 kPa	3.8	3.2	3.8	3.8	
Soil Organic Matter, g kg ⁻¹	23.4 ± 0.10	8.0 ± 0.19	4.7 ± 0.14	4.1 ± 0.10	
Tot N, g kg ⁻¹	1.3 ± 0.04	0.4 ± 0.08	0.3 ± 0.05	0.4 ± 0.07	

Table 2. N inputs (kg N ha⁻¹) from organic and mineral fertilizers (three-years average \pm std

rror) for Italian ryegrass and maize (MA = Cattle Manure; SL = Cattle Slurry; SM = Cattle

714 Slurry + Mineral; MI = Mineral).

	Italian ryegrass	Maize
Treatments	2009-12	2009-2011
MA	151 ±24	379 ±38
SL	317 ±115	469 ±213
SM	231 ±53	374 ±60
MI	130 ±0	316 ±0

715

Table 3. Indices used for the evaluation of the model performances (RRMSE= relative root mean

square error; EF=modeling efficiency) for calibration and validation in Italian ryegrass, maize

718 and ground water table depth.

		RRMSE	EF	Slope	Intercept	\mathbf{R}^2
	Best	0.0	1.0	1.0	0.0	1.0
	Min	0.0	$-\infty$	$-\infty$	$\infty - \infty$	$-\infty$
	Max	$+\infty$	1.0	$+\infty$	$\infty + \infty$	$\infty + \infty$
Silage maize	Calibration	5.90	0.50	0.70	6.95	0.57
yield	Validation	8.02	0.24	0.46	12.00	0.31
It. ryegrass	Calibration	17.55	0.55	0.65	2.68	0.64
yield	Validation	19.40	0.25	0.75	2.21	0.57
Groundwater	Calibration	25.75	0.37	1.13	-0.01	0.72
table depth	Validation	19.50	0.54	0.89	0.17	0.67

719

720

721

723 Table 4. Average three years N balance estimated from N input, removal and leaching as influenced by the treatments (MA = Cattle Manure; SL =

724 Cattle Slurry; SM = Cattle Slurry+ Mineral; MI = Mineral). The residual N is calculated by difference. Means followed by the same letter were not

	N input		N removal		N leaching		Residual N	
Treatments	(kg N ha ⁻¹ ± err std)	(%)	(kg N ha ⁻¹)	(%)	(kg N ha ⁻¹)	(%)	(kg N ha ⁻¹ ± err std)	(%)
MA	530 ± 64	100	216 b	41	42 b	8	272 ± 41 c	51
SL	813 ± 249	100	255 ab	31	110 a	14	447 ± 246 a	55
SM	602 ± 88	100	281 a	47	94 a	16	$226\pm95~b$	38
MI	446 ± 0	100	301 a	68	89 a	20	56 ± 44 c	12
Mean	599	100	264	47	84	15	250	39
CV (%)	20	-	14	-	29	-	83	-









