

Melatonin treatment in winter and spring and reproductive recovery in Sarda breed sheep

Questa è la versione Post print del seguente articolo:

Original

Melatonin treatment in winter and spring and reproductive recovery in Sarda breed sheep / Mura, Maria Consuelo; Luridiana, Sebastiano; Farci, Federico; DI STEFANO, Maria Veronica; Daga, Cinzia; Pulinas, Luisa; Starià, Joàñe; Carcangiu, Vincenzo. - In: ANIMAL REPRODUCTION SCIENCE. - ISSN 0378-4320. - 185:(2017), pp. 104-108. [10.1016/j.anireprosci.2017.08.009]

Availability:

This version is available at: 11388/181771 since: 2022-05-25T13:14:32Z

Publisher:

Published

DOI:10.1016/j.anireprosci.2017.08.009

Terms of use:

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

Publisher copyright

note finali coverpage

(Article begins on next page)

1 **Melatonin treatment in winter and spring and reproductive recovery in Sarda breed sheep**

2

3 Maria Consuelo Mura^a, Sebastiano Luridiana^a, Federico Farci^a, Maria Veronica Di Stefano^a, Cinzia
4 Daga^a, Luisa Pulinas^a, Jože Starič^b, Vincenzo Carcangiu^{a*}

5 ^aDipartimento di Medicina Veterinaria, Università di Sassari, Via Vienna 2, 07100, Sassari, Italia

6 ^bClinic for reproduction and large animals - section for ruminants, Veterinary Faculty, University of
7 Ljubljana, Cesta v Mestni log 47, 1000 Ljubljana, Slovenia

8

9 *Corresponding author: Vincenzo Carcangiu, Dipartimento di Medicina Veterinaria, Università di
10 Sassari, Via Vienna 2, 07100, Sassari, Italia, Tel. +39079229436; Fax +39079229592, E-mail
11 address: endvet@uniss.it,

12

14 **Abstract**

15 This study aimed to evaluate the effect of melatonin treatment carried out between late
16 winter and early spring on reproductive response in Sarda breed sheep and whether the photo-
17 refractoriness can influence this reproductive response. The study was conducted on 3,200 adult
18 ewes, aged 3 to 6 years old, with body condition score (BCS) 2.5-4.0, from 16 commercial sheep
19 farms in Northern Sardinia. In each farm 200 animals were enrolled and subdivided into 2 groups (n
20 = 100 each): Group M (treated with one 18 mg melatonin implant), and group C (untreated). The
21 melatonin treatments were performed on 10th February; 10th March; 10th April and on 10th May each
22 time in 4 different randomly selected farms. Adult males, treated with 3 melatonin implants 1 week
23 before females, were introduced in each flock 35 days after ewes' treatment, and removed after 45
24 days of cohabitation with females. Pregnancy was determined by transabdominal ultrasonography
25 examination between 45th and 90th day after ram introduction. Data showed that melatonin
26 treatment increased the fertility rate significantly ($P<0.05$), with the higher fertility rate in the ewes
27 treated in April and May. The average time in days from male introduction to lambing was shorter
28 in treated than in control ewes ($P<0.05$). Further, at the 160th and 170th day after male introduction
29 the group M showed a higher number of lambed ewes compared to C ($P<0.01$). This effect was
30 observed at 180th and 190th days after ram introduction, also, but with lower significance ($P<0.05$).
31 In conclusion, melatonin treatment improved reproductive efficiency and advanced breeding season
32 in Sarda sheep, especially when ewes were treated in spring.

33

34 Keywords: Sarda sheep, seasonal reproduction, melatonin implant, fertility

35

36 **Introduction**

37 Reproductive seasonality of dairy sheep at the Mediterranean basin latitudes is a major
38 constraint for production (Chemineau et al., 2007). Furthermore, these areas have a mild climate
39 during winter and autumn, rainfall mainly concentrated in spring and a very dry summer. Thus, the

40 sheep production cycle is closely linked to pasture growth which occurs predominantly in autumn
41 and spring (Carcangiu et al., 2011). Therefore, to increase the length of lactation (6-7 months), it is
42 necessary to mate sheep in spring to obtain lambing in autumn. The timing of reproductive
43 seasonality in many small ruminant species is controlled by the shortening of photoperiod
44 (Chemineau et al., 1992) i.e., while the long photoperiods inhibit, the short photoperiods stimulate
45 the reproductive activity (Bittman and Karsh 1984; Boden and Kennaway, 2006). Melatonin shows
46 a low blood-concentration during daylight and a high concentration during darkness, thus it can be
47 considered as an organic informer of the annual photoperiodic trend (Carcangiu et al., 2005,
48 Carcangiu et al., 2013). Melatonin subcutaneous implants are used in Europe and other parts of the
49 world, to advance reproductive activity of sheep in the spring (Abecia et al., 2006; Papachristoforou
50 et al., 2007; deNicolo et al., 2008; Carcangiu et al., 2012). At high latitudes, the administration of
51 melatonin by slow-release implants is usually performed around the summer solstice (Haresign et
52 al., 1990), whereas, in the Mediterranean areas, around the spring equinox (Chemineau et al., 1996;
53 Forcada et al., 1999). Several experiments have been carried out to determine which is the best time
54 for treatment with melatonin, but the responses vary with breed (Staples et al., 1992; Abecia et al.,
55 2007). The Sarda sheep (more than 3.0 million ewes farmed in Sardinia) is the main dairy breed in
56 Italy and its production is parallels the pasture growth pattern (Carta et al., 2009). It exhibits a short
57 anoestrous period, generally from early February until late March (personal observation).
58 Administration of melatonin implants has been reported to improved fertility and advance lambing
59 (Carcangiu et al., 2012; Luridiana et al., 2015). However, even in Sarda sheep the reproductive
60 responses after treatment with melatonin are sometimes unsatisfactory, probably due to treatment
61 fallen in the period of refractoriness (Abecia et al., 2007; Sweeney et al., 1997). This study aimed to
62 evaluate the effect of melatonin treatment carried out between late winter and early spring on
63 reproductive response in Sarda breed sheep and to determine whether photorefractoriness influences
64 this response.

66 **2. Materials and Methods**

67 **2.1 Experimental design**

68 This study was conducted on 3,200 Sarda breed ewes from 16 farms, located in North
69 Sardinia (40°N). In order to minimize the farm effect, farms were selected according to comparable
70 features such as total number of animals, milk production, flock health, veterinary
71 recommendations (deworming, vaccinations and use of other pharmacological treatments), nutrition
72 and reproductive management. All farms were located in the same area, within a 50 km radius of
73 Sassari. During the day the animals grazed on legumes and grasses and they also received 300 g of
74 commercial concentrate feed per head (crude protein 20.4% and 12.5 MJ ME/kg DM) during
75 milking. The sheep were penned during night, and received hay (crude protein 11.1% and 7.2 MJ
76 ME/kg DM) and water *ad libitum*. Each farm reared approximately 400 sheep, and 200 clinically
77 healthy ewes were chosen from each farm for the study, accounting for a total of 3,200 head. Ewes
78 enrolled in the study were lactating, 3 to 6 years old, with body condition score (BCS) 2.5-4.0 and
79 with a single lamb born between 1st November and 10th December, 2013. The individual rumen
80 bolus number was recorded, and each ewe was individually marked with a numbered collar to avoid
81 mistakes in the animals' identification.

82

83 **2.2 Treatment and registration data**

84 On each farm the enrolled animals were randomly subdivided into 2 groups, each containing
85 100 ewes. The ewes in group M received one melatonin implant (18 mg) (Melovine, CEVA Sanità
86 Animale, Agrate Brianza, Milano) in the left retro-auricular area; the other group of ewes served as
87 control and remained untreated (group C). In the first 4 randomly selected farms melatonin implants
88 were administered on 10th February, in the second 4 on 10th March, in the third 4 on 10th April and
89 in the last 4 on 10th May; controls and treated animals were kept together all the time. Adult rams
90 (male/female ratio 1/20) were introduced on each farm 35 days after treatment of ewes and
91 remained in the flock for 45 days. Before ram introduction, ewes were separated from rams for at

92 least 3 months. Rams received 3 melatonin implants, in the left retro-auricular area, 1 week before
93 ewes. Gestation was diagnosed starting from 45 days and up to 90 days after the rams introduction
94 by transabdominal ultrasonography examination using Esaote Piemedical Tringa linear equipment
95 (Esaote Europe B.V., Maastricht, The Netherlands) provided with a 5.0 – 7.5 MHz multiple
96 frequency linear probe. For each ewe the lambing date and number of new-born lambs were
97 recorded from 150th to 190th day after ram introduction.

98

99 **2.3 Statistical analysis**

100 All statistical analysis was performed using the computing environment R, version 3.3.2 (R Core
101 Team, 2016). A logit-link Hierarchical Linear Model (HLM) appropriate for binomial (lambing/not
102 lambing) data was used to analyse the fertility of different treatment time. Variables considered
103 were treatment and treatment time. To analyse the effect of treatment time on period in days from
104 male introduction to lambing a HLM procedure according to the following model was performed.

$$105 Y_{ijk} = \mu + T_m(Pe)_{ij} + \varepsilon_{ijk}$$

106 where Y_{ijk} is the period in days from male introduction to lambing, μ is the overall mean, T_m is the
107 fixed effect of treatment, Pe is the nested time effect within treatment, and ε_{ijk} is the error effect.

108 The same model was used to analyse the litter size. A P value <0.05 was considered statistically
109 significant. Multiple comparisons of the means were performed using Tukey's method.

110

111 **3. Results**

112 The number of ewes diagnosed as pregnant differed from the number of lambed ewes of
113 about 3%. Melatonin treatment increased the fertility rate significantly in the four treatment periods
114 compared to control group ($P < 0.05$). The higher fertility rate was seen in the ewes treated in April
115 and May compared to those treated in the other two periods ($P < 0.05$) (Table 1). The average time in
116 days from male introduction to lambing was shorter in the treated animals compared to controls, in
117 all the experimental periods (Table1). Melatonin treatment did not affect litter size, and no

118 difference was observed between the treated groups and controls in different periods of observation.
119 By daily observation of lambing trend, melatonin treated ewes lambbed approximately ten days
120 earlier compared to controls. At the 160th and 170th day from male introduction, in all four treatment
121 periods the treated group showed a higher number of lambbed ewes compared to control ($P<0.01$).
122 At the 180th and 190th day from male introduction the treated ewes exhibited a higher number of
123 lambbed ewes compared to controls ($P<0.05$), although the difference between treated and control
124 group was smaller than in the first 20 days of lambing. In all four treatment times, the lambing peak
125 was registered at day 170, in treated ewes, and at day 180 from male introduction, in controls (Table
126 2).

127

128 **4. Discussion**

129 The results from this study show that melatonin was able to advance the reproductive
130 activity in a dairy sheep. This finding is in agreement with other observations where melatonin
131 improves the reproductive efficiency in different sheep breeds (Chemineau et al., 1991; Haresign,
132 1992; Staples et al., 1992; Carcangiu et al., 2012). However, in many experiments melatonin
133 treatments were performed from the end of April till September, and thus are not completely
134 comparable to our experiment (Haresign et al., 1990; Forcada et al., 1995; Scott et al., 2009).

135 However, our data indicate that the reproductive response is influenced by the treatment
136 time; in fact, fertility was higher when the animals were implanted in April and May (early spring)
137 than in February-March (late winter). This result does not agree with the findings of Abecia et al.
138 (2007) in three sheep breeds reared in Spain, where fertility of the Rasa Aragonesa breed was not
139 improved when the administration of melatonin was performed in January and February (winter)
140 compared to April and May (spring). This finding from Abecia et al. (2007) is also in agreement
141 with previous study carried out by Forcada et al. (2002) and Zuniga et al. (2002) always on the
142 Rasa Aragonesa breed .

143 In the Assaf breed melatonin administration in January, February (winter) and May (spring)
144 increased fertility, but not when it was administered in April (spring). In the same experiment the
145 administration of melatonin in the Merino breed in February (winter) and April (spring) led to
146 increased fertility but not in January (winter) and May (spring) (Abecia et al., 2007). This finding is
147 presumably due to the different sensitivity to photoperiod in the different sheep breeds. Indeed, after
148 a short-day period, in several sheep breeds, a photo-refractory period occurs, corresponding to the
149 deep anestrous that can have a different length among sheep breeds (Chemineau et al., 2010).
150 Therefore the breed differences in response to melatonin and the male effect will impact on
151 reproductive. Our results show that in the four treatment times, reproductive response was different,
152 the more effective treatments being those performed in spring, compared to those performed in
153 winter.

154 The onset of the breeding season is a consequence of the number of long days experienced
155 by the ewes (Malpaux et al., 1989), and it is possible that our Sarda sheep experienced a number of
156 long or increasing days sufficient to respond to a short-day treatment such as that resulting from
157 melatonin implants (O'Callaghan et al., 1994; Malpaux et al., 1997). Sweeney et al. (1997)
158 postulated that ewes are able to reinitiate reproductive activity in response to 35 long days
159 (photorefractoriness period) followed by short days or melatonin treatment at any stage between the
160 winter and summer solstices. In fact in the present study the first melatonin treatment was
161 performed on February 10, thus approximately 50 days after the winter solstice. It is therefore
162 plausible to think that the ewes were already at a late-stage of their photorefractoriness period. It is
163 reasonable to think that the progressive increase in the fertility rate in the other three doses was due
164 to an increasing sensitivity of the reproductive system to melatonin.

165 In different experiments the reproductive response to the male effect in melatonin treated
166 ewes was affected by the farm/flock as well (Chemineau et al., 1991; Haresign, 1992; Scott et al.,
167 2009).

168 The differences in fertility rate found among treated and control ewes are mainly mediated
169 by the direct effect of melatonin on the ovary (where are located melatonin receptors) (Yie et al.,
170 1995), as well as by its effect on gonadotropin secretion at pituitary-hypothalamic level (Viguié et
171 al., 1995). Moreover, in all four treatment times, implanted sheep exhibited a reduced number of
172 days from ram introduction to lambing than the controls. Subsequently, the treated animals had a
173 more ready response to the male effect compared to controls. In our experiment a difference
174 between the melatonin-treated and untreated groups was registered in the lambing periods. That
175 difference was attributable to the fact that treated ewes were mated within the first 17 days of the
176 mating period, and had a higher proportion of cyclic ewes at ram introduction than the untreated
177 group. Then the administration of melatonin was able to stimulate the resumption of reproductive
178 activity, and this result agrees with the findings from Abecia et al. (2006). Lambing records showed
179 that the melatonin treated animals exhibited the onset of reproductive activity approximately one
180 week before the untreated ones. In fact, at the 160th and 170th day after male introduction the
181 number of lambed ewes was much higher in treated than in control animals. Conversely, at the 180th
182 and 190th day from male introduction the number of lambed ewes increased considerably in control
183 compared to treated group. This trend probably is related to the "carry-over" effect, that occur when
184 the treated and control animals are kept together (Abecia et al., 2006). When melatonin treated and
185 untreated ewes are kept together the onset of reproductive activity occurred earlier in control ewes
186 as well compared to isolated controls (Kennaway et al. 1987). Moreover, the introduction of estrus
187 ewes, exposed to controlled photoperiod, to a flock of anestrus ewes triggers reproductive activity,
188 indicating that female—female social interactions can influence the timing of reproductive
189 transitions in ewes (O’Callaghan et al., 1994). Establishing the number of sheep in the flock to be
190 treated to obtain the triggering effect on reproductive activity could be of considerable importance
191 for farmers, since the treatment of a part of the flock is able to improve the reproductive efficiency
192 of the entire flock. However, even without an economic calculation, it is clear that in the present
193 research, the treated ewes had an advance in the onset of lactation compared to controls, and this

194 results in an economic profit for farmers. In fact, the treated animals exhibited on average 10 days
195 shorter time between male introduction and lambing than controls.

196 The same litter size was observed in treated and control animals in all four treatment times
197 in this study, as established also in our previous works (Carcangiu et al., 2012; Luridiana et al.,
198 2015). Our data did not agree with Abecia et al. (2002) and Scott et al. (2009), who found an
199 increase in litter size in the treated animals. Some other studies found no effect of melatonin
200 treatment on litter size (Rajkumar et al., 1989; Schoeman and Botha, 1995; Gates et al., 1998).
201 Moreover, Abecia et al. (2007) after melatonin implants found an increased litter size in Rasa
202 Aragonesa, but not in Assaf breed. Thus, since Sarda and Assaf are both dairy breeds, we can
203 hypothesize that the absence of melatonin effect on litter size could be due to the genetic features of
204 these breeds.

205

206 **5. Conclusion**

207 Melatonin treatment improved the reproductive efficiency and advanced the breeding season
208 in dairy sheep. These effects were more pronounced when ewes were treated in spring than in
209 winter. So, to obtain an optimal reproductive efficiency in Sarda ewes melatonin implants should be
210 placed after the spring solstice. The present data also indicate that the period of photorefractoriness
211 in this breed could be confined between January and early February since the melatonin
212 administration on February 10 is able to stimulate the reproductive activity.

213

214 **Acknowledgments**

215 This research was supported by grants from the RAS research project entitled
216 MIGLIOVIGENSAR and by research project entitled "Study of different methods for seasonal
217 adjustment of sheep in Sardinia" developed at the University of Sassari, by way of the Temporary
218 Researcher Contract financed with resources from the P.O.R: SARDEGNA F.S.E. 2007-2013 -

219 Objective Regional Competitiveness and Employment, Asse IV Capitale umano, Linea di Attività
220 1.3.1.

221

222 **Conflict of interest**

223 None of the authors have any conflicts of interest to declare.

224

225 **References**

226 Abecia, J.A., Forcada, F., Zuniga, O., 2002. The effect of melatonin on the secretion of
227 progesterone in sheep and on the development of ovine embryos in vitro. *Vet. Res.*
228 *Commun.* 26, 51-58.

229 Abecia, J.A., Palacín, I., Forcada, F., Valares, J.A., 2006. The effect of melatonin treatment on the
230 ovarian response of ewes to the ram effect. *Domest. Anim. Endocrinol.* 31, 52-62.

231 Abecia, J.A., Valares, J.A., Forcada, F., Palacín, I., Martín, S., Martino, A., 2007. The effect of
232 melatonin on the reproductive performance of three sheep breeds in Spain. *Small Rum. Res.*
233 69, 10-16.

234 Bittman, E.L., Karsch, F.J., 1984. Nightly duration of pineal melatonin secretion determines the
235 reproductive response to inhibitory daylength in the ewe. *Biol. Reprod.* 30, 585–593.

236 Boden, M.J., Kennaway, D.J., 2006. Circadian rhythms and reproduction. *Reproduction* 132, 379-
237 392.

238 Carcangiu, V., Mura, M.C., Bini, P.P., Vacca, G.M., Daga, C., Luridiana, S., 2012. Can advance of
239 first lambing induced by melatonin implants influence the next lambing time in Sarda breed
240 sheep? *Can. J. Anim. Sci.* 92, 67-71.

241 Carcangiu, V., Mura, M.C., Parmeggiani, A., Piccione, G., Bini, P.P., Cosso, G., Luridiana, S.,
242 2013. Daily rhythm of blood melatonin concentrations in sheep of different ages. *Biol.*
243 *Rhythm Res.* 44, 908-915.

244 Carcangiu, V., Luridiana, S., Vacca, G.M., Daga, C., Mura, M.C., 2011. A polymorphism at the
245 melatonin receptor 1A (MTNR1Aa) gene in Sarda ewes affects fertility after AI in the
246 spring. *Reprod. Fert. Dev.* 23, 376-380.

247 Carcangiu, V., Vacca, G.M., Parmeggiani, A., Mura, M.C., Bini, P.P., 2005. Blood melatonin levels
248 relating to the reproductive activity of Sarda does. *Small Ruminant Res.* 59, 7-13.

249 Carta, A., Casu, S., Salaris, S., 2009. Invited review: current state of genetic improvement in dairy
250 sheep. *J. Dairy Sci.* 92, 5814-5833.

251 Chemineau, P., Guillaume, D., Migaud, M., Thiéry, J.C., Pellicer-Rubio, M.T., Malpoux, B., 2008.
252 Seasonality of reproduction in mammals: Intimate regulatory mechanisms and practical
253 implications. *Reprod. Dom. Anim.* 43 (Suppl. 2), 40-47.

254 Chemineau, P., Malpoux, B., Delgadillo, J.A., Guerin, Y., Ravault, J.P., Thimonier, J., Pelletier, J.,
255 1992. Control of sheep and goat reproduction: use of light and melatonin. *Anim. Reprod.*
256 *Sci.*, 30, 157-184.

257 Chemineau, P., Bodin, L., Migaud, M., Thiery, J.C., Malpoux, B., 2010. Neuroendocrine and
258 Genetic Control of Seasonal Reproduction in Sheep and Goats. *Reprod. Dom. Anim.* 45
259 (Suppl. 3), 42-49.

260 Chemineau, P., Malpoux, B., Brillard, J. P., Fostier, A., 2007. Seasonality of reproduction and
261 production in farm fishes, birds and mammals. *Animal* 1, 419-432.

262 Chemineau, P., Malpoux, B., Pelletier, J., Leboeuf, B., Delgadillo, J.A., Deletang, F., Pobel, T.,
263 Brice, G., 1996. Emploi des implants de melatonine et des traitements photoperiodiques
264 pour maitriser la reproduction saisonniere chez les ovins et les caprins (use of melatonin
265 implants and photoperiodic treatments to resuce seasonal reproduction in sheep and goats).
266 *INRA Prod. Anim.* 9, 45-60.

267 Chemineau, P., Vandaele, E., Brice, G., Jardon, C., 1991. Utilisation des implants de mélatonine
268 pour l'amélioration des performances de reproduction chez la brebis (use of melatonin
269 implants to improve sheep reproductive performances). *Rec. Méd. Vét.* 167, 227-239.

270 deNicolo, G., Morris, S.T., Kenyon, P.R., Morel, P.C., Parkinson, T.J., 2008. Melatonin-improved
271 reproductive performance in sheep bred out of season. *Anim. Reprod. Sci.* 109, 124-133.

272 Forcada, F., Abecia, J.A., Lozano, J.M., Ferrer, L.M., Lacasta, D., 1999. The effects on
273 reproductive performance in the short and medium term of the combined use of exogenous
274 melatonin and progestagen pessaries in ewes with a short seasonal anoestrous period. *Vet.*
275 *Res. Commun.* 23, 257-263.

276 Forcada, F., Abecia, J.A., Zuniga, O., Lozano, J.M., 2002. Variation in the ability of melatonin
277 implants inserted at two different times after the winter solstice to restore reproductive
278 activity in reduced seasonality ewes. *Aust. J. Agric. Res.* 53, 167–173.

279 Forcada, F., Zarazaga, L., Abecia, J.A., 1995. Effect of exogenous melatonin and plane of nutrition
280 after weaning on estrous activity, endocrine status and ovulation rate in Salz ewes lambing in
281 the seasonal anoestrus. *Theriogenology* 43, 1179–1193.

282 Gates, P.J., Henningson, T., Tengroth, G., Forsberg, M., 1998. Effects of melatonin, progestagens,
283 and the ram on out-of-season reproduction in Swedish Landrace finewool sheep. *Acta Vet.*
284 *Scand.* 39, 499-510.

285 Goldman, B.D., 2001. Mammalian photoperiodic system: formal properties and neuroendocrine
286 mechanisms of photoperiodic time measurement. *J. Biol. Rhythms* 16, 283-301.

287 Haresign, W., 1992. The effect of implantation of lowland ewes with melatonin on the time of
288 mating and reproductive performance. *Anim. Prod.* 54, 31–39.

289 Haresign, W., Peters, A.R., Staples, L.D., 1990. The effect of melatonin implants on breeding
290 activity and litter size in commercial sheep flocks in the UK. *Anim. Prod.* 50, 111–121.

291 Kennaway, D.J., Dunstan, E.A., Staples, L.D., 1987. Photoperiodic control of the onset of breeding
292 activity and fecundity in ewes *J. Reprod. Fert. (Suppl.)* 34, 187-199.

293 Luridiana, S., Mura, M.C., Daga, C., Farci, F., Di Stefano, M.V., Zidda, F., Carcangiu, V., 2015.
294 Melatonin treatment in spring and reproductive recovery in sheep with different body
295 condition score and age. *Anim. Reprod. Sci.* 160, 68-73.

296 Malpaux, B., Robinson, J.E., Wayne, N.L., Karsch, F.J., 1989. Regulation of the onset of the
297 breeding season of the ewe: importance of long days and of an endogenous reproductive
298 rhythm. *J. Endocrinol.* 122, 269–278.

299 Malpaux, B., Viguié, C., Skinner, D.C., Thyéry, J.C., Chemineau, P., 1997. Control of the
300 circannual rhythm of reproduction by melatonin in the ewe. *Brain Res. Bull.* 44, 431–438.

301 O’Callaghan, D., Donovan, A., Sunderland, S.J., Boland, M.P., Roche, J.F., 1994. Effect of the
302 presence of male and female flockmates on reproductive activity in ewes. *J. Reprod. Fertil.*
303 100, 497-503.

304 Papachristoforou, C., Koumas, A., Photiou, C., 2007. Initiation of the breeding season in ewe lambs
305 and goat kids with melatonin implants. *Small Ruminant Res.* 73, 122-126.

306 Rajkumar, R.R., Argo, C.M., Rodway, R.G., 1989. Fertility of ewes given either melatonin or
307 progesterone sponges. *Vet. Rec.* 124, 215-217.

308 Schoeman, S.J., Botha, M.A., 1995. The effect of melatonin treatment on reproductive efficiency in
309 an accelerated lambing system. *J. S. Afr. Vet. Assoc.* 66, 230-234.

310 Scott, P.R., Sargison, N.D., Macrae, A.I., Gough, M.R., 2009. Melatonin treatment prior to the
311 normal breeding season increases fetal number in United Kingdom sheep flocks. *Vet. J.* 182,
312 198-202.

313 Staples, L.D., McPhee, S., Kennaway, D.J., Williams, A.H., 1992. The influence of exogenous
314 melatonin on the seasonal patterns of ovulation and oestrus in sheep. *Anim. Reprod. Sci.* 30,
315 185–223.

316 Sweeney, T., Donovan, A., Roche, J.F., O’Callaghan, D., 1997. Variation in the ability of a long
317 day followed by a short-day photoperiod signal to initiate reproductive activity in ewes at
318 different times of the year. *J. Reprod. Fertil.* 109, 121–127.

319 Viguié, C., Caraty, A., Locatelli, A., Malpaux, B., 1995. Regulation of luteinizing hormone-
320 releasing hormone (LHRH) secretion by melatonin in the ewe: II. Changes in N-Methyl-

321 D,L-aspartic acid induced LHRH release during the stimulation of LH secretion by short
322 days. *Biol. Reprod.* 52, 1156-1161.

323 Yie, S.M., Niles, L.P., Younglai, E.V., 1995. Melatonin receptors on human granulosa cell
324 membranes. *J. Clin. Endocrinol. Metab.* 80, 1747-1749.

325 Zuniga, O., Forcada, F., Abecia, J.A., 2002. The effect of melatonin implants on the response to the
326 male effect and on the subsequent cyclicity of Rasa Aragonesa ewes implanted in April.
327 *Anim. Reprod. Sci.* 15, 165–174.

Table 1

Fertility rate, litter size and distance in days from rams introduction to parturition (DRIP) in the treated (M) and control (C) ewes in the four period (total ewes n = 3,200).

	M	C		M	C	M	C	
Time	Fertility rate	Fertility rate		Litter size	Litter size	DRIP	DRIP	
10 th Febr.	68% ^a	61% ^a	*	1.29	1.18	164.96	175.57	*
10 th March	75% ^a	66% ^a	*	1.20	1.22	165.66	174.77	*
10 th April	81% ^b	72% ^b	*	1.18	1.17	165.62	175.07	*
10 th May	83% ^b	76% ^b	*	1.19	1.22	165.90	175.02	*

Means within a column with different lowercase are significantly different (P<0.05); In group comparison means within a row with * differ significantly for P<0.05.

Table 2

Cumulative number of lambed ewes from 150th to 190th day after rams introduction (every 10 days) in the treated (M) and control (C) Sarda breed sheep, in the four observed periods (total ewes n=3,200).

	150-160 DARI ^a			161-170 DARI ^a			171-180 DARI ^a			181-190 DARI ^a		
	M	C		M	C		M	C		M	C	
10 th February	83	17	**	213	57	**	253	177	*	272	244	*
10 th March	77	21	**	230	68	**	276	201	*	300	264	*
10 th April	83	22	**	246	75	**	217	226	*	324	288	*
10 th May	83	20	**	248	78	**	302	231	*	332	304	*

^aDARI = days from rams introduction to parturition. In group comparison means within a row with

* or ** differ significantly for P<0.05 and for P<0.01 respectively.

Conflict of interest

None of the authors have any conflicts of interest to declare.