The weather dictates the rhythms: Alpine chamois activity is well adapted to ecological conditions

Questa è la versione Post print del seguente articolo:

Original

The weather dictates the rhythms: Alpine chamois activity is well adapted to ecological conditions / Brivio, Francesca; Bertolucci, C; Tettamanti, F; Filli, F; Apollonio, Marco; Grignolio, S.. - In: BEHAVIORAL ECOLOGY AND SOCIOBIOLOGY. - ISSN 1432-0762. - 70:8(2016), pp. 1291-1304. [10.1007/s00265-016-2137-8]

Availability: This version is available at: 11388/85037 since: 2022-05-28T23:00:24Z

Publisher:

Published DOI:10.1007/s00265-016-2137-8

Terms of use:

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

Publisher copyright

note finali coverpage

(Article begins on next page)

```
Manuscript
```

1	The weather dictates the rhythms: Alpine chamois activity is well
2	adapted to ecological conditions
3	
4	Francesca Brivio ^{*a} , Cristiano Bertolucci ^b , Federico Tettamanti ^a , Flurin Filli ^c , Marco
5	Apollonio ^a , Stefano Grignolio ^a
6	
7	^a University of Sassari, Department of Science for Nature and Environmental Resources, via
8	Muroni 25, I-07100, Sassari, Italy
9	
10	^b Department of Biology and Evolution, University of Ferrara, Via L. Borsari 46, I-44121
11	Ferrara, Italy
12	
13	^c Parc Naziunal Svizzer, 7530 Zernez, Switzerland
14	
15	* Corresponding author: Francesca Brivio (e-mail: <u>fbrivio@uniss.it</u> , Tel. (+39) 079 -228667,
16	Fax (+39) 079-228665)
17	
18	
19	
20	
21	Author Contributions: SG, FB, MA originally formulated the idea; FF and FT conducted
22	fieldwork; SG, CB, FB and MA and collaborated in imaging analysis; FB, CB and SG
23	performed statistical analyses; FB wrote the manuscript; other authors provided editorial
24	advice.

25 Abstract

26 Activity rhythms play an important role in the ecological relations of a species and form part 27 of its evolutionary adaptation. Such rhythms are strongly synchronised with the annual cyclic 28 changes by environmental stimuli, the so-called Zeitgebers. Animals' reliance on 29 environmental stimuli is highly species-specific and allows behavioural adjustments to be 30 made in preparation for the conditions expected in each season. Here we investigated daily 31 and annual activity rhythms of Alpine chamois (Rupicapra rupicapra) by analysing high-32 resolution long-term data from females and males monitored with GPS-collars. This first 33 detailed field study of chamois activity showed that this species exhibited clear daily and 34 annual activity rhythms entrained to the light-dark cycle. Chamois were more active during 35 spring-summer and less active during winter, likely in response to the variation in the 36 availability of food resources: both sexes appeared to maximise energy intake during the 37 season offering the highest amount of food resources to compensate for poor food supply 38 during winter. Daily activity was significantly influenced by climatic factors and poorly 39 affected by individual variables. As activity was strongly influenced by temperature and 40 radiation we conjectured that it was also critically dependent upon animals' thermal balance. 41 Our analyses showed a negative correlation between daily activity and adverse climatic 42 conditions (i.e., precipitation and snow cover). In conclusion, our study pointed out that 43 chamois is well adapted to Alpine environment and seasonality. However, it can be 44 susceptible to climate change, which may produce a mismatch between its annual activity 45 rhythms, evolved over thousands of years.

46

47 Key words: Activity rhythms, Alpine environment, climatic conditions, GPS-collars,
48 *Rupicapra rupicapra*

49 Introduction

50 Animals living in temperate habitats face pronounced changes in climatic conditions 51 and food availability. This is particularly true for the species inhabiting Alpine regions where 52 seasonal changes are marked. Alpine winter is generally characterised by a sharp decrease in 53 temperature and the presence of snow which lead to an increase in energetic demands at a 54 time when food resources are extremely scarce. Conversely, during spring and summer the 55 highly structured geomorphology of Alpine environments entails high spatial heterogeneity 56 in climatic conditions, as well as in the distribution of natural resources (i.e., high-quality 57 forage, Pettorelli et al. 2005).

To cope with these annual cyclic changes, animals undergo several physiological,
morphological and behavioural adjustments (Lovegrove 2005; Paul et al. 2008). Significant
examples of these seasonal responses are: the evolution of winter-specific physiological and
behavioural patterns to conserve energy (i.e., a highly insulative pelage: Martinet et al. 1992;
hibernation: Kondo et al. 2006; huddling: Gilbert et al. 2010), seasonal migrations (e.g.,
Kumar et al. 2010), and the periodic repetition of the rutting season (e.g., Rutberg 1987;
Noordwijk et al. 1995).

65 The seasonal coordination of these phenotypic modifications is regulated by 66 endogenous clocks, which help animals to anticipate and prepare for environmental changes 67 (Pittendrigh 1993; Paul et al. 2008). Circadian oscillators (i.e., endogenous clocks) are 68 synchronised (*Entrained*) with the time of the year by periodical changes in environmental 69 stimuli, the so-called Zeitgebers (from German, Zeit: "time"; geber: "to give"; Pittendrigh 70 1993). Day length (i.e., photoperiod), the most accurate natural predictor of annual phase, is 71 one of the main Zeitgebers (Gorman et al. 2001; Zucker 2001; Gwinner 2003). However, 72 other periodical factors in the animals' environment can also act as time cues: in particular, 73 food and water availability, ambient temperature and social signals have been shown to affect 74 seasonal traits (reviewed in Paul et al. 2008). Besides the entrainment effect, environmental

variables (geophysical and biological) usually have also a direct inhibitory or enhancing effect on the different physiological and behavioural patterns, resulting in a masking effect on the circannual and the circadian rhythms (Aschoff et al. 1982). Animals' reliance on all these cues is highly species-specific and allows physiological adjustments to be made in preparation for the conditions expected in each season (Paul et al. 2008).

80 Behavioural adaptations to the change in environmental and social conditions have 81 been shown to respond more rapidly than physiological or morphological ones (Van Buskirk 82 2012). Therefore, one of the first reactions to seasonal acclimatisation that vertebrates can 83 implement is the modification of their behavioural rhythms. The activity rhythms of several 84 species are highly entrained to the variation in environmental cues and strongly influenced by 85 individual characteristics, such as sex, age and body size (e.g., Prates and Bicca-Marques 86 2008; Owen-Smith and Goodall 2014). Activity rhythms are important in controlling the 87 energy balance of a species (Aschoff 1979), thus resulting in a complex trade-off between 88 optimal foraging time, social activities and environmental constraints (Aschoff 1963). These 89 rhythms play a major role in the ecological relations of a species and form part of its 90 evolutionary adaptation (Aschoff 1958; Pittendrigh 1993).

91 Nowadays, the global climate change is altering seasonal patterns. For instance, global 92 meta-analyses documented a significant mean shift towards earlier spring timing of 2.3 93 days/decade (Parmesan and Yohe 2003). Consequently, animal rhythms may become out of 94 phase with the periodic challenges posed by their environment (e.g., Pettorelli et al. 2007; 95 Plard et al. 2014). It has been demonstrated that earlier spring has already caused 96 phenological modifications in most taxonomic groups (Parmesan and Yohe 2003; Root et al. 97 2003). The Alpine ecosystems are among the most vulnerable to rapid climate change 98 (Ernakovich et al. 2014). Their climate is expected to be particularly affected by global 99 warming: not only by rising temperatures (about 0.25 °C /decade by the mid- 21st century), 100 but also by changes in the seasonal cycle of precipitation, global radiation, humidity,

temperature and precipitation extremes, as well as by closely related impacts such as floods,droughts, snow cover and natural hazards (Gobiet et al. 2014).

103 Studies on the current relationship between climate and animal behaviour should shed light on the species' ability to track and adapt to climate change (Dunbar 1998; Lehmann et 104 105 al. 2008; Korstjens et al. 2010). Alpine chamois is a relatively eurythermic mountain 106 ungulate, which has adapted to a wide range of temperatures and is distributed over a broad 107 altitudinal range (500-3100 m; Shackleton 1997; Spitzenberger et al. 2001). Indeed, this 108 species is the most widespread ungulate in the Alps (Apollonio et al. 2010) and is well 109 adapted to their environment and seasonality. It is characterised by nearly monomorphic 110 sexes, with only a weak and highly seasonal dimorphism in body mass (from approx. 40% 111 before the rut, to approx. 6% in winter and 4% in spring; Garel et al. 2009; Rughetti and 112 Festa-Bianchet 2011). The seasonal changes in body mass, together with the very high 113 survival rate of both sexes (Corlatti et al. 2012), are indicative of a conservative strategy, 114 more pronounced in males than in females. Indeed, during summer chamois accumulate fat 115 resources (Pérez-Barbería et al. 1998) to be used during the rut (i.e., November), in order to 116 reduce over-winter mortality costs. In other words, this ungulate is an ideal case-study to 117 investigate the plasticity in the response of a species to the variation in the environmental 118 conditions of mountainous regions.

Here we investigated daily and annual activity rhythms of Alpine chamois. In particular, we compared activity levels of chamois of different sex and age class throughout the years in order to investigate individual strategies of adaptation to seasonal and climatic variations. Since in mountainous regions the harshest conditions occur during the winter season, we focused a part of our analyses only on winter data to better understand how chamois managed to control their energy balance in response to different external factors under severe environmental conditions.

126

127 Materials and methods

128 Study Area and Population

This study was carried out in the Swiss National Park (SNP; 46°40'10.74" N, 10°9'15.15" E), in southeastern Switzerland. The SNP is an area of integral protection where hunting is not allowed. The area covers 170 km² and elevation ranges from 1500 to 3170 m above sea level (a.s.l.). The landscape consists of forests of Arolla pine (*Pinus cembra*), larch (*Larix deciduos*) and mountain pine (*Pinus mugo*) below the tree line (approx. 2200 m a.s.l.) and alpine grasslands and rocky slopes above it. Annual precipitation amounts to 700-1200 mm.

The only predator of chamois is the Golden eagle (Aquila chrysäetos) that typically 136 137 preys upon small kids. During the study period (2010-2013), population census showed that 138 chamois density was relatively stable: during the census of 2013, the rangers counted 6.2 139 individuals/Km² (523 females, 274 males and 267 kids + yearlings). Between 2010 and 2013, 140 11 females and 6 males were captured with box traps and equipped with GPS Plus Collar 141 (Vectronic Aerospace GmbH, Berlin, Deutschland). The exact year of birth was determined 142 by counting the number of annual incremental growth rings on the horns (Schröder and von 143 Elsner-Schack 1985). The age at capture ranged between 3 and 15 years for females and 144 between 6 and 13 years for males.

145

146 **Data collection**

147 The locomotor activity was recorded continuously from the capture of the individuals 148 (the first one in March 2010) until the conclusion of data collection (November 2013) or until 149 the collar stopped working. We collected an average of 514 days/chamois (minimum 190 – 150 maximum 780 days). The activity was recorded by means of an analogue accelerometer (i.e., 151 activity sensor) installed on the GPS-collars, which measured the activity based on the actual 152 acceleration they experienced. The accelerometer had a dynamic range from -2G to +2G

(G=gravitational constant) and measured activity as the change of static acceleration (gravity) 153 154 and dynamic acceleration (collar) four times/second. The activity values were given within a 155 relative range between 0 and 255. Value 255 was equal to -2G / +2G and it indicated the 156 maximum acceleration, whereas 0 indicated no acceleration at all. Measurements were 157 averaged over sampling intervals of 5 minutes and stored with the date and time associated. 158 Weather data (daily mean temperature, daily mean radiation, daily precipitation 159 amount, daily mean wind speed and daily snow cover) were provided by the Swiss 160 Meteorological Institute (Begert et al. 2005) from the Buffalora station located at approx.13 161 km from the study area at an altitude of 1968 m a.s.l. (46°38'53.29" N; 10°16'1.86" E).

162

163 Data analysis

164 The presence of periodicity in the locomotor activity rhythms was determined by means of the Lomb-Scargle periodogram (LSP) analysis and confirmed with the χ^2 165 periodogram analysis, by using the ActogramJ software for circadian analysis (Schmid et al. 166 167 2011). The significance was set to p < 0.05. For each chamois, analyses were performed each month with intervals of 10 days. To exclude that 8- and 12-hour periods were harmonics of 168 169 daily (24 hours) periods or vice versa, we compared the amplitude (PN and Qp) of the peaks 170 (van Oort et al. 2005). Data were presented graphically as actograms by using the Activity 171 Pattern ver. 1.3.1 software (Vectronic Aerospace GmbH, Berlin, Deutschland; Fig. 1, 172 Supplementary material Appendix 1 Fig. A1-A15). 173 On each day of data collection, mean activity $(\pm SD)$ during the day was calculated for each chamois by using the Activity Pattern ver. 1.3.1 software (Vectronic Aerospace GmbH, 174 175 Berlin, Deutschland). We focused on the daily mean activity (DMA) of each individual to 176 assess the effect of intrinsic and extrinsic factors on chamois activity patterns. To allow for 177 the likelihood of a non-linear response to the covariates, we chose to fit a non-parametric 178 model to the data in which the exact functional form was determined from the data

179 themselves and not specified *a priori*. Generalised Additive Mixed Models (GAMMs) are an 180 ideal tool for such analysis as they are flexible in modelling the shape of non-linear 181 relationships. Non-parametric smoothing functions are used on sections of the data and the 182 response curves are connected at their end-points to generate an overall smooth curve (Wood 183 2006). In addition to the non-parametric smoothing functions, parametric fixed and random 184 predictor terms may also be included, as in a conventional mixed-effects model. The DMA 185 was modelled as the response variable by fitting alternative GAMMs, implemented within 186 the mgcv package (version 1.7-26; Wood 2006) in R (version 3.0.2; R Core Team 2014). 187 Female and male DMA were modelled separately by using their identity as a random effect. 188 The predictor variables considered were: temperature, radiation, precipitation, wind speed, 189 snow cover and chamois age class. To subdivide our sample size homogeneously, we used 190 four uniform age classes: 1) 3-5 years old; 2) 6-8 years old; 3) 9-11 years old; 4) older than 191 11 years. Precipitation and wind speed were log-transformed in order to improve the 192 homogeneity of data distribution. Moreover, in order to identify the fluctuating pattern of 193 variation of DMA throughout the year, we inserted the Julian date as a continuous fixed 194 effect in the models. To check for multicollinearity between the explanatory variables, we 195 calculated the variance inflation factors (VIF) for each of them. Temperature was found to be 196 correlated to radiation, as well as snow cover to precipitation and, consequently, they were 197 used separately in alternative models. The VIF values of the other variables were < 3, thus 198 indicating no severe multicollinearity between them (Zuur et al. 2009). Effects of continuous 199 predictor variables were initially modelled as natural cubic spline functions and the optimal 200 roughness of the smoothing terms was determined by minimising the generalised cross-201 validation value. When the effective degrees of freedom (*edf*) of a predictor variable were 1 202 and the graphical inspection confirmed a linear relationship with the response variable, we 203 refitted the model omitting the smoothing function. Between these two models, we selected 204 those with the lower AIC value. Effects of the Julian date were modelled as a cyclic cubic

regression spline in order to ensure that the value of the smoother at the far left point (1January) was the same as the one at the far right point (31 December).

207 Analyses were performed using the information-theoretic approach (Dochtermann and 208 Jenkins 2011). In the light of theory, biological relevance and previous research on related 209 species, we defined a set of alternative hypotheses, which were used to construct 47 *a priori* 210 GAMM models (Supplementary material Appendix 2 Table A1). The alternative models 211 were ranked and weighted with the minimum AIC criterion (Symonds and Moussalli 2011). 212 We confirmed the global goodness-of-fit (i.e., homoscedasticity, normality of errors and 213 independence) of the best models by visual inspection of residuals (Zuur et al. 2009). 214 In order to point out the influence of snow cover on female and male DMA, we 215 constructed two sets of 24 a priori models (one for each sex, Supplementary material 216 Appendix 2 Table A2) by using only data from 1 October to 30 April (i.e., the period in 217 which snow was potentially present). Then, we followed the same approach described above, 218 in order to identify the best model among the alternative ones.

219

220 **Results**

221 A total of 5884 days/chamois for females and a total of 2869 days/chamois for males 222 were obtained during data collection. All chamois showed a clear daily rhythm of locomotor 223 activity entrained to the natural light-dark cycle (Fig. 1, Supplementary material Appendix 1 224 Fig. A1-A15). Their activity was not exclusively diurnal and a nocturnal activity was also 225 present during most of the year (Fig. 1-2, Supplementary material Appendix 1 Fig. A1-A15). 226 The pattern of locomotor activity showed a marked seasonal variation in both females males 227 and gradually changed from unimodal (with a single peak in December-May/June) to 228 bimodal/trimodal during summer and autumn (Fig. 2). During summer, the periodogram analysis showed a significant rhythm with a period of 12 hours (P<0.05, LSP; Fig. 2B, F) for 229 230 both males and females. The bimodal pattern of activity peaked at dawn and dusk, with the

activity at dusk being higher than that at dawn (Fig. 1, Supplementary material Appendix 1
Fig. A1-A15). During autumn, the period shorted to 8 hours (*P*<0.05, LSP; Fig. 2C, G) and
the pattern of activity showed a marked increase of nocturnal activity with a peak at midnight
(Fig. 1, Supplementary material Appendix 1 Fig. A1-A15).

235 Global models – The best model selected (with the lowest AIC value) for females 236 included all predictor variables: age class, day of the year, temperature, precipitation and 237 wind speed. For males, the best model included all variables except wind speed (Table 1). 238 Results showed that age class did not significantly affect the DMA of both female and male 239 chamois. Throughout the year, the DMA of both sexes followed a bell-shaped pattern, with a maximum of activity at the beginning of summer (females: peak on the 178th day of the year 240 -27 June; males: peak on the 185th day of the year -1 July) and minimum values in January 241 242 (Fig 3A). A second peak of activity was detected from mid-October to the end of November. 243 This second peak was higher in males than in females. Temperatures significantly affected 244 chamois DMA. The pattern of the relationship was similar for females and males: DMA 245 increased with increasing daily mean temperatures until 3-5 °C and then decreased with a 246 slight difference between the two sexes (Fig. 3B). Female and male DMA was also 247 significantly influenced by precipitation: DMA decreased linearly with increasing daily 248 precipitation (females: $\beta = -2.93 \pm 0.68$; males: $\beta = -1.02 \pm 0.14$). Finally, female DMA 249 decreased linearly with increasing wind speed ($\beta = -3.79 \pm 0.68$). 250 Winter models - For both sexes the best models selected included: radiation, wind 251 speed and snow cover (Table 2). Only for females, the model selected contained also age 252 class but with no significant statistical effect. Results showed that in this period (1 October -

253 1 April) the variation in chamois DMA was better explained by radiation than by

temperature, with a positive non-linear effect for female (Fig. 4A) and a positive linear effect

for males ($\beta = 0.03 \pm 0.002$). Chamois DMA considerably decreased with increasing snow

cover (Fig. 4B), while the relationship between DMA and wind speed followed a hump-

shaped pattern, with a peak of activity around 1.19 m/s for females and around 1.18 m/s formales (Fig. 4C).

259

260 Discussion

This first detailed field study on Alpine chamois activity showed that this species exhibited clear daily and annual activity rhythms entrained to the light-dark cycle. Daily activity was significantly influenced by climatic factors (i.e., temperature, radiation, snow cover and wind speed) and poorly affected by individual variables, such as sex and age class. As a matter of fact, even though females were more active during all seasons, the two sexes had a similar reaction to the environmental factors analysed, with no significant difference among individuals of different ages.

268 Daily activity rhythms varied from a unimodal pattern during winter to a bimodal pattern 269 during summer and autumn. It is no coincidence that the findings on DMA showed that 270 chamois were more active during spring and early summer and less active during winter. 271 Previous studies indicated that chamois typically spend about 67-70% of their active time 272 foraging (Rüttimann et al. 2008). Thus, in accordance with other studies on ungulate 273 behaviour (Collins et al. 1978; Risenhoover 1986; Beier and McCullough 1990), we can 274 assume that changes in chamois activity mainly reflected changes in the time spent feeding. 275 Hence, the variation in the activity patterns indicated the response of chamois to seasonal 276 variations in the availability of food resources. In mountainous areas, the rapid growth of 277 fresh plant forage typically begins immediately after snowmelt (April–June; earlier at lower 278 altitudes), thus providing an abundant and protein-rich source of food for a relatively brief 279 period. The shift from unimodal to bimodal pattern during spring and summer may be a 280 physiological consequence related to rumination rhythms. Whenever ruminants eat more 281 abundant forage, in fact, they typically fill their rumen more rapidly and therefore have to lie 282 down to ruminate and digest plant material earlier in the day (VanSoest 1994). In addition, if

283 the forage is protein-rich, the digestion is faster so that animals are able to engage in a second 284 bout of foraging activity during the day (VanSoest 1994). Moreover, even the modifications 285 in chamois daily activity levels pointed out by our analyses resulted to be a good strategy of 286 adaptation to the availability of food resources. As reported for other northern-temperate 287 mammals (Tyler and Blix 1990; Adamczewski et al. 1997; Parker et al. 2009), the peak of 288 activity during the spring-summer months may be an adaptation that enables chamois to use 289 the most favourable forage conditions in order to storage fat tissue to survive during long, 290 cold and snowy winters. Accordingly, the reduction of activity during winter and the use of 291 the unimodal pattern showed by our analyses may be seen as an adaptation strategy to 292 conserve energy in response to the unfavourable conditions (i.e., harsh weather, decline in 293 forage quality and decreased food availability due to snow cover). For both female and male 294 chamois, the decrease in forage quality and quantity may be offset by a decrease in metabolic 295 rate and the use of the stored energy reserves. The reduction of food intake and, in general, of 296 metabolic rate during winter is widespread amongst northern ungulates as a strategy to cope 297 with cold conditions and low availability of food resources (reviewed in Arnold et al. 2004). 298 At first glance, seasonal variations in the activity pattern coincided with the variation in 299 the photoperiod, thus suggesting that the latter is the main Zeitgeber. Nevertheless, 300 throughout the year activity rhythms can deviate from the photoperiod cycle, therefore 301 suggesting that other biological and environmental factors may act as time cues for the 302 activity regulation of chamois. In November, for example, a peak of activity -more 303 pronounced in males than in females- disrupted the pattern of decrease from mid-summer to 304 winter. Indeed, the chamois' mating season occurs in this period, with a peak between 16 and 305 25 November (Corlatti et al. 2013b). This peak of activity could not be ascribed to foraging 306 behaviour as males typically reduce forage intake during the rut (Willisch and Ingold 2007). 307 Therefore, in accordance with Corlatti et al. (2013b), the high activity levels found in this 308 study were likely due to social activities aimed at gaining mating opportunities. Our results

309 showed that chamois nocturnal activity increased during autumn, with a third peak of daily 310 activity detected at midnight. Thus, our analyses match with the previous knowledge on this 311 species and contribute to explain why males lose weight faster than females during the 312 rutting period (Garel et al. 2009; Rughetti and Festa-Bianchet 2011). In fact, during the rut 313 they spent significantly less time foraging (reduction of food intake; Willisch and Ingold 314 2007; Corlatti et al. 2013a) and were active for a longer period (i.e., increase in energy 315 expenditure).

316 Climate is another time cue for chamois activity: they adjusted their activity responding to variations in temperature, as well as to changes in precipitation levels, wind speed and, 317 318 during winter, snow cover and solar radiation. Air temperature appeared to significantly 319 influence chamois activity rhythms, either reducing or stirring daily activity levels. Similar 320 correlations between daily activity and air temperature have been found for other ungulate 321 species (Belovsky and Slade 1986; Beier and McCullough 1990; Schmitz 1991; Owen-Smith 322 1998; Loe et al. 2007; Pipia et al. 2008; Signer et al. 2011). The response of chamois activity 323 to temperature was parabolic, with the highest activity levels at temperatures between 3 and 324 5° C and a decline in activity levels at higher and lower temperatures for both sexes (Fig. 3B). 325 Therefore, activity appeared to be strictly dependent upon animal thermal balance. On the 326 one hand, when air temperatures are significantly below the animals' thermoneutral zone, 327 decreased activity can be seen as a strategy to lower the costs of thermoregulation by seeking 328 shelter (in time budget terms, by resting) in order to reduce heat loss. Accordingly, Moen 329 (1976) and Gates and Hudson (1979) showed that lying position and inactivity significantly 330 reduce the metabolic costs of thermoregulation during cold weather. On the other hand, when 331 the air temperature rises significantly above the species' thermoneutral zone, the reduction of 332 activity is likely an attempt to avoid thermal overload, by reducing heat-generating activities 333 (i.e., feeding, moving and maybe even social interaction). This last result is consistent with a 334 previous observational study on Alpine chamois, which reported that animals allocated less

335 time to foraging with increasing temperatures during summer (Mason et al. 2014). Several 336 authors showed that thermal stress places an upper limit on the time animals may devote to 337 daily activity (e.g., Orthoptera: Parker 1982; Chappell 1983; Corvus brachyrhynchos, Sciurus 338 carolinensis: Kilpatrick 2003; Castor Canadensis: Belovsky 1984b; Lepus americanus: 339 Belovsky 1984a; Alces alces: Belovsky 1978; Tragelaphus strepsiceros: Owen-Smith 1998). 340 Each species developed a tolerance to heat depending on the habitat in which it lives. 341 Therefore, ungulates living in temperate regions appear to have their activity levels reduced 342 by lower temperatures in comparison with species living in the African savannah (e.g., 343 Belovsky and Jordan 1978; Belovsky and Slade 1986; Owen-Smith 1998). Our results 344 showed that chamois started to decrease their activity already at 3-5°C. Given the rapid 345 alteration in the seasonal cycle of climatic variables characterising the last decades, the annual activity rhythms of chamois, which have evolved over thousands of years, may 346 347 become out of phase with the future periodic challenges posed by their environment. As a 348 consequence of the rising temperatures predicted for the Alpine regions (about 0.25 $^{\circ}C$ / 349 decade, Gobiet et al. 2014), chamois daily activity is expected to decrease, particularly during 350 summer, likely compromising their ability to accumulate body reserves. This may affect the 351 animals' ability to cope with the food shortages of the Alpine winter season, and therefore it may have important effects on individual life history. This scenario could be worsened by the 352 353 forecasts on global radiation along the Alpine ridge, which indicate an increase of radiation 354 during summer and a decrease during winter (Gobiet et al. 2014). On the one hand, 355 increasing radiation during summer may exacerbate the reduction of chamois activity in 356 order to avoid thermal overload. On the other hand, the decrease of radiation during winter may increase thermoregulation costs. Indeed, we found that chamois activity was positively 357 358 related to radiation during winter months, thus we conjectured that during winter chamois 359 were able to benefit from the absorption of solar radiation. By using sunny areas and taking 360 advantage of exogenous heat for thermoregulation they managed to reduce the amount of

and genous heat produced to maintain constant body temperature. Consistently, Signer et al.
(2011) showed that radiant heat-assisted rewarming play a key-role as a strategy to endure
harsh over-wintering conditions in Alpine ibex. In this framework, it is useful to note the role
of the black coat of chamois during the winter season. Our findings suggested that this
specific evolutionary characteristic favoured a greater absorption of solar radiation
(exogenous heat) and thus an increase of daily activity.

367 Conversely, chamois winter activity may benefit from the significant decrease in alpine 368 snow duration and abundance resulting from the global climate change (Gobiet et al. 2014). 369 Our analyses on winter data highlighted a strong negative correlation between daily activity 370 and snow cover. Several studies analysed the effect of snow cover on spatial behaviour 371 (Tyler and Blix 1990), activity rhythms (Cederlund 1981; Beier and McCullough 1990), life 372 history and population dynamics (Jacobson et al. 2004; Apollonio et al. 2013; Willisch et al. 373 2013). It is well known that deep snow causes a higher energy expenditure for large 374 mammals, likely as a consequence of both lower food availability and increased energy 375 expenditure related to locomotion. Our study showed that chamois reduced their activity 376 when the thickness of snow increased: indeed, when snow cover was thicker, chamois 377 adopted a safe strategy by reducing their total activity, likely because of the reduced food 378 availability, but also to avoid mobility problems and to reduce accident risks (i.e., 379 avalanches). As chamois have an interdigital membrane to increase the distribution of weight 380 and favour mobility on snow surfaces, we may expect a reduced effect of snow cover on their 381 behaviour. Nonetheless, this anatomic adaptation to snow surfaces does not seem sufficient 382 to maintain regular activity rhythms in case of deep snow cover. Further research should 383 assess the effect of such activity reduction on individual life history and population 384 dynamics.

In conclusion, our study pointed out that well-detailed information on activity may be
used to analyse how behavioural strategies evolved in animals to adapt to their environment,

take advantage of available resources and respond to environmental changes (e.g., global

388 climate change). In our case, we showed that Alpine chamois is well adapted to high

389 mountain environment and seasonality, even though it can be susceptible to climate change.

390 In this respect, our work showed the complexity of potential behavioural responses to the

391 variation in multiple environmental factors, such as climatic variables and food resources.

392

393 Acknowledgments

394 We wish to thank all the park rangers of the Swiss National Park who captured, marked and

395 monitored the chamois. This project was supported by the Italian Ministry of Education,

396 University and Research (PRIN 2010–2011, 20108 TZKHC, J81J12000790001). SG had the

397 support of the fund "P.O.R. F.S.E. 2007–2013—Obiettivo competitività regionale e

398 occupazione. Asse IV Capitale umano—Linea di attività 1.3.1". The English version was
399 edited by C. Pole.

400

401 **Conflict of Interest:** The authors declare that they have no conflict of interest.

402

403 **Ethical standards** Female and male Alpine chamois were tagged by rangers of the Swiss

404 National Park under the legal authorization of the Swiss Veterinary Office. All applicable

405 institutional and/or national guidelines for the care and use of animals were followed.

406 **References**

407	Adamczewski JZ, Flood PF, Gunn A (1997) Seasonal patterns in body composition and
408	reproduction of female muskoxen (Ovibos moschatus). J Zool 241:245-269
409	Apollonio M, Andersen R, Putman R (2010) European ungulates and their management in
410	the 21st century. Cambridge University Press, Cambridge, UK
411	Apollonio M, Brivio F, Rossi I, Bassano B, Grignolio S (2013) Consequences of snowy
412	winters on male mating strategies and reproduction in a mountain ungulate. Behav
413	Process 98:44-50
414	Arnold W, Ruf T, Reimoser S, Tataruch F, Onderscheka K, Schober F (2004) Nocturnal
415	hypometabolism as an overwintering strategy of red deer (Cervus elaphus). Am J
416	Physiol Regul Integr Comp Physiol 286:R174-81
417	Aschoff J (1958) Tierische Periodik unter dem Einfluß von Zeitgebern. Z Tierpsychol 15:1-
418	30
419	Aschoff J (1963) Comparative Physiology: Diurnal Rhythms. Annu Rev Physiol 25:581-600
420	Aschoff J (1979) Circadian rhythms: influences of internal and external factors on the period
421	measured in constant conditions. Z Tierpsychol 49:225-49
422	Aschoff J, Daan S, Honma KI (1982) Zeitgebers, entrainment, and masking: some unsettled
423	questions. In: Aschoff J, Daan S, Groos G (eds) Vertebrate Circadian Systems.
424	Springer Berlin Heidelberg, pp 13-24
425	Begert M, Schlegel T, Kirchhofer W (2005) Homogeneous temperature and precipitation
426	series of Switzerland from 1864 to 2000. Int J Climat 25:65-80
427	Beier P, McCullough DR (1990) Factors influencing white-tailed deer activity patterns and
428	habitat use. Wildl Monogr 109:3-51
429	Belovsky GE (1978) Diet optimization in a generalist herbivore: The moose. Theor Popul
430	Biol 14:105-134

- Belovsky GE (1984a) Moose and snowshoe hare competition and a mechanistic explanation
 from foraging theory. Oecologia 61:150-159
- 433 Belovsky GE (1984b) Summer diet optimization by beaver. Am Midl Nat 111:209-222
- 434 Belovsky GE, Jordan PA (1978) The time-energy budget of a moose. Theor Popul Biol
 435 14:76-104
- 436 Belovsky GE, Slade JB (1986) Time budgets of grassland herbivores: body size similarities.
 437 Oecologia 70:53-62
- 438 Cederlund G (1981) Daily and seasonal activity pattern of roe deer in a boreal habitat. Swed
 439 Wildl Res 11:315-348
- Chappell MA (1983) Thermal limitations to escape responses in desert grasshoppers. Anim
 Behav 31:1088-1093
- 442 Collins WB, Urness PJ, Austin DD (1978) Elk diets and activities on different lodgepole pine
 443 habitat segments. J Wildl Manag 42:799-810
- 444 Corlatti L, Bassano B, Valencak TG, Lovari S (2013a) Foraging strategies associated with
 445 alternative reproductive tactics in a large mammal. J Zool 291:111-118
- 446 Corlatti L, Caroli M, Pietrocini V, Lovari S (2013b) Rutting behaviour of territorial and
- 447 nonterritorial male chamois: is there a home advantage? Behav Process 92:118-24
- 448 Corlatti L, Lebl K, Filli F, Ruf T (2012) Unbiased sex-specific survival in Alpine chamois. Z
 449 Saugetierkd 77:135-139
- 450 Dochtermann NA, Jenkins SH (2011) Developing multiple hypotheses in behavioral ecology.
 451 Behav Ecol Sociobiol 65:37-85
- 452 Dunbar RIM (1998) Impact of global warming on the distribution and survival of the gelada
- 453 baboon: a modelling approach. Global Change Biol 4:293-304
- 454 Ernakovich JG, Hopping KA, Berdanier AB, Simpson RT, Kachergis EJ, Steltzer H,
- Wallenstein MD (2014) Predicted responses of arctic and alpine ecosystems to altered
 seasonality under climate change. Global Change Biol 20:3256-3269

458	Growth in Alpine Chamois. J Mammal 90:954-960
459	Gates CC, Hudson RJ (1979) Effects of posture and activity on metabolic responses of
460	Wapiti to cold. J Wildl Manag 43:564-567
461	Gilbert C, McCafferty D, Le Maho Y, Martrette JM, Giroud S, Blanc S, Ancel A (2010) One
462	for all and all for one: the energetic benefits of huddling in endotherms. Biol Rev
463	Camb Philos Soc 85:545-69
464	Gobiet A, Kotlarski S, Beniston M, Heinrich G, Rajczak J, Stoffel M (2014) 21st century
465	climate change in the European Alps - A review. Sci Total Environ 493:1138-1151
466	Gorman MR, Goldman BD, Zucker I (2001) Mammalian photoperiodism. In: Takahashi JS,
467	Turek FW, Moore RY (eds) Handbook of behavioral neurobiology: circadian clocks.
468	Kluwer Academic/Plenum Publishers, New York, pp 481–508
469	Gwinner E (2003) Circannual rhythms in birds. Curr Opin Neurobiol 13:770-8
470	Jacobson AR, Provenzale A, von Hardenberg A, Bassano B, Festa-Bianchet M (2004)
471	Climate forcing and density dependence in a mountain ungulate population. Ecology
472	85:1598-1610
473	Kilpatrick AM (2003) The impact of thermoregulatory costs on foraging behaviour: a test
474	with American Crows (Corvus brachyrhynchos) and eastern grey squirrels (Sciurus
475	carolinensis). Evol Ecol Res 5:781-786
476	Kondo N, Sekijima T, Kondo J, Takamatsu N, Tohya K, Ohtsu T (2006) Circannual control
477	of hibernation by HP complex in the brain. Cell 125:161-72
478	Korstjens AH, Lehmann J, Dunbar RIM (2010) Resting time as an ecological constraint on
479	primate biogeography. Anim Behav 79:361-374
480	Kumar V, Wingfield JC, Dawson A, Ramenofsky M, Rani S, Bartell P (2010) Biological
481	clocks and regulation of seasonal reproduction and migration in birds. Physiol

Garel M, Loison A, Jullien J-M, Dubray D, Maillard D, Gaillard J-M (2009) Sex-specific

482 Biochem Zool 83:827-35

457

- 483 Lehmann J, Korstjens AH, Dunbar RIM (2008) Time and distribution: a model of ape
 484 biogeography. Ethol Ecol Evol 20:337-359
- Loe LE, Mysterud A, Stien A, Steen H, Evans DM, Austrheim G (2007) Positive short-term
 effects of sheep grazing on the alpine avifauna. Biol Lett 3:110-112
- 487 Lovegrove B (2005) Seasonal thermoregulatory responses in mammals. J Comp Physiol B
 488 175:231-247
- 489 Martinet L, Mondain-Monval M, Monnerie R (1992) Endogenous circannual rhythms and
 490 photorefractoriness of testis activity, moult and prolactin concentrations in mink

491 (*Mustela vison*). J Reprod Fert 95:325-38

492 Mason THE, Stephens PA, Apollonio M, Willis SG (2014) Predicting potential responses to
493 future climate in an alpine ungulate: interspecific interactions exceed climate effects.

494 Global Change Biol 20:3872-3882

- 495 Moen AN (1976) Energy conservation by white-tailed deer in the winter. Ecology 57:192496 198
- 497 Noordwijk AJV, McCleery RH, Perrins CM (1995) Selection for the timing of great tit
- 498 breeding in relation to caterpillar growth and temperature. J Anim Ecol 64:451-458
- 499 Owen-Smith N (1998) How high ambient temperature affects the daily activity and foraging
- 500 time of a subtropical ungulate, the greater kudu (*Tragelaphus strepsiceros*). J Zool
 501 246:183-192
- 502 Owen-Smith N, Goodall V (2014) Coping with savanna seasonality: comparative daily
 503 activity patterns of African ungulates as revealed by GPS telemetry. J Zool 293:181-
- 504 191
- 505 Parker KL, Barboza PS, Gillingham MP (2009) Nutrition integrates environmental responses
 506 of ungulates. Funct Ecol 23:57-69
- 507 Parker MA (1982) Thermoregulation by diurnal movement in the barberpole grasshopper
 508 (*Dactylotum bicolor*). Am Midl Nat 107:228-237

- 509 Parmesan C, Yohe G (2003) A globally coherent fingerprint of climate change impacts
- 510 across natural systems. Nature 421:37-42
- 511 Paul MJ, Zucker I, Schwartz WJ (2008) Tracking the seasons: the internal calendars of

512 vertebrates. Phil Trans R Soc B 363:341-361

- 513 Pérez-Barbería F, Mutuberría G, Nores C (1998) Reproductive parameters, kidney fat index,
- and grazing activity relationships between the sexes in Cantabrian chamois *Rupicapra pyrenaica parva*. Acta Theriol 43:311–324
- 516 Pettorelli N, Mysterud A, Yoccoz NG, Langvatn R, Stenseth NC (2005) Importance of
- 517 climatological downscaling and plant phenology for red deer in heterogeneous
- 518 landscapes. Proc R Soc Lond B 272:2357-2364
- 519 Pettorelli N, Pelletier F, Hardenberg Av, Festa-Bianchet M, Côté SD (2007) Early onset of
- 520 vegetation growth vs. rapid green-up: impacts on juvenile mountain ungulates.
 521 Ecology 88:381-390
- 522 Pipia A, Ciuti S, Grignolio S, Luchetti S, Madau R, Apollonio M (2008) Influence of sex,
 523 season, temperature and reproductive status on daily activity patterns in Sardinian
- 524 mouflon (*Ovis orientalis musimon*). Behaviour 145:1723-1745
- 525 Pittendrigh CS (1993) Temporal organization: reflections of a Darwinian clock-watcher.
- 526 Annu Rev Physiol 55:16-54
- 527 Plard F, Gaillard J-M, Coulson T, Hewison AJM, Delorme D, Warnant C, Bonenfant C
- 528 (2014) Mismatch Between Birth Date and Vegetation Phenology Slows the
- 529 Demography of Roe Deer. PLoS Biol 12:e1001828
- 530 Prates H, Bicca-Marques J (2008) Age-sex analysis of activity budget, diet, and positional
- behavior in Alouatta caraya in an orchard forest. Int J Primatol 29:703-715
- 532 R Core Team (2014) R: A Language and Environment for Statistical Computing. In. R
- 533 Foundation for Statistical Computing, Vienna, Austria

- Risenhoover KL (1986) Winter activity patterns of moose in interior Alaska. J Wildl Manag
 50:727-734
- Root TL, Price JT, Hall KR, Schneider SH, Rosenzweig C, Pounds JA (2003) Fingerprints of
 global warming on wild animals and plants. Nature 421:57-60
- Rughetti M, Festa-Bianchet M (2011) Seasonal changes in sexual size dimorphism in
 northern chamois. J Zool 284:257-264
- 540 Rutberg AT (1987) Adaptive hypotheses of birth synchrony in ruminants: an interspecific
 541 test. Am Nat 130:692-710
- 542 Rüttimann S, Giacometti M, McElligott A (2008) Effect of domestic sheep on chamois
- 543 activity, distribution and abundance on sub-alpine pastures. Eur J Wildl Res 54:110544 116
- 545 Schmid B, Helfrich-Forster C, Yoshii T (2011) A new ImageJ plug-in "ActogramJ" for
 546 chronobiological analyses. J Biol Rhythms 26:464-7
- 547 Schmitz OJ (1991) Thermal constraints and optimization of winter feeding and habitat choice
 548 in white-tailed deer. Holarct Ecol 14:104-111
- 549 Schröder W, von Elsner-Schack I (1985) Correct age determination in chamois. In: Lovari S
- (ed) The biology and management of mountain ungulates. Croom Helm, London, pp67–70
- 552 Shackleton DM (1997) Wild sheep and goats and their relatives: status survey and
- conservation action plan for Caprinae. World Conservation Union IUCN, Gland,
 Switzerland
- Signer C, Ruf T, Arnold W (2011) Hypometabolism and basking: the strategies of Alpine
 ibex to endure harsh over-wintering conditions. Funct Ecol 25:537-547
- 557 Spitzenberger F, Bauer K, Mayer A, Weiß E, Preleuthner M, Sackl P, Sieber J (2001) Die
- 558 Säugetierfauna Österreichs, Grüne Reihe des Bundesministeriums für Land-und
- 559 Forstwirtschaft, Umwelt und Wasserwirtschaft. Austria medien service, Graz

560	Symonds MRE, Moussalli A (2011) A brief guide to model selection, multimodel inference
561	and model averaging in behavioural ecology using Akaike's information criterion.
562	Behav Ecol Sociobiol 65:13-21
563	Tyler NJC, Blix AS (1990) Survival strategies in arctic ungulates. Rangifer Special Issue
564	3:211–230
565	Van Buskirk J (2012) Behavioural plasticity and environmental change. In: Candolin U,
566	Wong BBM (eds) Behavioural responses to a changing world. Oxford University
567	Press, Oxford (UK), pp 145-158
568	van Oort BEH, Tyler NJC, Gerkema MP, Folkow L, Blix AS, Stokkan K-A (2005) Circadian
569	organization in reindeer. Nature 438:1095-1096
570	VanSoest PJ (1994) Nutritional ecology of the ruminant, Second edn. Cornell University
571	Press, Ithaca, New York
572	Willisch CS, Bieri K, Struch M, Franceschina R, Schnidrig-Petrig R, Ingold P (2013)
573	Climate effects on demographic parameters in an unhunted population of Alpine
574	chamois (Rupicapra rupicapra). J Mammal 94:173-182
575	Willisch CS, Ingold P (2007) Feeding or resting? The strategy of rutting male Alpine
576	chamois. Ethology 113:97-104
577	Wood S (2006) Generalized Additive Models: An Introduction with R. Chapman and
578	Hall/CRC Boca Raton, FL
579	Zucker I (2001) Circannual rhythms. In: Takahashi JS, Turek FW, Moore RY (eds)
580	Handbook of behavioral neurobiology: circadian clocks. Kluwer Academic/Plenum
581	Publishers, New York, pp 509–527
582	Zuur A, Ieno E, Walker N, Saveliev A, Smith G (2009) Mixed effects models and extensions
583	in ecology with R. Springer, New York, USA
584	

585 Figure captions

587	Figure 1: Representative actograms of daily activity of one A) radio-collared female Alpine
588	chamois and B) one radio-collared male Alpine chamois in the Swiss National Park
589	(Switzerland). Vertical bars represent their activity levels (over intervals of 5 minutes), the
590	color of the bar being a function of activity level: from white (=0) to black for maximum
591	values (i.e., 255). Black vertical lines indicate dawn and dusk according to civil twilight.
592	Records are double plotted on a 48 hours time scale to help the interpretation.
593	
594	Figure 2: Periodogram analysis of locomotor activity rhythms of one radio-collared female
595	Alpine chamois (g500: A, B, C, D) and one radio-collared male Alpine chamois (g319: E, F,
596	G, H) in the Swiss National Park (Switzerland). Lomb-Scargle periodogram analysis
597	(confidence level, 95%) for actograms shown in Figure 1 performed over intervals of 10 days
598	in 4 different periods of the year. Periodogram analysis showed the presence and periodicity
599	of rhythms of locomotor activity. Periodicity confirmed the presence of unimodal (A, D, E,
600	H), bimodal (B,F) and trimodal (C,G) patterns in both the male and the female in the same
601	month of the year. The periodogram indicates the normalised power (PN) of the rhythm
602	explained by each period analysed within a range of 5–30 hours. The straight horizontal lines
603	represent the threshold of significance, set at $P=0.05$. Amplitudes of the peaks are used to
604	indicate the main period (dotted lines).
605	
606	Figure 3: Value predicted by the best Generalised Additive Mixed Model (see the text for
607	more details) of daily mean activity of Alpine chamois in the Swiss National Park
608	(Switzerland). The figure A) shows the effect of Julian day and B) daily mean temperature.
609	In the graphs males are represented by continuous line, females by broken line.

- 611 **Figure 4:** Value predicted by the best Generalised Additive Mixed Model (see the text for
- 612 more details) of Alpine chamois daily activity during winter months (1 October-30 April) in
- 613 the Swiss National Park (Switzerland). The figure A) shows the effect of radiation, B) snow
- 614 cover, and C) wind speed. In the graphs males are represented by continuous line, females by
- 615 broken line.
- 616

Tables

 Table 1: Top 5 Generalised Additive Mixed Models predicting daily mean activity in

 Alpine chamois throughout the year in the Swiss National Park, Switzerland.

model #	variables in the model	AIC	deltaAIC
FEMALES			
45	age class+ temperature+ precipitation+ wind speed+ Julian day	42086.07	0
38	age class+ temperature+ precipitation+ Julian day	42104.48	18.41
43	temperature+ precipitation+ wind speed+ Julian day	42141.61	55.54
31	temperature+ precipitation+ Julian day	42156.78	70.71
37	age class+ temperature+ wind speed+ Julian day	42172.75	86.68
MALES			
38	age class+ temperature+ precipitation+ Julian day	19479.10	0
45	age class+ temperature+ precipitation+ wind speed+ Julian day	19479.34	0.24
43	temperature+ precipitation+ wind speed+ Julian day	19486.58	7.48
31	temperature+ precipitation+ Julian day	19486.62	7.52
37	age class+ temperature+ wind speed+ Julian day	19522.41	43.31

Best models were selected with the $\Delta AIC \ge 2$ criterion. [AIC = Akaike information criterion; ΔAIC = difference in AIC value between the AIC of a given model and the best model (lowest AIC)].

Table 2: Top 5 Generalised Additive Mixed Models predicting daily mean activity in Alpine chamois during winter months (1 October-30 April) in the Swiss National Park, Switzerland.

2.37	0
2.37	0
) 52	0
	38.15
.61	39.24
3.83	76.46
).02	287.65
483	0
079	1.596
038	10.555
877	12.394
037	74.554
	.483 .079 .038 .877 .037

Best models were selected with the $\Delta AIC \ge 2$ criterion. [AIC = Akaike information criterion; ΔAIC = difference in AIC value between the AIC of a given model and the best model (lowest AIC)].







