

Effects of grape seed supplementation, alone or associated with linseed, on ruminal metabolism in Sarda dairy sheep

Questa è la versione Post print del seguente articolo:

Original

Effects of grape seed supplementation, alone or associated with linseed, on ruminal metabolism in Sarda dairy sheep / Correddu, F; Nudda, Anna; Battacone, Gianni; Boe, R; Francesconi, Ahd; Pulina, Giuseppe. - In: ANIMAL FEED SCIENCE AND TECHNOLOGY. - ISSN 0377-8401. - 199:(2015), pp. 61-72. [10.1016/j.anifeedsci.2014.11.002]

Availability:

This version is available at: 11388/46047 since: 2022-05-30T08:46:57Z

Publisher:

Published

DOI:10.1016/j.anifeedsci.2014.11.002

Terms of use:

Chiunque può accedere liberamente al full text dei lavori resi disponibili come "Open Access".

Publisher copyright

note finali coverpage

(Article begins on next page)

1 Effects of grape seed supplementation, alone or associated with linseed, on ruminal metabolism in
2 Sarda dairy sheep

3 F. Correddu, A. Nudda*, G. Battacone, R. Boe, A.H.D. Francesconi, G. Pulina

4 Dipartimento di Agraria, Sezione di Scienze Zootecniche, University of Sassari, Viale Italia 39,
5 07100, Sassari, Italy

6 * Corresponding author at: Dipartimento di Agraria, University of Sassari, Viale Italia 39, 07100,
7 Sassari, Italy. Tel.: +39 079229390. *E-mail address*: anudda@uniss.it

8 **ABSTRACT**

9 Grape seed is a by-product of the winery and distillery industry which could be used in animal
10 nutrition. To test the hypothesis that dietary supplementation with this by-product can decrease the
11 biohydrogenation (BH) of healthy fatty acids (FA), the present study evaluated the effects of grape
12 seed supplementation, alone or combined with linseed, on ruminal BH processes in dairy sheep. In
13 this 60-d trial, twenty-four lactating Sarda dairy ewes were assigned to 4 homogeneous groups and
14 fed as follows: (1) control diet (CON), (2) a diet supplemented with 300 g/d per head of grape seed
15 (GS), (3) a diet supplemented with 220 g/d per head of extruded linseed (LIN), (4) and a diet
16 supplemented with a mix of both grape seed and linseed (300 and 220 g/d per head, respectively)
17 (MIX). Ammonia, pH, volatile fatty acids (VFA) and FA composition were determined in rumen
18 liquor at 3 sampling dates (20, 40 and 60 d). Rumen pH was not influenced by diet ($P>0.05$). The
19 ammonia content was increased ($P<0.05$) in GS and MIX compared with LIN and CON. The molar
20 proportions of acetate and propionate and their ratio were not affected by the diet ($P>0.05$), whereas
21 the molar proportion of butyrate was the lowest in MIX. Rumenic acid (RA; CLA *cis*-9,*trans*-11)
22 concentration increased in GS compared with CON (0.78 vs. 0.45 mg/100 mg FA; $P<0.05$), whereas
23 the percentage of vaccenic acid (VA; C18:1 *trans*-11) tended to increase ($P<0.10$) in GS compared
24 with CON. The concentration of VA was higher in MIX than in CON (8.18 vs. 3.77 mg/100 mg FA;

25 P<0.05), whereas RA did not differ between the same groups. The concentration of linoleic acid
26 (LA; C18:2 *n*-6) decreased and stearic acid (SA; C18:0) increased in all supplemented groups,
27 whereas linolenic acid (LNA; C18:3 *n*-3) decreased in the two groups receiving grape seed
28 compared with CON and LIN. The concentration of total odd- and branched-chain fatty acids
29 (OBCFA) decreased in all supplemented groups compared with CON (P<0.05), evidencing that
30 grape seed and linseed supplementation influenced the ruminal BH processes. Grape seed was able
31 to increase the accumulation of RA when supplemented alone, and of VA when combined with
32 linseed; however, the rumen accumulation of SA in both groups supplemented with grape seed
33 evidenced that this by-product was not effective in decreasing the BH of dietary polyunsaturated
34 fatty acids (PUFA).

35 **Keywords:** Biohydrogenation, Fatty acid, Grape seed, Linseed, Rumen fermentation, Sheep.

36 **1. Introduction**

37 Grape seed is a by-product derived from the winery and distillery industries. In countries where
38 the wine industry is an important activity, the large production of by-products and wastes is a
39 serious problem, because of the high cost of their management and disposal. As many other agro-
40 industrial by-products, grape seeds are rich in polyphenols (Schieber et al., 2001), especially mono-,
41 oligo- and polymeric proanthocyanidins (Shrikhande, 2000), which are well known for their
42 antioxidant properties (Riceevans et al., 1995; Bagchi et al., 1997).

43 In the last few decades, a lot of attention has been directed to the content of healthy fatty acids
44 (FA), especially polyunsaturated fatty acids belonging to the family of *n*-3 (PUFA *n*-3), such as
45 alpha-linolenic acid (C18:3 *n*-3, LNA), and conjugated linoleic acid (CLA), such as rumenic acid
46 (CLA *cis*-9, *trans*-11, RA), in ruminant milk and dairy products. A lot of studies have demonstrated
47 that diet composition and rumen microbial biohydrogenation (BH) strongly influence the fatty acid
48 profile of milk and dairy products (Lourenço et al., 2010; Buccioni et al., 2012). Supplementation

49 with linseed, which is rich in LNA, is often used to improve the FA profile in sheep, goat and cow
50 milk and cheese, by increasing the concentration of healthy FA (Caroprese et al., 2010; Mughetti et
51 al., 2012; Nudda et al., 2013). Lipids from the diet are involved in a sequence of reactions,
52 performed by the rumen microbial population, including hydrolysis of esterified lipids to free FA,
53 whose double bonds can be partly isomerized and hydrogenated. These reactions can be explained
54 by a detoxification mechanism to defend microorganisms from the toxicity of unsaturated fatty
55 acids (UFA) as reported by Dehority (2003) and Maia et al. (2010). Because C18:2 *cis*-9, *cis*-12
56 (linoleic acid, LA, *n*-6 series) and LNA provided with the diet are greatly reduced (by 80% and
57 92%, respectively) in the rumen (Doreau and Ferlay, 1994), several strategies have been tested to
58 protect dietary FA from rumen BH. The encapsulation of lipids in a protein matrix is one of the
59 techniques proposed to protect FA (Tymchuk et al., 1998; Hawkins et al., 2013). However, the
60 occasional lack of efficiency of this method (Petit, 2003), its high costs and the need for
61 formaldehyde utilization limit its use. Another technique that has been studied is the decrease in BH
62 by chemical modifications of UFA, such as the transformation of UFA to fatty acyl amides
63 (Jenkins, 1998) or to calcium salts (Lundy Iii et al., 2004). The presence of some plant compounds,
64 such as polyphenols, in the diet of ruminants can also influence the BH process, by inhibiting the
65 activity of rumen microbes (Cabiddu et al., 2009; Vasta et al., 2009). The use of grape seed as a
66 supplement in ruminant nutrition could be an alternative for the expensive management and
67 disposal of this winery by-product. However, the literature available on the use of grape by-
68 products in ruminant nutrition is limited (Moate et al., 2014; Santos et al., 2014) and, to our
69 knowledge, the effects of the dietary supplementation with grape seed on the BH of PUFA in the
70 rumen has not been explored yet.

71 To test the hypothesis that grape seed supplementation decreases the BH activity of rumen
72 microbes, thus boosting the effect of linseed supplementation, this work investigated the effect of

73 grape seed supplementation, alone or associated with linseed, on rumen BH processes in dairy
74 sheep.

75

accepted

76 **2. Material and methods**

77 *2.1. Animals, experimental design and treatments*

78 Twenty four Sarda dairy ewes in the first part of lactation (<50 days in milk, DIM) were
79 assigned to 4 groups of 6 animals each, homogeneous for milk production, body weight, DIM, and
80 lactation order. Groups were confined in four boxes and randomly assigned to one of the 4
81 experimental diets (Table 1): control (CON) diet, a diet supplemented with 300 g/d per head of
82 grape seed (GS), a diet supplemented with 220 g/d per head of extruded linseed (LIN), and a diet
83 supplemented with a mix of 300 g/d per head of grape seed and 220 g/d per head of linseed (MIX).
84 The extruded linseed dose of 220 g/d was used to supply 70 g/d of fat per head. Considering that the
85 total phenolic content of grape seed was 333.3 ± 10.1 mg gallic acid equivalent (GAE)/100 g of dry
86 matter (DM; mean \pm S.E.), the grape seed dose of 300 g/d per head was used to provide
87 approximately 1 g/d per head of total polyphenols (approximately 0.4 g polyphenols/kg DM of
88 diet). The grape seed used was obtained from different red grape varieties after distillation in the
89 winemaking process. Grape seeds were ground before administration. The chemical composition
90 and FA profile of the grape seed and linseed are reported in Table 2.

91 All animals were offered the same basal ration consisting of beet pulp, a commercial
92 concentrate, dehydrated alfalfa hay and mixed hay. In addition, they received a mixed meal
93 composed of corn, soybean, pea, grape seed and linseed, in different proportions depending on the
94 dietary treatments, to obtain isoenergetic and isonitrogenous diets. Dehydrated alfalfa hay and
95 mixed hay were offered to each group of six ewes, whereas all other dietary ingredients were
96 provided to each animal by using individual feeders. The commercial concentrate (500 g/d per
97 head) was provided at the two daily milkings (7:30 and 17:30). The mixed meals were provided two
98 hours after each milking, and subsequently beet pulp (400 g/d per head) and dehydrated alfalfa hay
99 (on average 800 g/d per head) were provided. The mixed hay (on average 200 g/d per head) was
100 offered during the night. Both hays were offered at a fixed amount to avoid selection by the

101 animals. Clean water was always available. Diets were formulated to meet the sheep energy and
102 protein requirements using the Small Ruminant Nutrition Model (Tedeschi et al., 2010). Diets were
103 offered in a fixed amount to ensure constant daily intakes of dietary ingredients and to maintain
104 energy balance. The experiment lasted 10 weeks, with two weeks of adaptation period and 8 weeks
105 of data collection.

106 *2.2. Rumen sample collection*

107 Samples of rumen liquor were collected from all animals on days 20, 40 and 60 of the trial
108 (Samplings 1, 2 and 3), 2 hours after the morning feeding, using a stomach tube and an evacuation
109 pump. The collection of rumen liquor samples was performed by two teams of qualified experience
110 and required approximately 30 min in total. In order to reduce saliva contamination, the first portion
111 of the liquor collected (about 30 mL) was discarded. After sampling and filtering the rumen liquor,
112 the pH value was immediately measured by a pH meter (Orion 250A, Orion Research Inc., Boston,
113 MA, USA), equipped with a glass electrode with Polysolve reference electrolyte (model 238405,
114 Hamilton Company, Reno, NV, USA), and a thermometer. The sample of rumen liquor of each
115 animal was then divided into 3 subsamples, which were immediately stored at -80°C until analysis
116 for ammonia, volatile fatty acids (VFA) and FA.

117 *2.3. Chemical analysis*

118 Dry matter content of feed ingredients was determined by oven-drying at 105°C for 24 h. Neutral
119 detergent fiber (NDF) and acid detergent lignin (ADL) analyses were performed following the
120 method of Van Soest et al. (1991), using an Ankom 220 fiber analyzer (AnkomTM technology,
121 Fairport, NY, USA); NDF was measured using heat stable amylase and expressed exclusive of
122 residual ash (aNDFom) and ADL was determined by solubilization of cellulose with sulphuric acid.
123 Crude protein (CP) content was measured according to the Kjeldahl method (proc. 988.05; AOAC,
124 2000), extract ether (EE) by the Soxhlet method (proc. 920.39; AOAC, 2005) and ash by using a

125 muffle at 550°C (proc. 942.05; AOAC, 2000). Non-fiber carbohydrates (NFC) were calculated
126 according to Weiss (1999) as follows: $\text{NFC (g/kg DM)} = 100 - (\text{NDF} + \text{CP} + \text{ash} + \text{EE})$.

127 To determine the phenolic content of grape seed, the seeds were powdered by a blender and an
128 aliquot of 1.5 g was homogenized with 50 mL of a mixture of acetone/water (70/30, v/v) using an
129 Ultra Turrax homogenizer (Ultra Turrax T25, Janke&Kunkel KG, Germany) at 3000 rpm, for 2
130 min, in a water/ice bath. The homogenate was then centrifuged (6,000 x g, 15 min, 4°C) and the
131 supernatant was filtered through Whatman 541 filter paper (Whatman, Maidstone, England) for
132 determination of phenolic compounds, using the Folin-Ciocalteu method described by Kim et al.
133 (2003) with some modifications. Briefly, 0.5 mL of the extract were added to 1 mL of Folin-
134 Ciocalteu phenol reagent (Sigma Chem. Co., St. Louis, MO, USA) and 9.5 mL of distilled water in
135 a 25 mL volumetric flask and shaken. After 7 min, 10 mL of 0.71 M sodium carbonate (Na_2CO_3)
136 were added, and then the mixture was diluted to a volume of 25 mL with distilled water and mixed
137 thoroughly. The mixture was then stored in the dark for 120 min at room temperature. The
138 absorbance was read at 750 nm wavelength versus a blank solution. Total phenolic content,
139 expressed as mg of GAE/100 g DM, was determined following a calibration curve obtained using
140 gallic acid (Sigma Chem. Co., St. Louis, MO, USA) as standard.

141 The FA profiles of the whole diets, grape seed, linseed and rumen liquor were determined using
142 the method of Kramer et al. (1997) with some modifications. Samples kept at -80°C were
143 lyophilized and powdered. After adding 2 mL of sodium methoxide 0.5 M in methanol (Sigma and
144 Aldrich, Spain) to the power, it was placed in a water bath at 50°C for 10 min. After cooling to
145 room temperature, the samples were placed in a water/ice bath; then 3 mL of HCl/methanol (3M),
146 prepared with acetyl chloride and methanol, were added. Subsequently, the samples were heated
147 again in a water bath at 50°C for 10 min and cooled to room temperature; then 1 mL of a solution
148 containing methyl nonadecanoate (C19:0) as internal standard (Sigma Chemical Co., St. Louis,
149 MO, USA) and, subsequently, 7.5 mL of a 0.43 M solution of K_2CO_3 were added. After quick

150 agitation the samples were centrifuged (1,500 x g, room temperature, 5 min) and each supernatant
151 was kept in a vial for GC analysis. Fatty acid methyl esters (FAME) were determined using a Turbo
152 3400 CX gas chromatograph (Varian Inc., Palo Alto, CA, USA), equipped with a flame ionization
153 detector (FID) and an automatic injector 8200 (CX Varian Inc., Palo Alto, CA, USA). The column
154 (CP-select CB for FAME; 100 m x 0.32 mm i.d., 0.25 µm film thickness, Varian Inc., Palo Alto,
155 CA, USA) was operated with the following program: 75°C for 1 min, increased at 5°C/min to
156 148°C and at 8°C/min to 165°C, held for 35 min; then increased at 5.5°C/min to 210 and, finally, at
157 3°C to 230°C, held for 14 min. Helium (1 mL/min flow rate) was used as carrier gas with a pressure
158 of 37.000 psi. Split ratio was 1:100. The injector and detector temperatures were held at 225 and
159 285°C, respectively. Varian Star 3.4.1 software was used to compute the retention time and area of
160 each individual FAME, identified by comparing their retention times with those of known standards
161 and with published studies as detailed by Nudda et al. (2008).

162 Ammonia content in rumen liquor was determined by colorimetric method, according to Chaney
163 and Marbach (1962) with one modification (the use of salicylate instead of phenol), using a UV-
164 Visible Spectrophotometer (Varian, Inc., Palo Alto, CA, USA).

165 The VFA analysis was performed by a high-performance liquid chromatography (HPLC)
166 method. Briefly, a sample of approximately 2 mL was defrozen and centrifuged (15,000 x g, 10
167 min, 4°C); the supernatant was then withdrawn by syringe and injected into a HPLC system (Varian
168 Inc., Palo Alto, California, USA) after filtration (PTFE 0.45 µm, 13 mm). The HPLC was equipped
169 with an auto sampler (Varian 9300), a degasser (Varian 9012 Q), a UV detector (Varian 906P
170 Polychrom) and an Aminex HPX 87H column (Biorad Laboratories, Hercules, CA, USA). The
171 column was operated at 55°C with 0.008 N H₂SO₄ at 0.6 mL/min as eluent. Concentrations of VFA
172 were estimated by comparison with a calibration curve obtained by injecting 5 µl of 5 standard
173 solutions (5.6, 11.25, 22.5, 45 and 90 mmol/L of acetic acid, and 5, 10, 20, 40 and 80 mmol/L of
174 propionic and butyric acid) obtained by appropriate dilutions of a standard mixture of VFA

175 containing 5.40, 5.76 and 7.02 mg/mL of acetic, propionic and butyric acids, respectively, in H₂SO₄
176 0.1 N. The concentration of total and single VFA were expressed as mmol/L and mol/100 mol of
177 total VFA, respectively.

178 2.4. Statistical analysis

179 Data of pH, ammonia, VFA and FA profile were analyzed with the PROC MIXED procedure of
180 SAS (2002). The model included the diet treatment (CON, GS, LIN and MIX), sampling (3
181 samplings on days 20, 40 and 60 of the trial) and their interaction as fixed effects, and the animal
182 nested within the treatment as random effect. Means were separated using Tukey test (P<0.05).

183 3. Results

184 The predominant FA of the lipid fraction in grape seed was LA (74 g/100 g FA), followed by
185 oleic acid (OA; C18:1 *cis*-9) and palmitic acid (C16:0) (9.6 and 8.5 g/100 g FA, respectively). The
186 main FA in linseed was LNA, which accounted for 56 g/100 g FA (Table 2).

187 All dairy ewes consumed the whole daily amount of feeds supplied, as a consequence of the
188 fixed amount of diets provided (Table 3). The intake of most FA varied with diets
189 (CON<GS<LIN<MIX; Table 3) and reflected the EE concentration of the diets (Table 1).

190 3.1. pH and ammonia in rumen liquor

191 Rumen pH was not influenced by the diets, ranging from 6.68 in LIN to 6.77 in MIX (Table 4).
192 Sampling affected pH, which was lower on sampling 1 (6.58) than on sampling 2 (6.85), and
193 intermediate on sampling 3 (6.75). A diet × sampling interaction (P<0.01) occurred, but no
194 significant differences between diets within sampling were detected.

195 Rumen ammonia was affected by diet and sampling date (P<0.05, Table 4), being higher in the
196 GS and MIX groups than in the CON and LIN groups. On average, rumen ammonia increased

197 (P<0.05) from sampling 2 (14.71 mg/dL) to sampling 3 (17.71 mg/dL), with sampling 1 (15.67
198 mg/dL) not differing from them.

199 3.2. Rumen VFA

200 Concentration of total VFA, molar proportions of individual VFA and acetate:propionate ratio in
201 the rumen are shown in Table 4. The concentration of total VFA and the molar proportions of
202 acetate and propionate were not affected by diet, whereas the concentration of butyrate was reduced
203 (P<0.05) by the MIX containing both grape seed and linseed. A time effect was observed (P<0.01)
204 for total VFA and individual VFA, with the highest concentrations of propionate, acetate and
205 butyrate being on samplings 1, 2, and 3, respectively (P<0.05; Fig. 1). The acetate:propionate ratio
206 was not affected by diet, but differed (P<0.05) among sampling 1 (3.14), sampling 2 (4.04) and
207 sampling 3 (3.54). This ratio was influenced by group × sampling interaction (P<0.05), but no
208 significant differences between diets within sampling were detected.

209 3.3. FA composition in rumen liquor

210 The fatty acid profile in the rumen liquor collected from the ewes of the different experimental
211 treatments is given in Table 5. The total concentration of short-chain fatty acids (SCFA) was lower
212 (P<0.05) in all supplemented groups than in CON, mainly due to a decrease in C4:0 and C5:0
213 (P<0.05). The total concentration of medium-chain fatty acids (MCFA) was affected by diet
214 (P<0.01), with a significant reduction in both groups supplemented with linseed (LIN and MIX)
215 compared with CON, mainly due to the lower concentration (P<0.05) of C16:0 and some C16:1
216 isomers in the LIN and MIX groups. The concentration of MCFA and C16:0 in the GS group was
217 intermediate, not differing significantly from CON and LIN.

218 The concentration of total odd- and branched-chain fatty acids (OBCFA) decreased (P<0.05) in
219 all supplemented groups compared with CON. The MIX group had the lowest concentration of
220 OBCFA, even if not significantly different from the LIN group. The reduction in OBCFA caused by

221 grape seed was mainly due to the reduction ($P<0.05$) in *anteiso* branched-chain fatty acids (BCFA)
222 and odd-chain fatty acids (OCFA), especially *anteiso* C15:0 and C5:0. The reduction in OBCFA
223 caused by linseed was related not only to the reduction in these FA, but also to the decrease
224 ($P<0.05$) in *iso* BCFA, especially isomers of heptadecanoid acid (C17:0, *iso* C17:0 and *anteiso*
225 C17:0). The total concentration of long-chain fatty acids (LCFA) increased with the grape seed and
226 linseed supplementation, alone and in combination, being the highest in the MIX group ($P<0.05$).

227 The concentration of C18:0 (stearic acid, SA) was higher ($P<0.05$) in all the supplemented
228 groups than in CON and, on average, increased ($P<0.05$) in the last sampling. The diet \times sampling
229 interaction was also significant ($P<0.05$). The temporal evolution of this FA (Fig. 2) showed that
230 MIX was higher than CON throughout the study, whereas GS and LIN were higher than CON on
231 sampling 2 ($P<0.05$).

232 The concentration of LA decreased in all supplemented groups, being the lowest in LIN and
233 MIX ($P<0.05$). The concentration of LNA was higher in the LIN and CON groups than in the GS
234 and MIX groups ($P<0.05$). The concentrations of the geometrical isomers of LA, C18:2 *trans*-
235 9,*trans*-12 and C18:2 *trans*-9,*cis*-12, were higher in LIN and MIX than in GS and CON ($P<0.05$).

236 The concentration of RA was the highest ($P<0.05$) in the GS group. A significant effect of
237 sampling ($P<0.01$) occurred, with increasing levels of this FA over time (0.11, 0.55 and 0.97 g/100
238 g FA, respectively, on samplings 1, 2 and 3). A significant diet \times sampling interaction occurred,
239 with GS having a higher concentration of this FA than the other supplemented groups on sampling
240 3 (Fig. 3).

241 The sum of total *trans* fatty acids (TFA) was influenced ($P<0.01$) by diet, being higher in both
242 groups receiving linseed supplementation (LIN and MIX) than in the CON, with GS not differing
243 from CON and LIN. In particular, the concentration of C18:1 *trans*-11 (vaccenic acid, VA) was the
244 highest in MIX and the lowest in GS and CON, with LIN being intermediate ($P<0.05$).

245 Many of the FA measured during the trial were influenced by sampling, but the pattern varied
246 among them. In general, most of the SCFA (C4:0, C5:0 and C6:0) and isomers of C18:1 showed a
247 significant decrease ($P < 0.05$) in the last sampling, whereas most of the MCFA increased ($P < 0.01$)
248 over time. The individual LCFA showed a variable pattern, without a significant sampling effect on
249 total LCFA ($P > 0.05$). The PUFA decreased over time, mainly due to a decrease in PUFA n-6 (LA)
250 ($P < 0.05$). Among OBCFA, *iso* BCFA and *anteiso* BCFA increased after sampling 1, whereas
251 OCFA decreased over time on the last sampling ($P < 0.05$).

252 4. Discussion

253 During the experiment, the daily amount of feeds supplied individually including those objective
254 of the study, i.e. grape seed and linseed, were completely eaten by the animals of all groups. Group
255 feeding of the forages was chosen because the length of the experiment (10 weeks in total)
256 suggested to avoid the inevitable stress caused by individual confinement of lactating ewes (Hutson
257 et al., 2007). However the group intake of the forages was similar among the groups and mostly
258 complete, with only the less digestible stems refused, suggesting that the inclusion of grape seed,
259 linseed or both in the diet of lactating ewes did not negatively affect their DM intake.

260 The pH values measured in all experimental groups throughout the trial were between 6.5 and
261 7.0, which is a normal range for rumen liquor pH in sheep (5.5-7.0; Dziuk, 1984). This is in
262 agreement with previous studies showing that rumen pH was not markedly affected when sheep
263 were supplemented with grape pomace by-products, as a source of polyphenols (Yidiz et al., 2005;
264 Abarghuei et al., 2010), or with extruded linseed, as a source of PUFA (Mughetti et al., 2007). The
265 increase in rumen ammonia observed in the groups supplemented with grape seed, alone or in
266 combination with linseed, is in contrast with studies showing that the inclusion of polyphenols in
267 the diet was usually associated with a decrease in protein degradation (Abarghuei et al., 2010;
268 Dschaak et al., 2011), because of their ability to bind proteins and reduce the activity of microbial
269 enzymes by decreasing the growth of proteolytic bacteria (Molan et al., 2001). This discrepancy

270 could be explained by two hypotheses. The first is that the level of polyphenols in the grape seed
271 used in the present work was too low to influence the activity of some strains of proteolytic
272 bacteria, compared with the doses of polyphenols (values higher than 5 g/kg DM) which influenced
273 rumen bacteria population in other studies (Hervás et al., 2003; Vasta et al., 2010; Anantasook et
274 al., 2014). In addition, in some studies low concentrations of tannins increased the enzymatic
275 activity and growth of some bacteria *in vitro* (Jones et al., 1994) and *in vivo* (Vasta et al., 2010),
276 likely because the interaction between proteins and tannins can cause conformational changes in the
277 protein structure, giving more accessible sites for some proteolytic bacteria (Mole and Waterman,
278 1985; Molan et al., 2001). The second hypothesis is that the high concentration of ADL in the grape
279 seed used (410 g/kg DM) reduced the growth of cellulolytic bacteria, considering that high amounts
280 of lignin in ruminant diets can reduce fiber digestion (Hartley, 1972; Jung and Fahey, 1984) by
281 decreasing the growth of this type of bacteria (Akin, 1982), whose favorite substrate for protein
282 synthesis is ammonia (Brayant, 1973; Van Soest, 1994), with a consequent accumulation of
283 ammonia in the rumen.

284 The lack of effect of grape seed or extruded linseed, alone or in combination, on the
285 concentration of total VFA and on the proportion of acetate and propionate and their ratio in dairy
286 ewes observed in our study is in accordance with previous studies on dairy sheep fed diets rich in
287 LA and supplemented with 10 or 20 g tannins/kg DM from quebracho (Toral et al., 2011, 2013), on
288 cattle supplemented with quebracho polyphenols (Beauchemin et al., 2007) and on dairy cows
289 supplemented with flax seed (Neveu et al., 2013). Differently, Ivan et al. (2013) found a reduction
290 in acetate in dairy cows fed diets supplemented with oilseeds rich in LA or LNA and an increase in
291 propionate in cows fed the latter supplement.

292 The reduction in the proportion of butyrate observed only in the rumen liquor of the ewes fed the
293 MIX could be explained by the synergistic effect of the two sources of LA (grape seed) and LNA
294 (linseed) and by the higher value of fat in the MIX diet than in the GS and LIN diets. Our finding is

295 in accordance with the meta-analysis of Patra (2014) showing a negative effect of increasing levels
296 of fat in the diets of sheep on the proportion of butyrate, likely due to the inhibition of
297 microorganisms (protozoa and *Butyvirbio fibrisolvens*) involved in its production (Hristov et al.,
298 2009).

299 Overall, the observed variations in rumen FA profile between diets suggest shifts in rumen
300 microbial population. Considering that OBCFA derive largely from rumen microflora (Fievez et al.,
301 2012), the observed decrease in total OBCFA in all supplemented groups indicates that the diets
302 affected the activity and the growth of ruminal microorganisms. It is well documented that
303 variations in dietary treatments influence the synthesis of OBCFA, by affecting the relative
304 abundance of specific ruminal bacterial population (Vlaeminck et al., 2006). The reduction in total
305 OBCFA by linseed supplementation is in accordance with previous findings on lactating sheep
306 supplemented with sunflower oil (Toral et al., 2012), likely because these two supplements are a
307 source of PUFA, which have detrimental effects on the ruminal microflora (Maia et al., 2007,
308 2010). The decrease in OBCFA caused by grape seed supplementation could be partly explained by
309 the depressive effects of tannins on the microorganism growth (Baah et al., 2007; Vasta et al.,
310 2010), and mainly by the high concentration of LA in this by-product. The lowest value of OBCFA
311 and the highest value of LCFA in the MIX group suggest a combined effect of the two supplements,
312 likely because of the high LA (74% of total FA) in grape seed and high LNA (56% of total FA) in
313 linseed.

314 The OBCFA profile has been proposed as diagnostic tool to predict shifts in microbial
315 population associated with the diet variation (Lee et al., 1999; Vlaeminck et al., 2004). In the
316 present work the reduction in the *iso* C17:0 and *anteiso* C17:0 in the rumen liquor of sheep
317 receiving linseed supplementation could be related to a negative effect of this supplement on the
318 protozoa population, which produces a greater proportion of these FA than the bacteria population
319 in the rumen (Or-Rashid et al., 2007). In fact, several authors found a negative effect of lipid

320 addition on the growth of the protozoa population (Broudicou et al., 1994; Doreau and Ferlay,
321 1995; Ivan et al., 2013). The decrease in *anteiso* C15:0 in GS compared with CON suggests a
322 negative effect of grape seed supplementation on the rumen bacterial population, which is the main
323 producer of this FA in the rumen (Or-Rashid et al., 2007). The decrease in both *anteiso* C17:0 and
324 *anteiso* C15:0 in ewes fed the MIX suggests that the combination of grape seed and linseed
325 influenced both protozoa and bacteria in the rumen, as confirmed by the lowest OBCFA level of
326 this treatment.

327 The greater proportion of TFA in the groups fed linseed, alone or in the MIX, than in CON was
328 expected and was likely a consequence of the higher concentration of most C18:1 and C18:2
329 isomers, which derive in part from the BH of LNA. This finding is in agreement with previous
330 studies showing that the concentration of C18:1 and C18:2 isomers increased in the rumen of steers
331 fed linseed oil (Shingfield et al., 2011). The greater accumulation of most *trans* and *cis* isomers of
332 C18:1 in the MIX group could be mainly a consequence of the BH of OA, before its reduction to
333 SA. The increase in C18:2 *trans*-9,*trans*-12 and C18:2 *trans*-9,*cis*-12 observed only in the two diets
334 containing linseed is in agreement with an *in vitro* study showing that the concentrations of these
335 geometrical isomers of LA were increased by LNA supplementation, but not by LA
336 supplementation (Jouany et al., 2007).

337 The significantly higher values of VA in MIX, and numerically higher in LIN, compared with
338 GS and CON reflected the pattern of intake of its precursor LNA. The increase in RA with time,
339 although with a different pattern among diets, reflected the observed reduction in its precursor LA
340 during the trial. The highest concentration of RA in GS was likely associated mainly with the intake
341 of its precursor (LA) and, to a lesser extent, with the presence of polyphenols in grape seed by-
342 product. Some authors found that the accumulation of RA and VA in the rumen was increased by
343 tannins (Vasta et al., 2010; Buccioni et al., 2011), which can inhibit the last step of the BH process
344 of VA to SA (Khiaosa-Ard et al., 2009; Vasta et al., 2009; Rana et al., 2012). The low content of

345 polyphenols in the grape seeds used in this trial was probably not high enough to markedly
346 influence this BH process, as supported by the high rumen accumulation of SA and by the decrease
347 in LA and LNA in both groups supplemented with grape seed. The higher content of SA in rumen
348 liquor of all supplemented groups compared with CON could be also a consequence of the high
349 extent of OA biohydrogenation, as evidenced by the low accumulation of OA in the rumen liquor,
350 despite **its high intake**. As expected, the linseed supplementation did not influence the accumulation
351 of RA, because this FA is not an intermediate of the rumen BH of LNA (Wilde and Dawson, 1966).

352 **5. Conclusions**

353 The rumen metabolism of lactating dairy ewes was markedly influenced by dietary
354 supplementation with grape seed alone or mixed with linseed. Although grape seed caused a
355 decrease in LA and an increase in SA, this by-product determined an accumulation of RA in the
356 rumen. When grape seed was mixed with linseed, it enhanced the BH of LNA, thus promoting a
357 high accumulation of VA in the rumen. Therefore, the use of grape seed obtained after distillation in
358 the winemaking process seems not to be effective in enhancing the effect of linseed
359 supplementation.

360 **Acknowledgements**

361 The authors thank Antonio Fenu, Antonio Mazza, Roberto Rubattu and Gesumino Spanu, from
362 the Dipartimento di Agraria of the University of Sassari, for giving technical assistance, the
363 Azienda Ledda for making the farm and animals available, and Giovanni Pinna of Cargill s.r.l.
364 (Animal Nutrition Division) for providing the feed ingredients. Fabio Correddu gratefully
365 acknowledges the Sardinia Regional Government for the financial support of his Ph.D. scholarship
366 (P.O.R. Sardegna F.S.E. Operational Programme of the Autonomous Region of Sardinia, European
367 Social Fund 2007-2013 - Axis IV Human Resources, Objective 1.3, Line of Activity 1.3.1.).

368 **References**

369 Abarghuei, M.J., Rouzbehan, Y., Alipour, D., 2010. The influence of the grape pomace on the
370 ruminal parameters of sheep. *Livest. Sci.* 132, 73–79.

371 Akin, D.E., 1982. Forage cell wall degradation and p-coumaric, ferulic, and sinapic acids. *Agron. J.*
372 74, 424–428.

373 Anantasook, N., Wanapat, M., Cherdthong, A., 2014. Manipulation of ruminal fermentation and
374 methane production by supplementation of rain tree pod meal containing tannins and saponins
375 in growing dairy steers. *J. Anim. Physiol. Anim. Nutr.* 98, 50–55.

376 Association of Official Analytical Chemists, 2000. *Official Methods of Analysis*, 17th ed. AOAC
377 Int., Arlington, VA, USA.

378 Association of Official Analytical Chemists, 2005. *Official Methods of Analysis*. 18th ed. AOAC
379 Int., Gaithersburg, MD, USA.

380 Baah, J., Ivan, M., Hristov, A.N., Koenig, K.M., Rode, L.M., McAllister, T.A., 2007. Effects of
381 potential dietary antiprotozoal supplements on rumen fermentation and digestibility in heifers.
382 *Anim. Feed Sci. Technol.* 137, 126–137.

383 Bagchi, D., Garg, A., Krohn, R.L., Bagchi, M., Tran, M.X., Stohs, S.J., 1997. Oxygen free radical
384 scavenging abilities of vitamins C and E, and a grape seed proanthocyanidin extract in vitro.
385 *Res. Comm. Mol. Pathol. Pharmacol.* 95, 179–189.

386 Beauchemin, K.A., McGinn, S.M., Martinez, T.F., McAllister, T.A., 2007. Use of condensed tannin
387 extract from quebracho trees to reduce methane emissions from cattle. *J. Anim. Sci.* 85,
388 1990–1996.

389 Brayant, M.P., 1973. Nutritional requirements of the predominant rumen cellulolytic bacteria. *Fed.*
390 *Proc.* 32, 1809–1813.

391 Broudiscou, L., Pochet, S., Poncet, C., 1994. Effect of linseed oil supplementation on feed
392 degradation and microbial synthesis in the rumen of ciliate-free and refaunated sheep. *Anim.*
393 *Feed Sci. Technol.* 49, 189–202.

394 Buccioni, A., Decandia, M., Minieri, S., Molle, G., Cabiddu, A., 2012. Lipid metabolism in the
395 rumen: New insights on lipolysis and biohydrogenation with an emphasis on the role of
396 endogenous plant factors. *Anim. Feed Sci. Technol.* 174, 1–25.

397 Buccioni, A., Minieri, S., Rapaccini, S., Antongiovanni, M., Mele, M., 2011. Effect of chestnut and
398 quebracho tannins on FA profile in rumen liquid- and solid-associated bacteria: an in vitro
399 study. *Animal* 5, 1521–1530.

400 Cabiddu, A., Molle, G., Decandia, M., Spada, S., Fiori, M., Piredda, G., Addis, M., 2009.
401 Responses to condensed tannins of flowering sulla (*Hedysarum coronarium* L.) grazed by
402 dairy sheep: Part 2: Effects on milk fatty acid profile. *Livest. Sci.* 123, 230–240.

403 Caroprese, M., Marzano, A., Marino, R., Gliatta, G., Muscio, A., Sevi, A., 2010. Flaxseed
404 supplementation improves fatty acid profile of cow milk. *J. Dairy Sci.* 93, 2580–2588.

405 Chaney, A.L., Marbach, E.P., 1962. Modified reagents for determination of urea and ammonia.
406 *Clin. Chem.* 8, 130–132.

407 Dehority, B.A., 2003. *Rumen Microbiology*. Nottingham University Press, Nottingham, United
408 Kingdom.

409 Doreau, M., Ferlay, A., 1994. Digestion and utilisation of fatty acids by ruminants. *Anim. Feed Sci.*
410 *Technol.* 45, 379–396.

411 Doreau, M., Ferlay, A., 1995. Effect of dietary lipids on nitrogen metabolism in the lumen: a
412 review. *Livest. Prod. Sci.* 43, 97–110.

413 Dschaak, C.M., Williams, C.M., Holt, M.S., Eun, J.S., Young A.J., Min, B.R., 2011. Effects of
414 supplementing condensed tannin extract on intake, digestion, ruminal fermentation, and milk
415 production of lactating dairy cows. *J. Dairy Sci.* 94, 2508–2519.

416 Dziuk, H.E., 1984. Digestion in the ruminant stomach, in: Melvin J. Swenson (Ed.), *Dukes’*
417 *Physiology of Domestic Animals*, tenth ed. Cornell University Press, Ithaca, New York, pp.
418 331–332.

419 Fievez, V., Colman, E., Castro-Montoya, J.M., Stefanov, I., Vlaeminck, B., 2012. Milk odd- and
420 branched-chain fatty acids as biomarkers of rumen function-An update. *Anim. Feed Sci.*
421 *Technol.* 172, 51–65.

422 Folch, J., Lees, M., Sloane Stanley G.H., 1957. A simple method for the isolation and purification
423 of total lipids from animal tissues. *J. Biol. Chem.* 226, 497–509.

424 Hartley, R.D., 1972. p-Coumaric and ferulic acid components of cell walls of ryegrass and their
425 relationships with lignin and digestibility. *J. Sci. Food Agric.* 23, 1347–1354.

426 Hawkins, A., Yuan, K., Armendariz, C.K., Highland, G., Bello, N.M., Winowiski, T., Drouillard, J.
427 S., Titgemeyer, E. C., Bradford, B. J., 2013. Effects of urea formaldehyde condensation
428 polymer treatment of flaxseed on ruminal digestion and lactation in dairy cows. *J. Dairy Sci.*
429 96, 3907–3915.

430 Hervás, G., Frutos, P., Giráldez, F.J., Mantecón, Á.R., Álvarez del Pino, M.C., 2003. Effect of
431 different doses of quebracho tannins extract on rumen fermentation in ewes. *Anim. Feed Sci.*
432 *Technol.* 109, 65–78.

433 Hristov, A.N., Vander Pol, M., Agle, M., Zaman, S., Schneider, C., Ndegwa, P., Vaddella, V.K.,
434 Johnson, K., Shingfield K.J., Karnati, S.K.R., 2009. Effect of lauric acid and coconut oil on
435 ruminal fermentation, digestion, ammonia losses from manure, and milk fatty acid
436 composition in lactating cows. *J. Dairy Sci.* 92, 5561–5582.

437 Ivan, M., Petit, H.V., Chiquette, J., Wright, A.D.G., 2013. Rumen fermentation and microbial
438 population in lactating dairy cows receiving diets containing oilseeds rich in C-18 fatty acids.
439 *Br. J. Nutr.* 109, 1211–1218.

440 Jenkins, T.C., 1998. Fatty acid composition of milk from Holstein cows fed oleamide or canola oil.
441 *J. Dairy Sci.* 81, 794–800.

442 Jones, G.A., McAllister, T.A., Muir, A.D., Cheng, K.J., 1994. Effects of sainfoin (*Onobrychis*
443 *Viciifolia* Scop.) condensed tannins on growth and proteolysis by 4 strains of ruminal
444 bacteria. *Appl. Environ. Microbiol.* 60, 1374–1378.

445 Jouany, J.P., Lassalas, B., Doreau, M., Glasser, F., 2007. Dynamic features of the rumen
446 metabolism of linoleic acid, linolenic acid and linseed oil measured in vitro. *Lipids* 42,
447 351–360.

448 Jung, H.G., Fahey, Jr. G.C., 1984. Influence of phenolic acids on forage structural carbohydrate
449 digestion. *Can. J. Anim. Sci.* 64, 50–51.

450 Khiaosa-Ard, R., Bryner, S.F., Scheeder, M.R.L., Wettstein, H.R., Leiber, F., Kreuzer, M., Soliva
451 C.R., 2009. Evidence for the inhibition of the terminal step of ruminal alpha-linolenic acid
452 biohydrogenation by condensed tannins. *J. Dairy Sci.* 92, 177–188.

453 Kim, D.O., Chun, O.K., Kim, Y.J., Moon, H.Y., Lee, C.Y., 2003. Quantification of polyphenolics
454 and their antioxidant capacity in fresh plums. *J. Agr. Food Chem.* 51, 6509–6515.

455 Kramer, J.K.C., Fellner, V., Dugan, M.E.R., Sauer, F.D., Mossoba, M.M., Yurawecz, M.P., 1997.
456 Evaluating acid and base catalysts in the methylation of milk and rumen fatty acids with
457 special emphasis on conjugated dienes and total *trans* fatty acids. *Lipids* 32, 1219–1228.

458 Lee, S.S., Chang, M.B., Scollan, N.D., Merry, R.J., Dhanoa, M.S., Hobbs, P.J., Theobald, V.J.,
459 Maeng, W.J., Dewhurst, R.J., 1999. The fatty acid composition of solid- and liquid-associated
460 rumen bacteria isolated from cows. *Proc. Br. Soc. Anim. Sci.*, Scarborough, UK. *Br. Soc.*
461 *Anim. Sci.*, Edinburgh, UK, pp. 30.

462 Lourenço, M., Ramos-Morales, E., Wallace R.J., 2010. The role of microbes in rumen lipolysis and
463 biohydrogenation and their manipulation. *Animal* 4, 1008–1023.

464 Lundy Iii, F.P., Block, E., Bridges Jr, W.C., Bertrand, J. A., Jenkins, T.C., 2004. Ruminal
465 biohydrogenation in Holstein cows fed soybean fatty acids as amides or calcium salts. *J.*
466 *Dairy Sci.* 87, 1038–1046.

467 Maia, M.R., Chaudhary, L.C., Bestwick, C.S., Richardson, A.J., McKain, N., Larson, T.R.,
468 Graham, I.A., Wallace, R.J., 2010. Toxicity of unsaturated fatty acids to the biohydrogenating
469 ruminal bacterium, *Butyrivibrio fibrisolvens*. BMC Microbiol. 10, 52.

470 Maia, M.R., Chaudhary, L.C., Figueres, L., Wallace, R.J., 2007. Metabolism of polyunsaturated
471 fatty acids and their toxicity to the microflora of the rumen. Antonie van Leeuwenhoek 91,
472 303–314.

473 Moate, P.J., Williams, S.R.O., Torok, V.A., Hannah, M.C., Ribaux, B.E., Tavendale, M.H., Eckard,
474 R.J., Jacobs, J.L., Auldist, M.J., Wales, W.J., 2014. Grape marc reduces methane emissions
475 when fed to dairy cows. J. Dairy Sci. 97, 5073–5087.

476 Molan, A.L., Attwood, G.T., Min, B.R., McNabb, W.C., 2001. The effect of condensed tannins
477 from *Lotus pedunculatus* and *Lotus corniculatus* on the growth of proteolytic rumen bacteria
478 in vitro and their possible mode of action. Can. J. Microbiol. 47, 626–633.

479 Mole, S., Waterman, P.G., 1985. Stimulatory effects of tannins and cholic-acid on tryptic hydrolysis
480 of proteins - Ecological implications. J. Chem. Ecol. 11, 1323–1332.

481 Mughetti, L., Acuti, G., Antonini, C., De Vincenzi, S., Olivieri, O., Trabalza Marinucci, M., 2007.
482 Effects of feeding raw or extruded linseed on the ruminal ecosystem of sheep. Ital. J. Anim.
483 Sci. 6, 27–329.

484 Mughetti, L., Sinesio, F., Acuti, G., Antonini, C., Moneta, E., Peparaiò, M., Trabalza-Marinucci,
485 M., 2012. Integration of extruded linseed into dairy sheep diets: Effects on milk composition
486 and quality and sensorial properties of Pecorino cheese. Anim. Feed Sci. Technol. 178,
487 27–39.

488 Neveu, C., Baurhoo, B., Mustafa, A., 2013. Effect of feeding extruded linseed with different
489 forage:concentrate ratios on the performance of dairy cows. J. Dairy Sci. 96, 3886–3894.

490 Nudda, A., Battacone, G., Atzori, A.S., Dimauro, C., Rassu, S.P., Nicolussi, P., Bonelli, P., Pulina,
491 G., 2013. Effect of extruded linseed supplementation on blood metabolic profile and milk
492 performance of Saanen goats. *Animal* 7, 1464–1471.

493 Nudda, A., Palmquist, D.L., Battacone, G., Fancellu, S., Rassu, S.P.G., Pulina, G., 2008.
494 Relationships between the contents of vaccenic acid, CLA and n-3 fatty acids of goat milk
495 and the muscle of their suckling kids. *Livest. Sci.* 118, 195–203.

496 Or-Rashid, M.M., Odongo, N.E., McBride, B.W., 2007. Fatty acid composition of ruminal bacteria
497 and protozoa, with emphasis on conjugated linoleic acid, vaccenic acid, and odd-chain and
498 branched-chain fatty acids. *J. Anim. Sci.* 85, 1228–1234.

499 Patra, A.K., 2014. A meta-analysis of the effect of dietary fat on enteric methane production,
500 digestibility and rumen fermentation in sheep, and a comparison of these responses between
501 cattle and sheep. *Livest. Sci.* 162, 97–103.

502 Petit, H.V., 2003. Digestion, milk production, milk composition, and blood composition of dairy
503 cows fed formaldehyde treated flaxseed or sunflower seed. *J. Dairy Sci.* 86, 2637–2646.

504 Rana, M.S., Tyagi, A., Hossain, S.A., Tyagi, A.K., 2012. Effect of tanniniferous *Terminalia chebula*
505 extract on rumen biohydrogenation, $\Delta 9$ -desaturase activity, CLA content and fatty acid
506 composition in longissimus dorsi muscle of kids. *Meat Sci.* 90, 558–563.

507 Riceevans, C.A., Miller, N.J., Bolwell, G.P., Bramley, P.M., Pridham, J.B., 1995. The relative
508 antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radic. Res.* 22, 375–383.

509 Santos, N.W., Santos, G.T.D., Silva-Kazama, D.C., Grande, P.A., Pintro, P.M., de Marchi, F.E.,
510 Jobim, C.C., Petit, H.V., 2014. Production, composition and antioxidants in milk of dairy
511 cows fed diets containing soybean oil and grape residue silage. *Livest. Sci.* 159, 37–45.

512 SAS, 2002. User's Guide: statistics. SAS Institute Inc., Cary, NC, USA.

513 Schieber, A., Stintzing, F.C., Carle, R., 2001. By-products of plant food processing as a source of
514 functional compounds – recent developments. *Trends Food Sci. Tech.* 12, 401–413.

515 Shingfield, K.J., Lee, M.R.F., Humphries, D.J., Scollan, N.D., Toivonen, V., Beever, D.E.,
516 Reynolds, C.K., 2011. Effect of linseed oil and fish oil alone or as an equal mixture on
517 ruminal fatty acid metabolism in growing steers fed maize silage-based diets. *J. Anim. Sci.*
518 89, 3728–3741.

519 Shrikhande, A.J., 2000. Wine by-products with health benefits. *Food Res. Int.* 33, 469–474.

520 Tedeschi, L.O., Cannas, A., Fox, D.G., 2010. A nutrition mathematical model to account for dietary
521 supply and requirements of energy and other nutrients for domesticated small ruminants: The
522 development and evaluation of the Small Ruminant Nutrition System. *Small Ruminant Res.*
523 89, 174–184.

524 Toral, P.G., Belenguer, A., Shingfield, K.J., Hervás, G., Toivonen, V., Frutos, P., 2012. Fatty acid
525 composition and bacterial community changes in the rumen fluid of lactating sheep fed
526 sunflower oil plus incremental levels of marine algae. *J. Dairy Sci.* 95, 794–806.

527 Toral, P.G., Hervás, G., Belenguer, A., Bichi, E., Frutos, P., 2013. Effect of the inclusion of
528 quebracho tannins in a diet rich in linoleic acid on milk fatty acid composition in dairy ewes.
529 *J. Dairy Sci.* 96, 431–439.

530 Toral, P.G., Hervás, G., Bichi, E., Belenguer, A., Frutos, P., 2011. Tannins as feed additives to
531 modulate ruminal biohydrogenation: Effects on animal performance, milk fatty acid
532 composition and ruminal fermentation in dairy ewes fed a diet containing sunflower oil.
533 *Anim. Feed Sci. Technol.* 164, 199–206.

534 Tymchuk, S.M., Khorasani, G.R., Kennelly, J.J., 1998. Effect of feeding formaldehyde- and heat-
535 treated oil seed on milk yield and milk composition. *Can. J. Anim. Sci.* 78, 693–700.

536 Van Soest P.J., 1994. *Nutritional Ecology of the Ruminant*, second ed. Cornell University Press,
537 Ithaca, New York.

- 538 Van Soest, P.J., Robertson, J.B., Lewis, B.A., 1991. Methods for dietary fiber, neutral detergent
539 fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74,
540 3583–3597.
- 541 Vasta, V., Makkar, H.P.S., Mele, M., Priolo, A., 2009. Ruminal biohydrogenation as affected by
542 tannins in vitro. *Br. J. Nutr.* 102, 82–92.
- 543 Vasta, V., Yanez-Ruiz, D.R., Mele, M., Serra, A., Luciano, G., Lanza, M., Biondi, L., Priolo, A.,
544 2010. Bacterial and protozoal communities and fatty acid profile in the rumen of sheep fed a
545 diet containing added tannins. *Appl. Environ. Microbiol.* 76, 2549–2555.
- 546 Vlaeminck, B., Fievez, V., Cabrita, A.R.J., Fonseca, A.J.M., Dewhurst, R.J., 2006. Factors affecting
547 odd- and branched-chain fatty acids in milk: A review. *Anim. Feed Sci. Technol.* 131,
548 389–417.
- 549 Vlaeminck, B., Fievez, V., Van Laar, H., Demeyer, D., 2004. Rumen odd and branched chain fatty
550 acids in relation to in vitro rumen volatile fatty acid productions and dietary characteristics of
551 incubated substrates. *J. Anim. Physiol. A. Anim. Nutr.* 88, 401–411.
- 552 Weiss, W.P., 1999. Energy prediction equations for ruminant feeds. *Proceedings of the Cornell*
553 *Nutrition Conference for Feed Manufacturers*, Ithaca, NY, pp. 176–185.
- 554 Wilde, P.F., Dawson, R.M.C., 1966. Biohydrogenation of alpha-linolenic acid and oleic acid by
555 rumen micro-organism. *Biochem. J.* 98, 469–475.
- 556 Yidiz, S., Kaya, I., Unal, Y., Elmaji, D.A., Kayab, S., Cenesiz, M., Kayaa, M., Oncuer, A., 2005.
557 Digestion and body weight change in Tuj lambs receiving oak (*Quercus hartwissiana*) leaves
558 with and without PEG. *Anim. Feed Sci. Technol.* 122, 159–172.

559

560 Abbreviations:

561 ADL, acid detergent lignin;

562 BCFA, branched-chain fatty acids;

563 BH, biohydrogenation;
564 CLA, conjugated linoleic acid;
565 CON, control diet;
566 CP, crude protein;
567 DHA, docosahexaenoic acid;
568 DM, dry matter;
569 DPA, docosapentaenoic acid;
570 EE, ether extract;
571 EPA, eicosapentaenoic acid;
572 FA, fatty acids;
573 FAME, fatty acid methyl esters;
574 GS, diet supplemented with grape seed;
575 LA, linoleic acid;
576 LCFA, long-chain fatty acids;
577 LIN: diet supplemented with linseed;
578 LNA, linolenic acid;
579 MCFA, medium-chain fatty acids;
580 MIX: diet supplemented with both grape seed and linseed;
581 MUFA, monounsaturated fatty acids;
582 NDF, neutral detergent fiber;
583 NFC, non-fiber carbohydrates;
584 OA, oleic acid;
585 OBCFA, odd- and branched-chain fatty acids;
586 OCFA, odd-chain fatty acids;
587 PUFA, polyunsaturated fatty acids;
588 RA, rumenic acid;

589 SA, stearic acid;
590 SCFA, short-chain fatty acids;
591 SFA, saturated fatty acids;
592 TFA, *trans* fatty acids;
593 UFA, unsaturated fatty acids;
594 VA, vaccenic acid;
595 VFA, volatile fatty acids;
596

accepted

598 **Table 1**
599 Ingredients, chemical composition and fatty acid profile of diets

	Diets ^a			
	CON	GS	LIN	MIX
Ingredients (kg/day per head, as fed)				
Corn	0.15	0.17	-	-
Soybean	0.12	0.24	0.04	0.16
Pea	0.25	0.09	0.15	0.02
Grape seed	-	0.30	-	0.30
Linseed	-	-	0.22	0.22
Beet pulp	0.40	0.40	0.40	0.40
Commercial concentrate ^b	0.50	0.50	0.50	0.50
Dehydrated alfalfa hay ^c	0.80	0.80	0.80	0.80
Mixed hay ^d	0.20	0.20	0.20	0.20
Chemical composition ^e				
Dry matter (DM, g/kg)	908	916	912	920
NDF (g/kg DM)	418	428	437	445
NFC (g/kg DM)	334	289	285	242
ADL (g/kg DM)	46	89	50	94
CP (g/kg DM)	180	179	179	179
Ash (g/kg DM)	78	74	81	76
EE (g/kg DM)	20	32	51	58
Major fatty acids (g/100 g of total FAME) ^f				
C16:0	18.98	14.88	11.99	11.50
C16:1 <i>cis</i> -7	0.69	0.35	0.36	0.27
C16:1 <i>cis</i> -9	0.31	0.00	0.13	0.09
C16:1 <i>cis</i> -10	0.21	0.09	0.14	0.15
C17:0	0.00	0.00	0.15	0.12
C18:0 (SA)	3.33	4.47	4.39	4.68
C18:1 <i>cis</i> -9 (OA)	22.79	23.52	21.78	21.91
C18:1 <i>cis</i> -11	0.74	0.81	0.65	0.61
C18:2 <i>n</i> -6 (LA)	41.53	47.50	23.84	33.46
C18:3 <i>n</i> -3 (LNA)	8.25	5.04	34.45	24.93
C20:0	0.67	0.50	0.39	0.38
C20:1 <i>n</i> -9	0.38	0.36	0.24	0.27
C20:3 <i>n</i> -6	0.00	0.12	0.06	0.07
C24:0	0.53	0.45	0.37	0.30
SFA	24.24	20.88	17.75	17.42
MUFA	25.84	26.02	23.70	23.94

600 ^a CON: control diet, GS: diet supplemented with grape seed, LIN: diet supplemented with linseed, MIX: diet
601 supplemented with both grape seed and linseed.

602 ^b Commercial concentrate containing the following ingredients: sunflower seed flour, wheat bran, dehydrated
603 alfalfa meal, corn gluten, rice husk, corn flour, soybean hulls, sugar beet molasses, calcium carbonate from
604 powdered calcium rocks, distilled wheat, sodium chloride, plant oil (palm), mineral supplement (ferrous sulfate
605 monohydrate at 106 mg/kg, calcium diiodate at 1.7 mg/kg, manganese oxide at 90 mg/kg, sodium selenite at
606 0.46 mg/kg, zinc oxide at 87 mg/kg, and sodium molybdate at 2.5 mg/kg), antioxidant (E310 propyl gallate at
607 4.3 mg/kg) and vitamin supplement (vitamin A at 17,971 IU/kg, vitamin D3 at 3,494 IU/kg, and vitamin E at 60
608 mg/kg).

609 ^c Dehydrated alfalfa hay composition: DM 936 g/kg, NDF 387 g/kg DM, CP 196 g/kg DM, EE 44 g/kg DM.

610 ^d Mixed hay composition: DM 873 g/kg, NDF 551 g/kg DM, CP 116 g/kg DM, EE 11 g/kg DM.

611 ^e NDF: neutral detergent fiber measured using heat stable amylase and expressed exclusive of residual ash,
612 NFC: non-fiber carbohydrates, ADL: acid detergent lignin determined by solubilization of cellulose with
613 sulphuric acid, CP: crude protein, EE: ether extract.

614 ^f FAME: fatty acid methyl esters; SA: stearic acid; OA: oleic acid; LA: linoleic acid; LNA: linolenic acid; SFA:
615 saturated fatty acids, sum of the individual saturated fatty acids; MUFA: monounsaturated fatty acids, sum of
616 the individual monounsaturated fatty acids; PUFA: polyunsaturated fatty acids, sum of the individual
617 polyunsaturated fatty acids.

618

619

620

accepted

621 **Table 2**

622 Chemical composition and FA profile of grape seed and linseed

Chemical composition ^a	Linseed	Grape seed
Dry matter (DM, g/kg)	911	974
NDF (g/kg DM)	240	539
NFC (g/kg DM)	105	231
ADL (g/kg DM)	31	410
CP (g/kg DM)	264	93
Ash (g/kg DM)	44	27
EE (g/kg DM)	347	109
Fatty acid (g/100 g of FAME)		
C16:0	5.5	8.5
C18:0 (SA)	4.5	4.9
C18:1 <i>cis</i> -9 (OA)	18.0	9.6
C18:2 <i>n</i> -6 (LA)	15.0	74.0
C18:3 <i>n</i> -3 (LNA)	56.3	0.3

623 ^a NDF: neutral detergent fiber measured using heat stable amylase and expressed exclusive of residual ash,
624 NFC: non-fiber carbohydrates, ADL: acid detergent lignin determined by solubilization of cellulose with
625 sulphuric acid, CP: crude protein, EE: ether extract, FAME: fatty acid methyl esters, SA: stearic acid, OA: oleic
626 acid, LA: linoleic acid, LNA: linolenic acid.
627

628 **Table 3**

629 Dry matter and fatty acids intake of Sarda dairy ewes

	Diets ^a			
	CON	GS	LIN	MIX
Dry matter intake (kg/d)	2.2	2.5	2.1	2.4
Fatty acid intake (g/d) ^b				
C16:0	8.34	11.79	12.88	15.96
C16:1 <i>cis</i> -7	0.30	0.27	0.38	0.37
C16:1 <i>cis</i> -9	0.14	0.00	0.14	0.12
C16:1 <i>cis</i> -10	0.09	0.07	0.15	0.21
C17:0	0.00	0.00	0.16	0.17
C18:0 (SA)	1.46	3.54	4.72	6.49
C18:1 <i>cis</i> -9 (OA)	10.02	18.63	23.40	30.42
C18:1 <i>cis</i> -11	0.32	0.64	0.70	0.84
C18:2 <i>n</i> -6 (LA)	18.26	37.62	25.62	46.45
C18:3 <i>n</i> -3 (LNA)	3.63	3.99	37.02	34.60
C20:0	0.30	0.40	0.42	0.53
C20:1 <i>n</i> -9	0.17	0.28	0.26	0.37
C20:3 <i>n</i> -6	0.00	0.09	0.06	0.10
C24:0	0.23	0.36	0.40	0.41
SFA	10.66	16.53	19.08	24.17
MUFA	11.36	20.61	25.47	33.23
PUFA	21.94	42.06	62.91	81.39

630 ^a CON: control diet, GS: diet supplemented with grape seed, LIN: diet supplemented with linseed, MIX: diet
631 supplemented with both grape seed and linseed.

632 ^b SA: stearic acid; OA: oleic acid; LA: linoleic acid; LNA: linolenic acid; SFA: saturated fatty acids, sum of the
633 individual saturated fatty acids; MUFA: monounsaturated fatty acids, sum of the individual monounsaturated
634 fatty acids; PUFA: polyunsaturated fatty acids, sum of the individual polyunsaturated fatty acids.

635

636

637 **Table 4**

638 Effect of experimental diets with grape seed and linseed, alone or in combination, sampling and their
 639 interaction on rumen fermentation parameters in Sarda dairy ewes

	Diet ^a				SEM ^b	P-value ^c	
	CON	GS	LIN	MIX		D	S
Rumen pH	6.75	6.71	6.68	6.77	0.06	0.807	<0.001
Ammonia (mg/dL)	13.66 ^b	18.40 ^a	13.88 ^b	18.18 ^a	0.62	0.040	0.014
Total VFA ^d (mmol/L)	57.52	59.18	58.67	54.10	2.30	0.720	<0.001
VFA ^d (mol/100 mol)							
Acetate	67.45	69.22	67.44	71.04	0.54	0.149	<0.001
Propionate	20.84	18.61	21.03	19.60	0.43	0.386	<0.001
Butyrate	11.71 ^a	12.16 ^a	11.54 ^a	9.36 ^b	0.37	0.003	<0.001
Acetate:Propionate	3.44	3.79	3.31	3.74	0.09	0.422	<0.001

640 Means within a row with different superscripts (a, b) are different (P<0.05).

641 ^a CON: control diet, GS: diet supplemented with grape seed, LIN: diet supplemented with linseed, MIX: diet
 642 supplemented with both grape seed and linseed.

643 ^b SEM: standard error of the mean.

644 ^c D: effect of experimental diet; S: effect of sampling.

645 ^d VFA: volatile fatty acids.

646

Fatty acid (mg/100 mg of FAME) ^d	Diet ^a				SEM ^b	P-value ^c	
	CON	GS	LIN	MIX		D	S
C4:0	15.15 ^a	9.59 ^b	10.25 ^b	8.79 ^b	0.475	<0.001	<0.001
C5:0	3.01 ^a	1.77 ^b	1.86 ^b	1.73 ^b	0.113	0.002	<0.001
C6:0	0.66	0.58	0.41	0.54	0.036	0.398	<0.001
C8:0	0.01	0.01	0.01	0.01	0.0005	0.154	0.207
C10:0	0.02	0.02	0.02	0.01	0.001	0.466	0.034
C11:0	0.02	0.01	0.01	0.01	0.001	0.479	0.008
C12:0	0.34	0.28	0.32	0.25	0.016	0.399	<0.001
<i>iso</i> C13:0	0.02	0.02	0.02	0.02	0.001	0.998	0.009
<i>anteiso</i> C13:0	0.04	0.04	0.03	0.04	0.002	0.542	0.005
C13:0	0.06	0.08	0.09	0.09	0.007	0.413	0.121
<i>iso</i> C14:0	0.19	0.19	0.18	0.17	0.008	0.914	0.009
C14:0	0.69	0.77	0.78	0.64	0.032	0.504	<0.001
C14:1 <i>cis</i> -9	0.00	0.01	0.01	0.03	0.005	0.292	0.109
<i>iso</i> C15:0	0.33	0.30	0.27	0.32	0.013	0.437	<0.001
<i>anteiso</i> C15:0	1.40 ^a	1.12 ^b	1.19 ^{ab}	0.98 ^b	0.031	0.002	0.101
C15:0	1.21	1.16	1.10	1.05	0.029	0.352	<0.001
<i>iso</i> C16:0	2.18 ^a	2.00 ^a	1.67 ^{ab}	1.28 ^b	0.060	0.001	<0.001
C16:0	18.34 ^a	16.84 ^{ab}	15.68 ^{bc}	14.21 ^c	0.310	0.001	<0.001
C16:1 <i>trans</i> -8	0.05 ^a	0.02 ^b	0.02 ^b	0.02 ^b	0.003	0.001	0.012
C16:1 <i>trans</i> -9	0.21 ^a	0.12 ^b	0.11 ^b	0.01 ^b	0.010	<0.001	0.053
C16:1 <i>cis</i> -7	0.62	0.58	0.73	0.56	0.043	0.649	<0.001
C16:1 <i>cis</i> -9	0.06 ^a	0.05 ^{ab}	0.05 ^{ab}	0.03 ^b	0.002	0.005	0.227
<i>iso</i> C17:0	0.31 ^a	0.29 ^{ab}	0.21 ^b	0.23 ^{ab}	0.010	0.007	<0.001
<i>anteiso</i> C17:0	0.71 ^a	0.58 ^{ab}	0.51 ^b	0.47 ^b	0.023	0.009	<0.001
C17:0	0.53 ^a	0.44 ^{ab}	0.39 ^b	0.41 ^b	0.014	0.011	0.021
C18:0 (SA)	19.81 ^b	31.11 ^a	28.02 ^a	32.81 ^a	0.797	<0.001	0.003
C18:1 <i>trans</i> -4	0.03 ^c	0.04 ^{bc}	0.05 ^a	0.05 ^{ab}	0.002	<0.001	<0.001
C18:1 <i>trans</i> -6 + <i>trans</i> -8	0.40 ^b	0.58 ^{ab}	0.78 ^a	0.83 ^a	0.030	<0.001	<0.001
C18:1 <i>trans</i> -9	0.23 ^c	0.36 ^{bc}	0.45 ^{ab}	0.50 ^a	0.018	<0.001	<0.001
C18:1 <i>trans</i> -10	0.78	0.89	0.91	1.21	0.072	0.485	0.002
C18:1 <i>trans</i> -11 (VA)	3.77 ^b	4.76 ^b	6.44 ^{ab}	8.18 ^a	0.302	0.003	0.836
C18:1 <i>trans</i> -12	0.25 ^c	0.73 ^b	0.67 ^b	0.91 ^a	0.036	<0.001	0.098
C18:1 <i>cis</i> -9 (OA)	7.23 ^a	5.97 ^a	7.39 ^a	5.89 ^b	0.157	0.011	0.567
C18:1 <i>trans</i> -15 + <i>cis</i> -10	0.46 ^b	0.59 ^b	0.85 ^a	0.96 ^a	0.035	<0.001	<0.001
C18:1 <i>cis</i> -11	0.77 ^c	0.97 ^{bc}	1.21 ^{ab}	1.22 ^a	0.031	<0.001	0.311
C18:1 <i>cis</i> -12	0.33 ^c	0.61 ^{ab}	0.44 ^{bc}	0.71 ^a	0.028	<0.001	0.001

C18:1 <i>cis</i> -13	0.03	0.03	0.05	0.05	0.003	0.116	0.639
C18:1 <i>cis</i> -14 + <i>trans</i> -16	0.08 ^b	0.16 ^a	0.19 ^a	0.20 ^a	0.010	<0.001	0.006
C18:2 <i>trans</i> -9, <i>trans</i> -12	0.39 ^b	0.50 ^b	0.78 ^a	0.76 ^a	0.024	<0.001	0.007
C18:1 <i>cis</i> -15	0.09 ^c	0.21 ^{bc}	0.26 ^{ab}	0.35 ^a	0.016	<0.001	0.052
C18:2 <i>trans</i> -8, <i>cis</i> -13	0.23 ^a	0.20 ^{ab}	0.16 ^{bc}	0.14 ^c	0.007	0.001	0.170
C18:2 <i>trans</i> -11, <i>cis</i> -15	0.03	0.03	0.01	0.02	0.003	0.097	0.138
C18:2 <i>trans</i> -9, <i>cis</i> -12	0.04 ^b	0.04 ^b	0.20 ^a	0.21 ^a	0.013	<0.001	0.151
C18:2 <i>n</i> -6 (LA)	5.95 ^a	4.66 ^b	3.77 ^c	3.26 ^c	0.161	<0.001	<0.001
C18:2 <i>n</i> -4	0.16	0.14	0.17	0.11	0.012	0.434	<0.001
C18:3 <i>n</i> -6	0.29	0.26	0.25	0.24	0.016	0.718	<0.001
C18:3 <i>n</i> -3 (LNA)	1.60 ^a	0.81 ^b	1.29 ^a	0.84 ^b	0.050	<0.001	0.348
CLA <i>cis</i> -9, <i>trans</i> -11 (RA)	0.45 ^b	0.78 ^a	0.49 ^b	0.47 ^b	0.057	0.009	<0.001
C18:4 <i>n</i> -3	0.20 ^a	0.11 ^b	0.09 ^b	0.08 ^b	0.008	<0.001	0.067
CLA <i>trans</i> -9, <i>cis</i> -11 + C20	0.60	0.61	0.47	0.43	0.024	0.025	0.190
CLA <i>trans</i> -10, <i>cis</i> -12	0.12 ^{ab}	0.10 ^b	0.17 ^a	0.15 ^{ab}	0.007	0.007	<0.001
CLA <i>trans</i> -11, <i>cis</i> -13	0.03 ^b	0.08 ^{ab}	0.14 ^a	0.11 ^{ab}	0.011	0.012	<0.001
CLA <i>trans</i> -11, <i>trans</i> -13	0.06 ^b	0.07 ^b	0.12 ^a	0.12 ^a	0.005	<0.001	0.062
CLA <i>trans</i> -9,11+ C20:1 <i>n</i> -9	0.13	0.14	0.14	0.13	0.004	0.763	0.377
C20:2 <i>n</i> -6	0.02	0.02	0.02	0.02	0.001	0.418	<0.001
C20:3 <i>n</i> -9	0.03 ^{ab}	0.03 ^a	0.03 ^b	0.03 ^{ab}	0.001	0.027	0.323
C22:0	0.36 ^a	0.30 ^{ab}	0.33 ^{ab}	0.28 ^b	0.009	0.025	0.201
C22:1 <i>n</i> -11	0.06 ^a	0.04 ^b	0.04 ^b	0.03 ^b	0.002	0.003	<0.001
C20:5 <i>n</i> -3 (EPA)	0.20 ^a	0.15 ^{ab}	0.17 ^{ab}	0.13 ^b	0.007	0.011	<0.001
C22:2 <i>n</i> -6	0.17 ^a	0.13 ^b	0.13 ^b	0.11 ^b	0.005	0.001	<0.001
C24:0	0.42 ^a	0.34 ^b	0.37 ^{ab}	0.31 ^b	0.009	0.002	0.131
C24:1 <i>cis</i> -15	0.53 ^a	0.41 ^a	0.38 ^a	0.33 ^b	0.020	0.039	0.221
C22:5 <i>n</i> -3 (DPA)	0.01	0.01	0.01	0.01	0.002	0.456	0.909
C22:6 <i>n</i> -3 (DHA)	0.03	0.02	0.01	0.01	0.002	0.068	0.072
SCFA	18.86 ^a	11.97 ^b	12.56 ^b	11.10 ^b	0.583	<0.001	<0.001
MCFA	27.65 ^a	25.16 ^{ab}	23.72 ^{bc}	21.31 ^c	0.463	0.001	<0.001
LCFA	46.37 ^c	57.01 ^b	57.52 ^b	62.22 ^a	0.799	<0.001	0.229
OBCFA	10.01 ^a	8.00 ^b	7.54 ^{bc}	6.81 ^c	0.164	<0.001	0.029
<i>iso</i> BCFA	3.03 ^a	2.80 ^{ab}	2.36 ^{bc}	2.03 ^c	0.074	<0.001	<0.001
<i>anteiso</i> BCFA	2.15 ^a	1.73 ^b	1.73 ^b	1.49 ^b	0.043	<0.001	<0.001
OCFA	4.84 ^a	3.47 ^b	3.46 ^b	3.30 ^b	0.106	<0.001	<0.001
TFA	6.90 ^c	8.86 ^{bc}	11.53 ^{ab}	13.87 ^a	0.447	0.001	0.033
SFA	65.82	67.83	63.73	64.77	0.450	0.175	<0.001
MUFA	16.11 ^c	17.12 ^{bc}	21.18 ^{ab}	22.30 ^a	0.497	0.005	0.060
PUFA	10.73 ^a	8.94 ^b	8.64 ^b	7.39 ^c	0.171	<0.001	0.017
<i>n</i> -6	6.44 ^a	5.09 ^b	4.17 ^b	3.64 ^c	0.160	<0.001	<0.001

<i>n</i> -3	2.03 ^a	1.10 ^c	1.57 ^b	1.08 ^c	0.059	<0.001	0.054
CLA	1.38 ^b	1.79 ^a	1.52 ^{ab}	1.42 ^{ab}	0.067	0.045	<0.001

649 Means within a row with different superscripts (a, b, c) are different (P<0.05).


650 ^a CON: control diet, GS: diet supplemented with grape seed, LIN: diet supplemented with linseed, MIX: diet
651 supplemented with both grape seed and linseed.

652 ^b SEM: standard error of the mean.


653 ^c D: effect of experimental diet; S: effect of sampling.

654 ^d FAME: fatty acid methyl esters; SA: stearic acid; VA: vaccenic acid; OA: oleic acid; LA: linoleic acid; LNA:
655 linolenic acid; RA: rumenic acid; EPA: eicosapentaenoic acid; DPA: docosapentaenoic acid; DHA:
656 docosahexaenoic acid; SCFA: short-chain fatty acids, sum of the individual fatty acids from C4:0 to C10:0
657 reported in this table; MCFA: medium-chain fatty acids, sum of the individual fatty acids from C11:0 to C17:0
658 reported in this table; LCFA: long-chain fatty acids, sum of the individual fatty acids from C18:0 to DHA
659 reported in this table; OBCFA: odd- and branched-chain fatty acids, sum of *iso* BCFA, *anteiso* BCFA, OCFA; *iso*
660 BCFA: *iso* branched-chain fatty acids, sum of *iso* C13:0, *iso* C14:0, *iso* C15:0, *iso* C16:0, *iso* C17:0; *anteiso* BCFA:
661 sum of *anteiso* C13:0, *anteiso* C15:0, *anteiso* C17:0; OCFA: odd-chain fatty acids, sum of C5:0, C9:0, C11:0,
662 C13:0, C15:0, C17:0; TFA: *trans* fatty acids, sum of the individual *trans* fatty acids reported in this table; SFA:
663 saturated fatty acids, sum of the individual saturated fatty acids reported in this table; MUFA:
664 monounsaturated fatty acids, sum of the individual monounsaturated fatty acids reported in Table 4; PUFA:
665 polyunsaturated fatty acids, sum of the individual polyunsaturated fatty acids reported in Table 4; *n*-6: sum of
666 individual *n*-6 fatty acids reported in this table; *n*-3: sum of individual *n*-3 fatty acids reported in this table; CLA:
667 sum of individual conjugated of linoleic acids reported in this table.
668

accepted



670
 671 **Fig. 1.** Temporal evolution of the molar proportion of acetate, propionate and butyrate in rumen liquor
 672 on 3 samplings (days 20, 40 and 60) during the trial. Different letters (a, b, c) within the same volatile
 673 fatty acid show statistical differences ($P < 0.05$) between the samplings.
 674



675
 676 **Fig. 3.** Temporal evolution of CLA *cis-9,trans-11* (RA; rumenic acid) in rumen liquor on 3 samplings (days
 677 20, 40 and 60) during the trial in dairy ewes fed a control diet (CON), a diet supplemented with grape
 678 seed (GS), a diet supplemented with linseed (LIN) and a diet supplemented with both grape seed and
 679 linseed (MIX). Different letters (a, b) within the same sampling show statistical differences ($P < 0.05$)
 680 between the diets.
 681

682

683